


Fibonacci
(Leonardo Pisano)
1170-1240?
Statue in Pisa Italy



FIBONACCI NUMBERS
GOLDEN RATIO,
RECURRENCES

Lecture 23
CS2110 – Fall 2016

Prelim tonight

You know already whether you are taking it at 5:30 or 7:30.

5:30 Exam:
A thru Te... Take in Uris Hall G01
Th... thru Z. Take it in lves 305

7:30 Exam:
A... thru De... Take it in lves 305
Dh... Thru Z. Take it in Uris Hall G01

Fibonacci function

fib(0) = 0
fib(1) = 1
fib(n) = fib(n-1) + fib(n-2) for n ≥ 2

0, 1, 1, 2, 3, 5, 8, 13, 21, ...

In his book in 120
titled *Liber Abaci*

*Has nothing to do with the
famous pianist Liberaci*

But sequence described
much earlier in India:
Virahaṅka 600–800
Gopala before 1135
Hemacandra about 1150

The so-called Fibonacci
numbers in ancient and
medieval India.
Parmanad Singh, 1985
pdf on course website

Fibonacci function (year 1202)

fib(0) = 0
fib(1) = 1
fib(n) = fib(n-1) + fib(n-2) for n ≥ 2

```
/** Return fib(n). Precondition: n ≥ 0.*/
public static int f(int n) {
    if ( n <= 1) return n;
    return f(n-1) + f(n-2);
}
```

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55


We'll see that this is a
lousy way to compute
f(n)

Golden ratio $\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

Find the golden ratio when we divide a line into two parts such
that

whole length / long part = long part / short part

Call long part *a* and short part *b*




$(a + b) / a = a / b$ Solution is called Φ

See webpage:
<http://www.mathsisfun.com/numbers/golden-ratio.html>

Golden ratio $\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

Find the golden ratio when we divide a line into two parts *a* and
b such that

$(a + b) / a = a / b = \Phi$



Golden rectangle


See webpage:
<http://www.mathsisfun.com/numbers/golden-ratio.html>

Golden ratio $\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

Find the golden ratio when we divide a line into two parts a and b such that

$$(a + b) / a = a / b = \Phi$$

Golden rectangle



a/b
 $8/5 = 1.6$
 $13/8 = 1.625\dots$
 $21/13 = 1.615\dots$
 $34/21 = 1.619\dots$
 $55/34 = 1.617\dots$

For successive Fibonacci numbers a, b, a/b is close to Φ but not quite it Φ . 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, ...

Find fib(n) from fib(n-1)

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55

Since $\text{fib}(n) / \text{fib}(n-1)$ is close to the golden ratio,


You can see that $(\text{golden ratio}) * \text{fib}(n-1)$ is close to $\text{fib}(n)$

We can actually use this formula to calculate $\text{fib}(n)$ From $\text{fib}(n-1)$


Golden ratio and Fibonacci numbers: inextricably linked

Fibonacci function (year 1202)

Downloaded from wikipedia




Fibonacci tiling




Fibonacci spiral


0, 1, 1, 2, 3, 5, 8, 13, 21, 34 ...



The golden ratio



golden rectangle




How to draw a golden rectangle

fibonacci and bees

Male bee has only a mother
 Female bee has mother and father

The number of ancestors at any level is a Fibonacci number



1 MB

1 FB

2 FB

3 MB

5 FB

8 MB

MB: male bee, FB: female bee

Fibonacci in Pascal's Triangle

13

0
1
2
3
4
5
6
7
8

1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
1 6 15 20 15 6 1
1 7 21 35 35 21 7 1

$p[i][j]$ is the number of ways i elements can be chosen from a set of size j

Suppose you are a plant

14

You want to grow your leaves so that they all get a good amount of sunlight. You decide to grow them at successive angles of 180 degrees

Pretty stupid plant!
The two bottom leaves get VERY little sunlight!

Suppose you are a plant

15

You want to grow your leaves so that they all get a good amount of sunlight. 90 degrees, maybe?

Where does the fifth leaf go?

Fibonacci in nature

16

The artichoke uses the Fibonacci pattern to spiral the sprouts of its flowers.

$360/(\text{golden ratio}) = 222.492$

The artichoke sprouts its leaves at a constant amount of rotation: 222.5 degrees (in other words the distance between one leaf and the next is 222.5 degrees).

topones.weebly.com/1/post/2012/10/the-artichoke-and-fibonacci.html

Blooms: strobe-animated sculptures

17

www.instructables.com/id/Blooming-Zoetrope-Sculptures/

Uses of Fibonacci sequence in CS

18

- Fibonacci search
- Fibonacci heap data structure
- Fibonacci cubes: graphs used for interconnecting parallel and distributed systems

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

25

$T(0) = a$ $T(1) = a$ $T(n) = T(n-1) + T(n-2)$ $T(0) = a \leq a * 2^0$ $T(1) = a \leq a * 2^1$ $T(2) \leq a * 2^2$ $T(3) \leq a * 2^3$ $T(4) \leq a * 2^4$	$T(n) \leq c * 2^n$ for $n \geq N$ $T(k)$ = <Definition> $a + T(k-1) + T(k-2)$ <look to the left> $a + a * 2^{k-1} + a * 2^{k-2}$ = <arithmetic> $a * (1 + 2^{k-1} + 2^{k-2})$ <arithmetic> $a * 2^k$
---	--

Caching

26

As values of $f(n)$ are calculated, save them in an ArrayList. Call it a **cache**.

When asked to calculate $f(n)$ see if it is in the cache. If yes, just return the cached value. If no, calculate $f(n)$, add it to the cache, and return it.

Must be done in such a way that if $f(n)$ is about to be cached, $f(0), f(1), \dots, f(n-1)$ are already cached.


The golden ratio

27

$a > 0$ and $b > a > 0$ are in the **golden ratio** if

$(a + b) / b = b/a$ call that value ϕ

$\phi^2 = \phi + 1$ so $\phi = (1 + \text{sqrt}(5)) / 2 = 1.618 \dots$

a

 b

ratio of sum of sides to longer side

=

ratio of longer side to shorter side

Can prove that Fibonacci recurrence is $O(\phi^n)$

28

We won't prove it.
 Requires proof by induction
 Relies on identity $\phi^2 = \phi + 1$

Linear algorithm to calculate fib(n)

29

```

/** Return fib(n), for n >= 0. */
public static int f(int n) {
    if (n <= 1) return 1;
    int p= 0; int c= 1; int i= 2;
    // invariant: p = fib(i-2) and c = fib(i-1)
    while (i < n) {
        int fibi= c + p; p= c; c= fibi;
        i= i+1;
    }
    return c + p;
}
    
```

Logarithmic algorithm!

30

$$\begin{matrix} f_0 = 0 \\ f_1 = 1 \\ f_{n+2} = f_{n+1} + f_n \end{matrix} \quad \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} f_{n+1} \\ f_{n+2} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f_{n+1} \\ f_{n+2} \end{pmatrix} = \begin{pmatrix} f_{n+2} \\ f_{n+3} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} f_{n+k} \\ f_{n+k+1} \end{pmatrix}$$

Logarithmic algorithm!

31

$$\begin{aligned} f_0 &= 0 \\ f_1 &= 1 \\ f_{n+2} &= f_{n+1} + f_n \end{aligned} \quad \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} f_{n+k} \\ f_{n+k+1} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} = \begin{pmatrix} f_k \\ f_{k+1} \end{pmatrix}$$

You know a logarithmic algorithm for exponentiation—recursive and iterative versions

Gries and Levin
Computing a Fibonacci number in log time.
IPL 2 (October 1980), 68-69.

Another log algorithm!

32

$$\text{Define } \phi = (1 + \sqrt{5}) / 2 \quad \phi' = (1 - \sqrt{5}) / 2$$

The golden ratio again.

Prove by induction on n that

$$f_n = (\phi^n - \phi'^n) / \sqrt{5}$$