

Fibonacci
(Leonardo Pisano)
1170-1240?
Statue in Pisa Italy



FIBONACCI
NUMBERS AND
RECURRENCES

Lecture 26
CS2110 – Spring 2016

Info about optional final on course website

We post course grade as soon after 10 May as possible.
You answer quiz on CMS: Accept letter grade or take final?
Walk into final room? You must complete the final.
Take only 1 prelim? Must take final.
Final may lower (rarely) or raise course grade.
Conflict? Email Megan Gatch mlg34@cornell.edu
Quiet room / extra time. Email Megan Gatch
Review session 1: Java. TBA
Data structures, algorithms, concurrency. TBA

Fibonacci function

```
fib(0) = 0
fib(1) = 1
fib(n) = fib(n-1) + fib(n-2) for n ≥ 2
```

0, 1, 1, 2, 3, 5, 8, 13, 21, ...

In his book in 1202 titled *Liber Abaci*

Has nothing to do with the famous pianist Liberaci

But sequence described much earlier in India:
Virahaṅka 600–800
Gopala before 1135
Hemacandra about 1150

The so-called Fibonacci numbers in ancient and medieval India.
Parmanad Singh, 1985

Fibonacci function (year 1202)

```
fib(0) = 0
fib(1) = 1
fib(n) = fib(n-1) + fib(n-2) for n ≥ 2
```

*/** Return fib(n). Precondition: n ≥ 0.*/*

```
public static int f(int n) {
    if ( n <= 1) return n;
    return f(n-1) + f(n-2);
}
```

0, 1, 1, 2, 3, 5, 8, 13, 21, ...


LOUSY WAY TO COMPUTE: $O(2^n)$

```
/** Return fib(n). Precondition: n ≥ 0.*/
public static int f(int n) {
    if ( n <= 1) return n;
    return f(n-1) + f(n-2);
}
```

20
19 18
18 17 17 16
17 16 16 15 16 15 15 14

Fibonacci function (year 1202)

Downloaded from wikipedia



Fibonacci tiling Fibonacci spiral

0, 1, 1, 2, 3, 5, 8, 13, 21, ...

fibonacci and bees

Male bee has only a mother
Female bee has mother and father

The number of ancestors at any level is a Fibonacci number

```

graph TD
    L1[MB] --- L2[FB]
    L2 --- L3[FB]
    L2 --- L3[MB]
    L3 --- L4[FB]
    L3 --- L4[MB]
    L4 --- L5[FB]
    L4 --- L5[MB]
    L5 --- L6[FB]
    L5 --- L6[MB]
    L6 --- L7[FB]
    L6 --- L7[MB]
    L7 --- L8[FB]
    L7 --- L8[MB]
    
```

MB: male bee, FB: female bee

Fibonacci in nature

The artichoke uses the Fibonacci pattern to spiral the sprouts of its flowers.

The artichoke sprouts its leaves at a constant amount of rotation: 222.5 degrees (in other words the distance between one leaf and the next is 222.5 degrees). You can measure this rotation by dividing 360 degrees (a full spin) by the inverse of the **golden ratio**. We see later how **the golden ratio is connected to fibonacci**.

topones.weebly.com/1/post/2012/10/the-artichoke-and-fibonacci.html

Fibonacci in Pascal's Triangle

0

1

2

3

4

5

6

7

8

$p[i][j]$ is the number of ways i elements can be chosen from a set of size j

Blooms: strobe-animated sculptures

www.instructables.com/id/Blooming-Zoetrope-Sculptures/

Uses of Fibonacci sequence in CS

- Fibonacci search
- Fibonacci heap data structure
- Fibonacci cubes: graphs used for interconnecting parallel and distributed systems

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

Calculates $f(15)$ four times! What is complexity of $f(n)$?

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

13

$T(0) = a$ “Recurrence relation” for the time
 $T(1) = a$ It's just a recursive function
 $T(n) = a + T(n-1) + T(n-2)$

We can prove that $T(n)$ is $O(2^n)$

It's a “proof by induction”.
 Proof by induction is not covered in this course.
 But we can give you an idea about why $T(n)$ is $O(2^n)$

$T(n) \leq c \cdot 2^n$ for $n \geq N$

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

14

$T(0) = a$ $T(n) \leq c \cdot 2^n$ for $n \geq N$
 $T(1) = a$
 $T(n) = a + T(n-1) + T(n-2)$

$T(2)$
 = <Definition>
 $a + T(1) + T(0)$
 \leq <look to the left>
 $a + a \cdot 2^1 + a \cdot 2^0$
 = <arithmetic>
 $a \cdot (4)$
 = <arithmetic>
 $a \cdot 2^2$

$T(0) = a \leq a \cdot 2^0$
 $T(1) = a \leq a \cdot 2^1$

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

15

$T(0) = a$ $T(n) \leq c \cdot 2^n$ for $n \geq N$
 $T(1) = a$
 $T(n) = T(n-1) + T(n-2)$

$T(3)$
 = <Definition>
 $a + T(2) + T(1)$
 \leq <look to the left>
 $a + a \cdot 2^2 + a \cdot 2^1$
 = <arithmetic>
 $a \cdot (7)$
 \leq <arithmetic>
 $a \cdot 2^3$

$T(0) = a \leq a \cdot 2^0$
 $T(1) = a \leq a \cdot 2^1$
 $T(2) = a \leq a \cdot 2^2$

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

16

$T(0) = a$ $T(n) \leq c \cdot 2^n$ for $n \geq N$
 $T(1) = a$
 $T(n) = T(n-1) + T(n-2)$

$T(4)$
 = <Definition>
 $a + T(3) + T(2)$
 \leq <look to the left>
 $a + a \cdot 2^3 + a \cdot 2^2$
 = <arithmetic>
 $a \cdot (13)$
 \leq <arithmetic>
 $a \cdot 2^4$

$T(0) = a \leq a \cdot 2^0$
 $T(1) = a \leq a \cdot 2^1$
 $T(2) = a \leq a \cdot 2^2$
 $T(3) = a \leq a \cdot 2^3$

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

17

$T(0) = a$ $T(n) \leq c \cdot 2^n$ for $n \geq N$
 $T(1) = a$
 $T(n) = T(n-1) + T(n-2)$

$T(5)$
 = <Definition>
 $a + T(4) + T(3)$
 \leq <look to the left>
 $a + a \cdot 2^4 + a \cdot 2^3$
 = <arithmetic>
 $a \cdot (25)$
 \leq <arithmetic>
 $a \cdot 2^5$

$T(0) = a \leq a \cdot 2^0$
 $T(1) = a \leq a \cdot 2^1$
 $T(2) = a \leq a \cdot 2^2$
 $T(3) = a \leq a \cdot 2^3$
 $T(4) = a \leq a \cdot 2^4$

WE CAN GO ON FOREVER LIKE THIS

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

18

$T(0) = a$ $T(n) \leq c \cdot 2^n$ for $n \geq N$
 $T(1) = a$
 $T(n) = T(n-1) + T(n-2)$

$T(k)$
 = <Definition>
 $a + T(k-1) + T(k-2)$
 \leq <look to the left>
 $a + a \cdot 2^{k-1} + a \cdot 2^{k-2}$
 = <arithmetic>
 $a \cdot (1 + 2^{k-1} + 2^{k-2})$
 \leq <arithmetic>
 $a \cdot 2^k$

$T(0) = a \leq a \cdot 2^0$
 $T(1) = a \leq a \cdot 2^1$
 $T(2) = a \leq a \cdot 2^2$
 $T(3) = a \leq a \cdot 2^3$
 $T(4) = a \leq a \cdot 2^4$

Recursion for fib: $f(n) = f(n-1) + f(n-2)$

19

$T(0) = a$
 $T(1) = a$
 $T(n) = T(n-1) + T(n-2)$
 $T(0) = a \leq a * 2^0$
 $T(1) = a \leq a * 2^1$
 $T(2) = a \leq a * 2^2$
 $T(3) = a \leq a * 2^3$
 $T(4) = a \leq a * 2^4$
 $T(5) = a \leq a * 2^5$

$T(n) \leq c * 2^n$ for $n \geq N$

Need a formal proof, somewhere.
Uses mathematical induction

“Theorem”: all odd integers > 2 are prime
 3, 5, 7 are primes? yes
 9? experimental error
 11, 13? Yes.
 That’s enough checking

The golden ratio

20

$a > 0$ and $b > a > 0$ are in the **golden ratio** if

$(a + b) / a = a/b$ call that value φ

$\varphi^2 = \varphi + 1$ so $\varphi = (1 + \text{sqrt}(5)) / 2 = 1.618 \dots$

1.618....

ratio of sum of sides to longer side

=

ratio of longer side to shorter side

The golden ratio

21

golden rectangle

How to draw a golden rectangle

The Parthenon

22

Can prove that Fibonacci recurrence is $O(\varphi^n)$

23

We won't prove it.
 Requires proof by induction
 Relies on identity $\varphi^2 = \varphi + 1$

Linear algorithm to calculate fib(n)

24

```

/** Return fib(n), for n >= 0. */
public static int f(int n) {
    if (n <= 1) return 1;
    int p = 0; int c = 1; int i = 2;
    // invariant: p = fib(i-2) and c = fib(i-1)
    while (i < n) {
        int fibi = c + p; p = c; c = fibi;
        i = i + 1;
    }
    return c + p;
}
    
```

Logarithmic algorithm!

25

$$f_0 = 0$$

$$f_1 = 1$$

$$f_{n+2} = f_{n+1} + f_n$$

You know a logarithmic algorithm for exponentiation—recursive and iterative versions

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} f_{n+1} \\ f_{n+2} \end{pmatrix} \quad \text{so:} \quad \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^n \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} = \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix}$$

Gries and Levin/ Computing a Fibonacci number in log time. IPL 2 (October 1980), 68-69.

Constant-time algorithm!

26

$$\text{Define } \phi = (1 + \sqrt{5}) / 2 \quad \phi' = (1 - \sqrt{5}) / 2$$

The golden ratio again.

Prove by induction on n that

$$f_n = (\phi^n - \phi'^n) / \sqrt{5}$$

We went from $O(2^n)$ to $O(1)$