


CS4670/5760: Computer Vision

Kavita Bala

Lecture 11: Transformations




Announcements

- HW 1 FAQ pinned on Piazza
- PA 1 Artifacts
 - Soon voting will be set up. Please vote
- PA 2 out this weekend
 - Will present in class on Monday
- Next Friday: in class review
- Prelim: Mar 5, Thu, 7:30pm
 - Call Auditorium, Kennedy Hall

Reading

- Szeliski: Chapter 6.1, 3.6

What is the geometric relationship between these two images?



Answer: Similarity transformation (translation, rotation, uniform scale)

All 2D Linear Transformations

- Linear transformations are combinations of ...

- Scale,
- Rotation,
- Shear, and
- Mirror

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Properties of linear transformations:

- Origin maps to origin
- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} \begin{bmatrix} i & j \\ k & l \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Homogeneous coordinates

Trick: add one more coordinate:

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image
coordinates



Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

Translation

- Solution: homogeneous coordinates to the rescue

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$

Affine transformations

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$



any transformation with
last row [0 0 1] we call an
affine transformation

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$

Basic affine transformations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Translate

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Scale

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D *in-plane* rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & sh_x & 0 \\ sh_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Shear


Affine Transformations

- Affine transformations are combinations of ...

- Linear transformations, and
- Translations

$$\begin{bmatrix} x' \\ y' \\ w \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

- Properties of affine transformations:
 - Origin does not necessarily map to origin
 - Lines map to lines
 - Parallel lines remain parallel
 - Ratios are preserved
 - Closed under composition



- Euclidean: translation, rotation, reflection
- Similarity: translation, rotation, uniform scale, reflection
- Affine: linear transformations + translation







| Transformation | Matrix | # DoF | Preserves | Icon |
|-----------------------|---|--------------|------------------|---|
| translation | $\begin{bmatrix} \mathbf{I} & & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 2 | orientation |  |
| rigid (Euclidean) | $\begin{bmatrix} \mathbf{R} & & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 3 | lengths |  |
| similarity | $\begin{bmatrix} s\mathbf{R} & & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 4 | angles |  |
| affine | $\begin{bmatrix} \mathbf{A} \end{bmatrix}_{2 \times 3}$ | 6 | parallelism |  |
| projective | $\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}_{3 \times 3}$ | 8 | straight lines |  |

Table 2.1 Hierarchy of 2D coordinate transformations. Each transformation also preserves the properties listed in the rows below it, i.e., similarity preserves not only angles but also parallelism and straight lines. The 2×3 matrices are extended with a third $[\mathbf{0}^T \ 1]$ row to form a full 3×3 matrix for homogeneous coordinate transformations.

2D image transformations




| Name | Matrix | # D.O.F. | Preserves: | Icon |
|-------------------|---|----------|-------------------|------|
| translation | $\begin{bmatrix} \mathbf{I} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 2 | orientation + ... | |
| rigid (Euclidean) | $\begin{bmatrix} \mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 3 | lengths + ... | |
| similarity | $\begin{bmatrix} s\mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 4 | angles + ... | |
| affine | $\begin{bmatrix} \mathbf{A} \end{bmatrix}_{2 \times 3}$ | 6 | parallelism + ... | |
| projective | $\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}_{3 \times 3}$ | 8 | straight lines | |

These transformations are a nested set of groups

- Closed under composition and inverse is a member


Homographies



Reading


- Szeliski: Chapter 3.6

Is this an affine transformation?



Where do we go from here?

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$




what happens when we
mess with this row?

affine transformation

Projective Transformations aka Homographies aka Planar Perspective Maps


$$\mathbf{H} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix}$$

Called a *homography*
(or *planar perspective map*)



Why do we care?

- What is the relation between a plane in the world and a perspective image of it?
- Can we reconstruct another view from one image?
- Relation between pairs of images
 - Need to make a mosaic




Homographies

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by + c \\ dx + ey + f \\ gx + hy + 1 \end{bmatrix}$$


Recap of 4620

Plane projection in drawing



[CS 417 Spring 2002]

Perspective projection



similar triangles: $(x', y', w') = (fx, fy, z)$

$$x' = fx/z, y' = fy/z$$

$$M_{proj} = \begin{pmatrix} f & 0 & 0 \\ 0 & f & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Homographies

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by + c \\ dx + ey + f \\ gx + hy + 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} \frac{ax+by+c}{gx+hy+1} \\ \frac{dx+ey+f}{gx+hy+1} \\ 1 \end{bmatrix}$$

Homographies


$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

What happens when
the denominator is 0?

\sim

$$\begin{bmatrix} \frac{ax+by+c}{gx+hy+1} \\ \frac{dx+ey+f}{gx+hy+1} \\ 1 \end{bmatrix}$$

Points at infinity







Implications of w

$$\begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} \sim \begin{bmatrix} wx \\ wy \\ wz \\ w \end{bmatrix}$$

- All scalar multiples of a 4-vector are equivalent
- When w is not zero, can divide by w
 - therefore these points represent “normal” affine points
- When w is zero, it’s a point at infinity, a.k.a. a direction
 - this is the point where parallel lines intersect
 - can also think of it as the vanishing point

Masaccio, Trinity, Florence





Consider a line through point (a, b) with slope m

$x' = a + m t$ (line through a along direction m)

if $t = 1/w$; $x' = (a w + m)/w$

In homogeneous coordinates $(aw+m, w)$

At $w = 0$, point at infinity $(m, 0)$ represents line with slope m






Image warping with homographies




Homographies








Homographies

- Homographies ...
 - Affine transformations, and
 - Projective warps
- $$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$
- Properties of projective transformations:
 - Origin does not necessarily map to origin
 - Lines map to lines
 - Parallel lines do not necessarily remain parallel
 - Ratios are not preserved
 - Closed under composition

2D image transformations



| Name | Matrix | # D.O.F. | Preserves: | Icon |
|-------------------|---|----------|-------------------|---|
| translation | $\begin{bmatrix} \mathbf{I} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 2 | orientation + ... |  |
| rigid (Euclidean) | $\begin{bmatrix} \mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 3 | lengths + ... |  |
| similarity | $\begin{bmatrix} s\mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$ | 4 | angles + ... |  |
| affine | $\begin{bmatrix} \mathbf{A} \end{bmatrix}_{2 \times 3}$ | 6 | parallelism + ... |  |
| projective | $\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}_{3 \times 3}$ | 8 | straight lines |  |

These transformations are a nested set of groups

- Closed under composition and inverse is a member






Image Warping

- Given a coordinate xform $(x',y') = T(x,y)$ and a source image $f(x,y)$, how do we compute an xformed image $g(x',y') = f(T(x,y))$?



Forward Warping

- Send each pixel $f(\mathbf{x})$ to its corresponding location $(\mathbf{x}', \mathbf{y}') = T(\mathbf{x}, \mathbf{y})$ in $g(\mathbf{x}', \mathbf{y}')$



procedure *forwardWarp*($f, h, \text{out } g$):


For every pixel x in $f(x)$

1. Compute the destination location $x' = h(x)$.
2. Copy the pixel $f(x)$ to $g(x')$.

Algorithm 3.1 Forward warping algorithm for transforming an image $f(x)$ into an image $g(x')$ through the parametric transform $x' = h(x)$.


Forward Warping

- Send each pixel $f(x,y)$ to its corresponding location $x' = h(x,y)$ in $g(x',y')$
 - What if pixel lands “between” pixels?
 - Answer: add “contribution” to several pixels, normalize later (*splatting*)
 - Problem? Can still result in holes




Inverse Warping

- Get each pixel $g(x',y')$ from its corresponding location $(x,y) = T^{-1}(x',y')$ in $f(x,y)$
- Requires taking the inverse of the transform
- What if pixel comes from “between” pixels?



Inverse Warping

- Get each pixel $g(\mathbf{x}')$ from its corresponding location $\mathbf{x}' = \mathbf{h}(\mathbf{x})$ in $f(\mathbf{x})$
- What if pixel comes from “between” two pixels?
- Answer: *resample* color value from *interpolated (prefiltered)* source image



procedure *inverseWarp*($f, h, \text{out } g$):

For every pixel x' in $g(x')$

1. Compute the source location $x = \hat{h}(x')$
2. Resample $f(x)$ at location x and copy to $g(x')$

Algorithm 3.2 Inverse warping algorithm for creating an image $g(x')$ from an image $f(x)$ using the parametric transform $x' = h(x)$.

Interpolation

- Possible interpolation filters:
 - nearest neighbor
 - bilinear
 - bicubic (interpolating)
 - sinc
- Needed to prevent “jaggies”
and “texture crawl”

(with prefiltering)

