



Perspective projection and Transformations

The pinhole camera



The pinhole camera



$$\lambda = 0 \Rightarrow Q(\lambda) = O$$

$$\lambda = 1 \Rightarrow Q(\lambda) = P$$

$$Q(\lambda)$$


$$= (0 + \lambda(X - 0), 0 + \lambda(Y - 0), 0 + \lambda(Z - 0))$$

$$= (\lambda X, \lambda Y, \lambda Z)$$

The pinhole camera

- Pinhole camera collapses *ray OP* to point *p*
- Any point on ray $OP = O + \lambda(P - O) = (\lambda X, \lambda Y, \lambda Z)$
- For this point to lie on $Z=-1$ plane:
$$\lambda^* Z = -1$$
$$\Rightarrow \lambda^* = \frac{-1}{Z}$$
- Coordinates of point *p*:

$$(\lambda^* X, \lambda^* Y, \lambda^* Z) = \left(\frac{-X}{Z}, \frac{-Y}{Z}, -1 \right)$$



The projection equation

- A point $P = (X, Y, Z)$ in 3D projects to a point $p = (x, y)$ in the image

$$x = \frac{-X}{Z}$$


$$y = \frac{-Y}{Z}$$

- But pinhole camera's image is inverted, invert it back!

$$x = \frac{X}{Z}$$


$$y = \frac{Y}{Z}$$

Another derivation



A virtual image plane

- A pinhole camera produces an inverted image
- Imagine a "virtual image plane" in the front of the camera



The projection equation

$$x = \frac{X}{Z}$$
$$y = \frac{Y}{Z}$$

Consequence 1: Farther away objects are smaller



Image of foot: $(\frac{X}{Z}, \frac{Y}{Z})$

Image of head: $(\frac{X}{Z}, \frac{Y + h}{Z})$

$$\frac{Y + h}{Z} - \frac{Y}{Z} = \frac{h}{Z}$$

Consequence 2: Parallel lines converge at a point

- Point on a line passing through point A with direction D :

$$Q(\lambda) = A + \lambda D$$

- Parallel lines have the same direction but pass through different points

$$Q(\lambda) = A + \lambda D$$

$$R(\lambda) = B + \lambda D$$

-



Consequence 2: Parallel lines converge at a point

- Parallel lines have the same direction but pass through different points

$$Q(\lambda) = A + \lambda D$$

$$R(\lambda) = B + \lambda D$$

- $A = (A_X, A_Y, A_Z)$
- $B = (B_X, B_Y, B_Z)$
- $D = (D_X, D_Y, D_Z)$



Consequence 2: Parallel lines converge at a point

- $Q(\lambda) = (A_X + \lambda D_X, A_Y + \lambda D_Y, A_Z + \lambda D_Z)$
- $R(\lambda) = (B_X + \lambda D_X, B_Y + \lambda D_Y, B_Z + \lambda D_Z)$
- $q(\lambda) = \left(\frac{A_X + \lambda D_X}{A_Z + \lambda D_Z}, \frac{A_Y + \lambda D_Y}{A_Z + \lambda D_Z} \right)$
- $r(\lambda) = \left(\frac{B_X + \lambda D_X}{B_Z + \lambda D_Z}, \frac{B_Y + \lambda D_Y}{B_Z + \lambda D_Z} \right)$
- Need to look at these points as Z goes to infinity
- Same as $\lambda \rightarrow \infty$



Consequence 2: Parallel lines converge at a point

- $q(\lambda) = \left(\frac{A_X + \lambda D_X}{A_Z + \lambda D_Z}, \frac{A_Y + \lambda D_Y}{A_Z + \lambda D_Z} \right)$
- $r(\lambda) = \left(\frac{B_X + \lambda D_X}{B_Z + \lambda D_Z}, \frac{B_Y + \lambda D_Y}{B_Z + \lambda D_Z} \right)$

$$\lim_{\lambda \rightarrow \infty} \frac{A_X + \lambda D_X}{A_Z + \lambda D_Z} = \lim_{\lambda \rightarrow \infty} \frac{\frac{A_X}{\lambda} + D_X}{\frac{A_Z}{\lambda} + D_Z} = \frac{D_X}{D_Z}$$

$$\lim_{\lambda \rightarrow \infty} q(\lambda) = \left(\frac{D_X}{D_Z}, \frac{D_Y}{D_Z} \right)$$

$$\lim_{\lambda \rightarrow \infty} r(\lambda) = \left(\frac{D_X}{D_Z}, \frac{D_Y}{D_Z} \right)$$

Consequence 2: Parallel lines converge at a point

- Parallel lines have the same direction but pass through different points

$$Q(\lambda) = A + \lambda D$$

$$R(\lambda) = B + \lambda D$$

- Parallel lines converge at the same point $\left(\frac{D_X}{D_Z}, \frac{D_Y}{D_Z}\right)$
- This point of convergence is called the *vanishing point*
- What happens if $D_Z = 0$?

Consequence 2: Parallel lines converge at a point



What about planes?



$$N_X X + N_Y Y + N_Z Z = d$$

$$\Rightarrow N_X \frac{X}{Z} + N_Y \frac{Y}{Z} + N_Z = \frac{d}{Z}$$


$$\Rightarrow N_X x + N_Y y + N_Z = \frac{d}{Z}$$

Take the limit as Z approaches infinity



$$N_X x + N_Y y + N_Z = 0$$

Vanishing line of
a plane



Changing coordinate systems





Changing coordinate systems





Changing coordinate systems





Changing coordinate systems



Changing coordinate systems



Changing coordinate systems



Rotations and translations

- How do you represent a rotation?
- A point in 3D: (X,Y,Z)
- Rotations can be represented as a matrix multiplication

$$\mathbf{v}' = R\mathbf{v}$$

- What are the properties of rotation matrices?

Properties of rotation matrices

- Rotation does not change the length of vectors

$$\mathbf{v}' = R\mathbf{v}$$

$$\|\mathbf{v}'\|^2 = \mathbf{v}'^T \mathbf{v}'$$

$$= \mathbf{v}^T R^T R \mathbf{v}$$

$$\|\mathbf{v}\|^2 = \mathbf{v}^T \mathbf{v}$$

$$\Rightarrow R^T R = I$$

Properties of rotation matrices

$$\Rightarrow R^T R = I$$

$$\Rightarrow \det(R)^2 = 1$$

$$\Rightarrow \det(R) = \pm 1$$

$$\det(R) = 1$$

Rotation

$$\det(R) = -1$$


Reflection

Rotation matrices

- Rotations in 3D have an axis and an angle
- Axis: vector that does not change when rotated

$$R\mathbf{v} = \mathbf{v}$$

- Rotation matrix has eigenvector that has eigenvalue 1



Rotation matrices from axis and angle

- Rotation matrix for rotation about axis \mathbf{v} and θ
- First define the following matrix

$$[\mathbf{v}]_{\times} = \begin{bmatrix} 0 & -v_z & v_y \\ v_z & 0 & -v_x \\ -v_y & v_x & 0 \end{bmatrix}$$

- Interesting fact: this matrix represents cross product

$$[\mathbf{v}]_{\times} \mathbf{x} = \mathbf{v} \times \mathbf{x}$$

Rotation matrices from axis and angle

- Rotation matrix for rotation about axis \mathbf{v} and θ
- Rodrigues' formula for rotation matrices

$$R = I + (\sin \theta)[\mathbf{v}]_{\times} + (1 - \cos \theta)[\mathbf{v}]_{\times}^2$$

Translations

$$\mathbf{x}' = \mathbf{x} + \mathbf{t}$$

- Can this be written as a matrix multiplication?

Putting everything together

- Change coordinate system so that center of the coordinate system is at pinhole and Z axis is along viewing direction

$$\mathbf{x}'_w = R\mathbf{x}_w + \mathbf{t}$$

- Perspective projection

$$\mathbf{x}'_w \equiv (X, Y, Z)$$

$$\mathbf{x}'_{img} \equiv (x, y)$$

$$x = \frac{X}{Z}$$

$$y = \frac{Y}{Z}$$

The projection equation

$$\begin{aligned}x &= \frac{X}{Z} \\y &= \frac{Y}{Z}\end{aligned}$$

- Is this equation linear?
- Can this equation be represented by a matrix multiplication?

Is projection linear?

$$X' = aX + b$$

$$Y' = aY + b$$

$$Z' = aZ + b$$

$$x' = \frac{aX + b}{aZ + b}$$

$$y' = \frac{aY + b}{aZ + b}$$

Can projection be represented as a matrix multiplication?

Matrix multiplication


$$\begin{bmatrix} a & b & c \\ p & q & r \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} aX + bY + cZ \\ pX + qY + rZ \end{bmatrix}$$

Perspective
projection

$$x = \frac{X}{Z}$$
$$y = \frac{Y}{Z}$$

The space of rays

- Every point on a ray maps it to a point on image plane
- Perspective projection maps rays to points
- All points $(\lambda x, \lambda y, \lambda)$ map to the same image point $(x, y, 1)$



Projective space

- Standard 2D space (plane) \mathbb{R}^2 : Each point represented by 2 coordinates (x,y)
- Projective 2D space (plane) \mathbb{P}^2 : Each “point” represented by 3 coordinates (x,y,z) , BUT:
 - $(\lambda x, \lambda y, \lambda z) \equiv (x, y, z)$
- Mapping \mathbb{R}^2 to \mathbb{P}^2 (points to rays):
$$(x, y) \rightarrow (x, y, 1)$$
- Mapping \mathbb{P}^2 to \mathbb{R}^2 (rays to points):
$$(x, y, z) \rightarrow \left(\frac{x}{z}, \frac{y}{z} \right)$$

Projective space and homogenous coordinates

- Mapping \mathbb{R}^2 to \mathbb{P}^2 (points to rays):

$$(x, y) \rightarrow (x, y, 1)$$

- Mapping \mathbb{P}^2 to \mathbb{R}^2 (rays to points):

$$(x, y, z) \rightarrow \left(\frac{x}{z}, \frac{y}{z} \right)$$


- A change of coordinates
- Also called *homogenous coordinates*

Homogenous coordinates


- In standard Euclidean coordinates
 - 2D points : (x,y)
 - 3D points : (x,y,z)
- In homogenous coordinates
 - 2D points : $(x,y,1)$
 - 3D points : $(x,y,z,1)$

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$



Homogenous
coordinates of
world point



Homogenous
coordinates of
image point

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$

$$P\vec{\mathbf{x}}_w = \vec{\mathbf{x}}_{img}$$

- Perspective projection is matrix multiplication in homogenous coordinates!

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

- Translation is matrix multiplication in homogenous coordinates!

Homogenous coordinates

$$\begin{bmatrix} a & b & c & t_x \\ d & e & f & t_y \\ g & h & i & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} aX + bY + cZ + t_x \\ dX + eY + fZ + t_y \\ gX + hY + iZ + t_z \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \begin{bmatrix} \mathbf{x}_w \\ 1 \end{bmatrix} = \begin{bmatrix} M\mathbf{x}_w + \mathbf{t} \\ 1 \end{bmatrix}$$

Homogenous coordinates

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \longrightarrow [I \quad \mathbf{0}]$$

Perspective projection in homogenous coordinates

$$\vec{\mathbf{x}}_{img} = \begin{bmatrix} I & \mathbf{0} \end{bmatrix} \begin{bmatrix} R & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \vec{\mathbf{x}}_w$$

$$\vec{\mathbf{x}}_{img} = \begin{bmatrix} R & \mathbf{t} \end{bmatrix} \vec{\mathbf{x}}_w$$

More about matrix transformations

$\begin{bmatrix} I & \mathbf{0} \end{bmatrix}$ 3 x 4 : Perspective projection

$\begin{bmatrix} I & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$ 4 x 4 : Translation


$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$ 4 x 4 : Affine transformation
(linear transformation + translation)

More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M^T M = I$$

Euclidean





More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M = sR$$

$$R^T R = I$$

Similarity
transformation





More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{bmatrix}$$



Anisotropic scaling and translation



More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

General affine transformation



Matrix transformations in 2D

