


Image formation

The projection equation



$$x = \frac{X}{Z}$$

$$y = \frac{Y}{Z}$$



Changing coordinate systems





Changing coordinate systems





Changing coordinate systems





Changing coordinate systems



Changing coordinate systems



Changing coordinate systems



Rotations and translations

- How do you represent a rotation?
- A point in 3D: (X,Y,Z)
- Rotations can be represented as a matrix multiplication

- What are the properties of rotation matrices?
 $\mathbf{v}' = R\mathbf{v}$

Properties of rotation matrices

- Rotation does not change the length of vectors

$$\mathbf{v}' = R\mathbf{v}$$

$$\|\mathbf{v}'\|^2 = \mathbf{v}'^T \mathbf{v}'$$

$$= \mathbf{v}^T R^T R \mathbf{v}$$

$$\|\mathbf{v}\|^2 = \mathbf{v}^T \mathbf{v}$$

$$\Rightarrow R^T R = I$$

Properties of rotation matrices

$$\Rightarrow R^T R = I$$

$$\Rightarrow \det(R)^2 = 1$$

$$\Rightarrow \det(R) = \pm 1$$

$$\det(R) = 1$$

Rotation

$$\det(R) = -1$$


Reflection

Rotation matrices

- Rotations in 3D have an axis and an angle
- Axis: vector that does not change when rotated

$$R\mathbf{v} = \mathbf{v}$$

- Rotation matrix has eigenvector that has eigenvalue 1



Rotation matrices from axis and angle

- Rotation matrix for rotation about axis \mathbf{v} and θ
- First define the following matrix

$$[\mathbf{v}]_{\times} = \begin{bmatrix} 0 & -v_z & v_y \\ v_z & 0 & -v_x \\ -v_y & v_x & 0 \end{bmatrix}$$

- Interesting fact: this matrix represents cross product

$$[\mathbf{v}]_{\times} \mathbf{x} = \mathbf{v} \times \mathbf{x}$$

Rotation matrices from axis and angle

- Rotation matrix for rotation about axis \mathbf{v} and θ
- Rodrigues' formula for rotation matrices

$$R = I + (\sin \theta)[\mathbf{v}]_{\times} + (1 - \cos \theta)[\mathbf{v}]_{\times}^2$$

Translations

$$\mathbf{x}' = \mathbf{x} + \mathbf{t}$$

- Can this be written as a matrix multiplication?

Putting everything together

- Change coordinate system so that center of the coordinate system is at pinhole and Z axis is along viewing direction

$$\mathbf{x}'_w = R\mathbf{x}_w + \mathbf{t}$$

- Perspective projection

$$\begin{aligned}\mathbf{x}'_w &\equiv (X, Y, Z) & x &= \frac{X}{Z} \\ \mathbf{x}'_{img} &\equiv (x, y) & y &= \frac{Y}{Z}\end{aligned}$$

The projection equation

$$\begin{aligned}x &= \frac{X}{Z} \\y &= \frac{Y}{Z}\end{aligned}$$

- Is this equation linear?
- Can this equation be represented by a matrix multiplication?

Is projection linear?

$$X' = aX + b$$

$$Y' = aY + b$$

$$Z' = aZ + b$$

$$x' = \frac{aX + b}{aZ + b}$$

$$y' = \frac{aY + b}{aZ + b}$$

Can projection be represented as a matrix multiplication?

Matrix multiplication


$$\begin{bmatrix} a & b & c \\ p & q & r \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} aX + bY + cZ \\ pX + qY + rZ \end{bmatrix}$$

Perspective
projection

$$x = \frac{X}{Z}$$
$$y = \frac{Y}{Z}$$

The space of rays

- Every point on a ray maps it to a point on image plane
- Perspective projection maps rays to points
- All points $(\lambda x, \lambda y, \lambda)$ map to the same image point $(x, y, 1)$



Projective space

- Standard 2D space (plane) \mathbb{R}^2 : Each point represented by 2 coordinates (x, y)
- Projective 2D space (plane) \mathbb{P}^2 : Each “point” represented by 3 coordinates (x, y, z) , BUT:
 - $(\lambda x, \lambda y, \lambda z) \equiv (x, y, z)$
- Mapping \mathbb{R}^2 to \mathbb{P}^2 (points to rays):
$$(x, y) \rightarrow (x, y, 1)$$
- Mapping \mathbb{P}^2 to \mathbb{R}^2 (rays to points):
$$(x, y, z) \rightarrow \left(\frac{x}{z}, \frac{y}{z} \right)$$

Projective space and homogenous coordinates


- Mapping \mathbb{R}^2 to \mathbb{P}^2 (points to rays):
$$(x, y) \rightarrow (x, y, 1)$$
- Mapping \mathbb{P}^2 to \mathbb{R}^2 (rays to points):
$$(x, y, z) \rightarrow \left(\frac{x}{z}, \frac{y}{z}\right)$$
- A change of coordinates
- Also called *homogenous coordinates*

Homogenous coordinates


- In standard Euclidean coordinates
 - 2D points : (x,y)
 - 3D points : (x,y,z)
- In homogenous coordinates
 - 2D points : $(x,y,1)$
 - 3D points : $(x,y,z,1)$

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$



Homogenous
coordinates of
world point



Homogenous
coordinates of
image point

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$

$$P\vec{\mathbf{x}}_w = \vec{\mathbf{x}}_{img}$$

- Perspective projection is matrix multiplication in homogenous coordinates!

Why homogenous coordinates?

$$\begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

- Translation is matrix multiplication in homogenous coordinates!

Homogenous coordinates

$$\begin{bmatrix} a & b & c & t_x \\ d & e & f & t_y \\ g & h & i & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} aX + bY + cZ + t_x \\ dX + eY + fZ + t_y \\ gX + hY + iZ + t_z \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \begin{bmatrix} \mathbf{x}_w \\ 1 \end{bmatrix} = \begin{bmatrix} M\mathbf{x}_w + \mathbf{t} \\ 1 \end{bmatrix}$$

Homogenous coordinates

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \equiv \begin{bmatrix} \frac{X}{Z} \\ \frac{Y}{Z} \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \longrightarrow [I \quad \mathbf{0}]$$

Perspective projection in homogenous coordinates

$$\vec{\mathbf{x}}_{img} = \begin{bmatrix} I & \mathbf{0} \end{bmatrix} \begin{bmatrix} R & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \vec{\mathbf{x}}_w$$

$$\vec{\mathbf{x}}_{img} = \begin{bmatrix} R & \mathbf{t} \end{bmatrix} \vec{\mathbf{x}}_w$$

More about matrix transformations



$$\begin{bmatrix} I & \mathbf{0} \end{bmatrix} \quad 3 \times 4 : \text{Perspective projection}$$
$$\begin{bmatrix} I & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \quad 4 \times 4 : \text{Translation}$$
$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix} \quad 4 \times 4 : \text{Affine transformation} \\ \text{(linear transformation + translation)}$$

More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M^T M = I$$

Euclidean





More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M = sR$$

$$R^T R = I$$

Similarity
transformation





More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

$$M = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{bmatrix}$$



Anisotropic scaling and translation



More about matrix transformations

$$\begin{bmatrix} M & \mathbf{t} \\ \mathbf{0}^T & 1 \end{bmatrix}$$

General affine
transformation



Matrix transformations in 2D

