

DSFA
Spring 2021

Lecture 22

Confidence Intervals


Announcements

- HW 4 due today, 5:59PM
 - Project 2 out today
 - Part 1 due 4/9, 5:59PM
 - Part 2 due 4/16, 5:59PM
 - Next two lab sessions (4/7-8, 4/14-15) will be dedicated to working on project
 - Can have a partner again; preferably from the same section
 - Prelim 2, 4/20, 8:30-10PM, here
-


The Bootstrap

Why the Bootstrap Works




population



sample



resamples




All of these look pretty similar, most likely.

Key to Resampling

- From the original sample,
 - draw at random
 - with replacement
 - as many values as the original sample contained
 - The size of the new sample has to be the same as the original one, so that the two estimates are comparable
-

Why the Bootstrap Works




Inference Using the Bootstrap




population



sample



resamples



All of these look pretty similar, most likely.

95% Confidence Interval

- Interval of **estimates of a parameter**
- Based on random sampling
- 95% is called the confidence level
 - Could be any percent between 0 and 100
 - Bigger means wider intervals
- The **confidence is in the process** that generated the interval:
 - It generates a “good” interval about 95% of the time.

(Demo)

Use Methods Appropriately

When *Not* to Use The Bootstrap

- If you're trying to estimate very high or very low percentiles, or min and max
 - If you're trying to estimate any parameter that's greatly affected by rare elements of the population
 - If the probability distribution of your statistic is not roughly bell shaped (the shape of the empirical distribution will be a clue)
 - If the original sample is very small (~ 15)
-

Can You Use a CI Like This?

By our calculation, an approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

True or False:

- About 95% of the mothers in the population were between 26.9 years and 27.6 years old.

When poll is active, respond at pollev.com/dsfa

Untitled true or false question

True

False



Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Can You Use a CI Like This?

By our calculation, an approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

True or False:

- About 95% of the mothers in the population were between 26.9 years and 27.6 years old.

Answer: False. We're estimating that their **average age** is in this interval.

(Demo)

Is This What a CI Means?

By our calculation, an approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

True or False:

- There is a 0.95 probability that the average age of mothers in the population is in the range 26.9..27.6 years.
-

There is a 0.95 probability that the average age of mothers in the population is in the range 26.9-27.6

True

False



Is This What a CI Means?

By our calculation, an approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

True or False:

- There is a 0.95 probability that the average age of mothers in the population is in the range 26.9..27.6 years.

Answer: False. It's not a probability; that's either true or false.

Confidence Interval Tests

Using a CI for Testing

- Null hypothesis: **Population mean = x**
 - Alternative hypothesis: **Population mean $\neq x$**
 - Cutoff for P-value: $p\%$
 - Method:
 - Construct a $(100-p)\%$ confidence interval for the population statistic
 - If x is not in the interval, reject the null
 - If x is in the interval, can't reject the null
-