

CS322  
Homework #1  
Due Friday, July 11, 2003

In the Beginning...

Answer all questions. Submit any pertinent hand calculations and/or Matlab code in a neat manner in order to help us give you partial credit. Readings: Van Loan 2.0-2.3, 3.0-3.3.

- 1) Describe briefly an example of each of the following:
  - a) A numerical method that is inefficient.
  - b) A numerical method that is inaccurate.

2) Describe briefly (a sentence or two for each) three topics that will be covered in this class. The text will be helpful here.

3) *A numerical analyst's answer to "hello world"*. Let  $f(x) = \pi \sin 3x + e^{-3} \cos 2x$  on the interval  $[0, 3\pi]$  and  $n = 3$ . Let  $V_n$  and  $N_n$  denote the Vandermonde and Newton representations of polynomial interpolants of  $f$ , respectively. (a) Should  $V_n$  be equal to  $N_n$  for all  $x \in [0, 3\pi]$ ? (b) Why or why not?

Let  $g$  be some function defined on the interval  $I$  and let  $p$  be some interpolant of  $g$ . Denote by  $ae$  the *absolute error* and define  $ae$  by

$$ae(x) = |g(x) - p(x)|,$$

where  $x$  is any point in  $I$ .

(c) Let  $\underline{x} = (x_1, x_2, x_3) = (0, \frac{3\pi}{2}, 3\pi)$  and  $y_i = f(x_i)$ , for  $i = 1, 2, 3$ . Find  $ae(2\pi)$  for our  $f$  using both  $V_n$  and  $N_n$ . I worked out the following crude upper bound for  $f$ 's third derivative,

$$\left| f^{(3)}(\eta) \right| \leq 85.168,$$

for some  $\eta \in [0, 3\pi]$ , (d) are your observed  $ae$ 's smaller than our theoretical bound?

4) *It's All About Who Represents You*. Recall the cubic representation given in the text:

if

$$q(z) = a + b(z - x_L) + c(z - x_L)^2 + d(z - x_L)^2(z - x_R),$$

then

$$\begin{aligned} q(x_L) &= y_L, & q(x_R) &= y_R, \\ q'(x_L) &= s_L, & q'(x_R) &= s_R. \end{aligned}$$

Consider the following, alternative representation, that has the same conclusion:

if

$$q_c(z) = A + Bz + Cz^2 + Dz^3,$$

then

$$\begin{aligned} q_c(x_L) &= y_L, & q_c(x_R) &= y_R, \\ q'_c(x_L) &= s_L, & q'_c(x_R) &= s_R. \end{aligned}$$

(a) Write down the matrix equations that result from this alternative. (b) Find  $D$ , thereby showing that

$$D = \frac{(x_L - x_R)(s_L + s_R) - 2(y_L - y_R)}{(x_L - x_R)^3}.$$

Consider how much work it took you to find  $D$ , and in at most two sentences (c) comment on the virtues of the original cubic representation. (d) Where else have we seen this virtue (in the readings and/or lecture)?