

# CS 3220: PRELIM 1 EXAMPLE QUESTIONS

Instructor: Anil Damle

## WHATS IN THIS DOCUMENT

These questions are intended to be somewhat representative of the type of questions that could arise on the first prelim.

---

### QUESTION 1

Here, we consider some properties of  $\|A\|_F$ . For what follows  $A \in \mathbb{R}^{n \times n}$

(a) Prove that

$$\|A\|_F^2 = \sum_{i=1}^n \|A(:, i)\|_2^2.$$

(b) For any two orthogonal matrices  $Q_1 \in \mathbb{R}^{n \times n}$  and  $Q_2 \in \mathbb{R}^{n \times n}$  prove that

$$\|Q_1 A Q_2\|_F = \|A\|_F$$

(c) Prove that

$$\|A\|_F = \sqrt{\sum_{i=1}^n \sigma_i^2},$$

where  $\sigma_1, \dots, \sigma_n$  are the singular values of  $A$ .

### SOLUTION

(a) We have that  $\|A\|_F^2 = \sum_{i,j} A_{i,j}^2$ , which is equivalent to

$$\|A\|_F^2 = \sum_{j=1}^n \sum_{i=1}^n A_{i,j}^2.$$

The inner loop is  $\|A(:, j)\|_2^2$ , thereby completing the proof.

(b) Since the two norm of vectors is invariant under orthogonal transformations, the prior part gives us immediately that

$$\begin{aligned} \|Q_1 A Q_2\|_F^2 &= \sum_{i=1}^n \|Q_1 (A Q_2)(:, i)\|_2^2 \\ &= \sum_{i=1}^n \|(A Q_2)(:, i)\|_2^2 \\ &= \|A Q_2\|_F^2. \end{aligned}$$

Using the fact that for any matrix  $B$ ,  $\|B\|_F^2 = \|B^T\|_F^2$  we can use the same argument to remove  $Q_2$ .

(c) If  $A = U\Sigma V^T$  we have that

$$\|A\|_F^2 = \|\Sigma\|_F^2.$$

Explicitly writing out the right hand side and taking the square root yields the result.

## QUESTION 2

Here, we ask you to interpret the condition number of a  $2 \times 2$  matrix geometrically. (Hint: pictures are useful here!)

1. We saw that the SVD of a  $2 \times 2$  matrix allows us to view any matrix  $A$  as mapping a circle to an ellipse. If  $A$  becomes increasingly ill-conditioned what is geometrically happening to the ellipse?
2. Geometrically argue why for an ill-conditioned matrix a small relative change in  $b$  can result in a big relative change in  $x$ .

## SOLUTION

- (a) The ratio of the length of the major axis to minor axis of the ellipse is going to infinity, so the ellipse is collapsing to a line segment.
- (b) Because of the elongated shape of the ellipse, changes to  $b$  in the direction of  $v_2$  can drastically alter the location along the ellipse.

## QUESTION 3

Say you are given a symmetric matrix  $A$  and tasked with computing the algebraically smallest eigenvalue. Using only the power method (applied to  $A$  or matrices related to  $A$ ), how might you go about doing this? (Hint: think about how the eigenvalues/vectors of  $A - \gamma I$  relate to those of  $A$  for any scalar  $\gamma \in \mathbb{R}$ .)

## SOLUTION

We can first use the power method to compute the largest eigenvalue in magnitude of  $A$ . If it is negative we are done, if not we need to do a bit more work. Let  $\mu$  be the eigenvalue of  $A$  that we previously computed. Now, we can simply use the power method to compute an eigenvalue of  $A - \mu I$ , call it  $\lambda$  and then  $\lambda + \mu$  is the algebraically smallest eigenvalue of  $A$ . This works because the eigenvalues of  $A - \mu I$  are simply those of  $A$  shifted right by  $\mu$ . Therefore the algebraically largest eigenvalue of  $A - \mu I$  is 0 and the largest magnitude one is necessarily the smallest algebraic.