

CS4620/5620: Lecture 28

Splines

Announcements


- Practicum
 - This week (PPA2)
 - Next week (PPA3)

Defining spline curves

- At the most general they are parametric curves

$$S = \{\mathbf{p}(t) \mid t \in [0, N]\}$$

- Generally $p(t)$ is a piecewise polynomial
 - the discontinuities are at the integers



Defining spline curves

- Generally $p(t)$ is a piecewise polynomial
 - the discontinuities are at the integers
 - e.g., a cubic spline has the following form over $[k, k + 1]$:


$$x(t) = a_x t^3 + b_x t^2 + c_x t + d_x$$

$$y(t) = a_y t^3 + b_y t^2 + c_y t + d_y$$

- Coefficients are different for every interval


Control of spline curves

- Specified by a sequence of control points
- Shape is guided by control points (aka control polygon)
 - interpolating: passes through points
 - approximating: merely guided by points



Trivial example: piecewise linear

- This spline is just a polygon
 - control points are the vertices
- But we can derive it anyway as an illustration
- Each interval will be a linear function
 - $x(t) = at + b$
 - constraints are values at endpoints
 - $b = x_0$; $a = x_1 - x_0$
 - this is linear interpolation



Trivial example: piecewise linear

- Basis function formulation
 - regroup expression by \mathbf{p} rather than t


$$\begin{aligned}\mathbf{p}(t) &= (\mathbf{p}_1 - \mathbf{p}_0)t + \mathbf{p}_0 \\ &= (1 - t)\mathbf{p}_0 + t\mathbf{p}_1\end{aligned}$$

- interpretation in matrix viewpoint

$$\mathbf{p}(t) = \begin{pmatrix} [t & 1] \begin{bmatrix} -1 & 1 \\ 1 & 0 \end{bmatrix} \end{pmatrix} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \end{bmatrix}$$


Trivial example: piecewise linear

- Vector blending formulation: “average of points”
 - blending functions: contribution of each point as t changes



Trivial example: piecewise linear


- Basis function formulation: “function times point”
 - basis functions: contribution of each point as t changes



- can think of them as blending functions glued together
- this is just like a reconstruction filter!

Hermite splines

- Less trivial example
- Form of curve: piecewise cubic
 - How many constraints?
- Constraints: endpoints and tangents (derivatives)



Defining spline curves

- Generally $p(t)$ is a piecewise polynomial
 - the discontinuities are at the integers
 - e.g., a cubic spline has the following form over $[k, k + 1]$:

$$x(t) = a_x t^3 + b_x t^2 + c_x t + d_x$$

$$y(t) = a_y t^3 + b_y t^2 + c_y t + d_y$$

- Coefficients are different for every interval

Hermite splines

- Solve constraints to find coefficients

$$x(t) = at^3 + bt^2 + ct + d$$

$$x'(t) = 3at^2 + 2bt + c$$

$$x(0) = x_0 = d$$

$$x(1) = x_1 = a + b + c + d$$

$$x'(0) = x'_0 = c$$

$$x'(1) = x'_1 = 3a + 2b + c$$

$$d = x_0$$

$$c = x'_0$$

$$a = 2x_0 - 2x_1 + x'_0 + x'_1$$

$$b = -3x_0 + 3x_1 - 2x'_0 - x'_1$$

Hermite splines

- Matrix form is much simpler

$$\mathbf{p}(t) = \begin{bmatrix} t^3 & t^2 & t & 1 \end{bmatrix} \begin{bmatrix} 2 & -2 & 1 & 1 \\ -3 & 3 & -2 & -1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \mathbf{v}_0 \\ \mathbf{v}_1 \end{bmatrix}$$

– coefficients = rows

– basis functions = columns

Matrix form of spline

$$\mathbf{p}(t) = \mathbf{a}t^3 + \mathbf{b}t^2 + \mathbf{c}t + \mathbf{d}$$

$$\begin{bmatrix} t^3 & t^2 & t & 1 \end{bmatrix} \begin{bmatrix} \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \\ \times & \times & \times & \times \end{bmatrix} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \mathbf{p}_2 \\ \mathbf{p}_3 \end{bmatrix}$$

$$\mathbf{p}(t) = \mathbf{b}_0(t)\mathbf{p}_0 + \mathbf{b}_1(t)\mathbf{p}_1 + \mathbf{b}_2(t)\mathbf{p}_2 + \mathbf{b}_3(t)\mathbf{p}_3$$

Hermite splines

- Matrix form is much simpler


$$\mathbf{p}(t) = \begin{bmatrix} t^3 & t^2 & t & 1 \end{bmatrix} \begin{bmatrix} 2 & -2 & 1 & 1 \\ -3 & 3 & -2 & -1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \mathbf{v}_0 \\ \mathbf{v}_1 \end{bmatrix}$$

– coefficients = rows

– basis functions = columns

Hermite splines


- Hermite basis functions



Longer Hermite splines


- Can only do so much with one Hermite spline
- Can use these splines as segments of a longer curve
 - curve from $t = 0$ to $t = 1$ defined by first segment
 - curve from $t = 1$ to $t = 2$ defined by second segment
- To avoid discontinuity, match derivatives at junctions
 - this produces a C^1 curve

Hermite basis



Hermite to Bézier

- Mixture of points and vectors is awkward
- Specify tangents as differences of points



Hermite to Bézier


$$p_0 = q_0$$

$$p_1 = q_3$$

$$v_0 = 3(q_1 - q_0)$$

$$v_1 = 3(q_3 - q_2)$$

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} q_0 \\ q_1 \\ q_2 \\ q_3 \end{bmatrix}$$



Bézier matrix

$$\mathbf{p}(t) = \begin{bmatrix} t^3 & t^2 & t & 1 \end{bmatrix} \begin{bmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \mathbf{p}_2 \\ \mathbf{p}_3 \end{bmatrix}$$

– note that these are the Bernstein polynomials

$$C(n,k) t^k (1-t)^{n-k}$$

and that defines Bézier curves for any degree

Bézier basis

