

ABSOLUTE CONTINUITY BETWEEN THE WIENER AND STATIONARY GAUSSIAN MEASURES

U. KEICH

ABSTRACT. It is known that the entropy distance between two Gaussian measures is finite if, and only if, they are absolutely continuous with respect to one another. Shepp [5] characterized the correlations corresponding to stationary Gaussian measures that are absolutely continuous with respect to the Wiener measure. By analyzing the entropy distance, we show that one of his conditions, involving the spectrum of an associated operator, is essentially extraneous, providing a simple criterion for finite entropy distance in this case.

1. INTRODUCTION

Let $C[1 - \tau, 1 + \tau]$ (where $0 < \tau < 1$) denote the space of continuous functions on $[1 - \tau, 1 + \tau]$. A standard Brownian motion observed between times $1 - \tau$ and $1 + \tau$ induces on $C[1 - \tau, 1 + \tau]$ the Wiener measure W^τ . As a Gaussian measure, it is characterized by its correlation $R(t, s) = t \wedge s$ for $t, s \in [1 - \tau, 1 + \tau]$, and by its vanishing mean.

A Gaussian measure, Q^τ , on $C[1 - \tau, 1 + \tau]$ is *stationary* if its mean is constant and its correlation is a Töeplitz function. That is, with $X \in C[1 - \tau, 1 + \tau]$ being the sample path,

$$\mu_t \stackrel{d}{=} E^{Q^\tau} X_t \equiv \text{const.} \quad t \in [1 - \tau, 1 + \tau],$$

and

$$S(t, s) \stackrel{d}{=} E^{Q^\tau} X_t X_s - \mu^2 = S(t - s) \quad t, s \in [1 - \tau, 1 + \tau],$$

where S denotes both the correlation $S(t, s)$, and the auto-correlation $S(r)$ with $r = t - s \in [-2\tau, 2\tau]$. Krein [4] showed that $S(r)$ can always be extended to \mathbb{R} as a continuous positive-definite function, thereby providing an extension of Q^τ to a stationary Gaussian measure, Q , on $C(\mathbb{R})$.

We would like to characterize the stationary measures Q^τ which are absolutely continuous with respect to W^τ . Since the measures are

AMS 1991 subject classifications. 60G10, 60G15, 60G30.

Key words and phrases. stationary Gaussian measures, entropy distance, absolute continuity, Wiener measure.

Gaussian, this question can be settled in terms of the mean μ and the correlation S of Q^τ . Furthermore, for $\tau < 1$, the Brownian paths b_t and $b_t + \mu$ with $t \in [1 - \tau, 1 + \tau]$ induce measures that are absolutely continuous with respect to one another, and therefore the constant mean of Q is irrelevant to the question of absolute continuity.

Shepp provides the following necessary and sufficient conditions for a stationary Gaussian measure Q^{τ_0} with correlation S to be absolutely continuous with respect to W^{τ_0} [5]:

(i) On $(0, 2\tau_0)$, S' is absolutely continuous and S'' satisfies

$$\int_0^{2\tau_0} S''(t)^2 (2\tau_0 - t) dt < \infty.$$

(ii) $S^+(0) = -1/2$ (the derivative from the right).

(iii) $-1 \notin \sigma(F)$, where $\sigma(F)$ is the spectrum of the integral operator defined by the kernel

$$F \stackrel{d}{=} \frac{\partial}{\partial s} \frac{\partial}{\partial t} \left[S(t-s) - \frac{S(t)S(s)}{S(0)} - t \wedge s \right] \quad t, s \in [0, 2\tau_0].$$

Shepp gives an example showing that (iii) is essential:

$$S(r) \stackrel{d}{=} \frac{1}{4} - \frac{|r|}{2} \quad \text{for} \quad -2\tau_0 \leq r \leq 2\tau_0.$$

Here (i) and (ii) hold for any $\tau_0 > 0$, but (iii) is valid only if $\tau_0 < \frac{1}{2}$ ($\tau_0 \leq \frac{1}{2}$ is required for S to be positive-definite). Indeed, with $\tau_0 = \frac{1}{2}$ and $X \in C[1 - \tau_0, 1 + \tau_0]$, we have $X_{1/2} = -X_{3/2}$ a.s. dQ , thus ruling out absolute continuity with respect to W^{τ_0} . However, a closer look at this example yields two interesting facts for $\tau_0 = \frac{1}{2}$:

- There exists only one positive-definite extension of S from $[-2\tau_0, 2\tau_0]$ to \mathbb{R} ([4]).
- S'' does not exist at $r = 2\tau_0$.

Both observations turn out to be the rule whenever (iii) is violated. This allows us to rid ourselves of the third condition in Shepp's theorem by paying a small price: (i) and (ii) suffice for absolute continuity for $\tau < \tau_0$. The precise statement is:

Theorem 1. *If Q^{τ_0} , a stationary Gaussian measure with correlation S , is absolutely continuous with respect to W^{τ_0} ($0 < \tau_0 < 1$), then (i) and (ii) hold. If, on the other hand, (i) and (ii) hold, then for any $0 < \tau < \tau_0$, Q^τ is absolutely continuous with respect to W^τ .*

Remarks:

- A simple scaling argument shows that the time interval can be centered about any point (not necessarily 1).

- Condition (ii) guarantees that the measure Q is supported on paths with the same quadratic variation as that of Brownian motion. This must be so, as it is an “almost sure” property of the Brownian path.

We next provide a brief overview of the proof which is entirely different from Shepp’s; it relies on our ability to estimate the entropy distance between Q^τ and W^τ . Let P and Q be Gaussian measures on $C[1 - \tau, 1 + \tau]$. It is known that the entropy distance between P and Q , $H(P, Q)$, is finite if and only if the two measures are absolutely continuous with respect to one another; in that case:

$$H(P, Q) = E^P\left(\log \frac{dP}{dQ}\right) + E^Q\left(\log \frac{dQ}{dP}\right),$$

where E^P is the expectation under the measure P and $\frac{dP}{dQ}$ is the Radon-Nikodym derivative of P with respect to Q (see e.g. [3]).

Let P_n and Q_n be the restrictions of P and Q to $n+1$ equally spaced points in $[1 - \tau, 1 + \tau]$, with correlations R_n and S_n . Then

$$H(P, Q) = \overline{\lim} H(P_n, Q_n).$$

Let K_n be a root of R_n , i.e., $R_n = K_n K_n^*$, and let $T_n = K_n^{-1} S_n K_n^{-*}$, with K^{-*} being a short for $(K^{-1})^*$. If λ_i^n are the eigenvalues of T_n , then

$$H(P_n, Q_n) = \frac{1}{2} \sum_{i=0}^n \frac{(\lambda_i^n - 1)^2}{\lambda_i^n}.$$

Thus, $H(W^{\tau_0}, Q^{\tau_0}) < \infty$ implies

$$(1) \quad \sup_{\tau \leq \tau_0} \overline{\lim} \sum (\lambda_i^n - 1)^2 < \infty.$$

In the case of W^{τ_0} ($R = t \wedge s$), R_n^{-1} is essentially a second order difference operator so we can choose K_n so that K_n^{-1} is basically a first order difference operator. Thus, the typical entry in $T_n = K_n^{-1} S_n K_n^{-*}$ is a second order difference of the sampled Töeplitz correlation S_n . Add to this the identity

$$\overline{\lim} \sum (\lambda_i^n - 1)^2 = \overline{\lim} \text{Tr} (T_n - I)^2,$$

and you will understand how (1) implies the existence of S' and S'' on $(0, 2\tau_0)$ as in (i) and (ii) of theorem 1.

This is the easier half of that theorem. As for the other half, with our choice of K_n we can readily show that (1) follows from (i) and (ii) of theorem 1. It is left to prove that for $\tau < \tau_0$, $\inf_{i,n} \lambda_i^n(\tau) > 0$ (this

is the analogue of Shepp's third condition, which we omit). Indeed, if $\inf_{i,n} \lambda_i^n(\tau) = 0$, then there exist $\lambda_k \rightarrow 0$ and $\mathbf{v}_k \in \mathbb{R}^{n_k}$, such that

$$T_{n_k} \mathbf{v}_k = \lambda_k \mathbf{v}_k.$$

Appropriately embedded in $\mathcal{H} \stackrel{d}{=} \mathbb{R} \oplus L^2[0, 2\tau]$, \mathbf{v}_k converge strongly to a non-zero limit $(\beta, f) \in \mathcal{H}$ with the property that almost surely dQ :

$$\beta X_0 + \int_0^{2\tau} f(t) dX_t = 0,$$

where Q is a stationary extension of Q^τ to $C(\mathbb{R})$ and X_t is the sample path. As we show, this implies that S has a unique positive-definite extension, from $[-2\tau, 2\tau]$ to \mathbb{R} . This unique extension has the property that $S'' \notin L^2(0, 2\tau + \varepsilon)$ for any $\varepsilon > 0$, contradicting (i).

2. PROOF OF THEOREM 1

The space $C_0 \stackrel{d}{=} C[1 - \tau_0, 1 + \tau_0]$, is equipped with the σ -field \mathcal{F} generated by the cylinder sets. Wiener measure (denoted by W) is defined on that space, so any measure Q that is absolutely continuous with respect to it must live there. On the other hand, an application of Kolmogorov-Čensov shows that a correlation S subject to (i) and (ii) of theorem 1 defines a probability measure Q on C_0 .

Remark. Since the mean of Q is irrelevant to our problem it may be assumed to vanish.

Let P and Q be Gaussian measures on C_0 . For $0 < \tau \leq \tau_0$, let P^τ , Q^τ and \mathcal{F}^τ be the restrictions of P , Q , respectively \mathcal{F} , to $C[1 - \tau, 1 + \tau]$. Let \mathcal{F}_n^τ be the σ -field obtained by sampling the paths on $[1 - \tau, 1 + \tau]$ at $n + 1$ equally spaced points. Let P_n^τ and Q_n^τ be the restrictions of P^τ and Q^τ to that σ -field. Let

$$H_n^\tau \stackrel{d}{=} H(P_n^\tau, Q_n^\tau).$$

Since \mathcal{F}^τ is generated by $\bigcup \mathcal{F}_n^\tau$, it follows that $H(P^\tau, Q^\tau) = \overline{\lim} H_n^\tau$ [3].

A first step is to express the entropy H_n^τ in terms of the $(n+1) \times (n+1)$ sampled correlations, S_n^τ and R_n^τ .

Claim 2.1. Let R and S be the $m \times m$ correlation matrices of the 0-mean Gaussian measures P , respectively Q on \mathbb{R}^m , and let K be a root of R so that $R = KK^*$. Then

$$H(P, Q) = \frac{1}{2} \text{Tr}(K^{-1}SK^{-*} + K^*S^{-1}K - 2I).$$

Proof.

$$\begin{aligned} E^Q \log \frac{dQ}{dP} &= E^Q \log \left\{ \frac{\sqrt{\det R}}{\sqrt{\det S}} \exp \left[-\frac{1}{2} \langle (S^{-1} - R^{-1}) \mathbf{x}, \mathbf{x} \rangle \right] \right\} \\ &= -\frac{1}{2} \log \det R^{-1} S - \frac{1}{2} E^Q \langle (S^{-1} - R^{-1}) \mathbf{x}, \mathbf{x} \rangle, \end{aligned}$$

with $\mathbf{x} \in \mathbb{R}^m$.

If A is any $m \times m$ matrix, then

$$E^Q \langle A \mathbf{x}, \mathbf{x} \rangle = E^Q \sum_{i,j} a_{ij} \mathbf{x}_i \mathbf{x}_j = \sum_i \sum_j a_{ij} s_{ji} = \text{Tr } AS,$$

so

$$E^Q \log \frac{dQ}{dP} = -\frac{1}{2} \log \det R^{-1} S + \frac{1}{2} \text{Tr}(R^{-1} S - I).$$

The same applies to $E^P \log \frac{dP}{dQ}$ so the entropy is:

$$H = \frac{1}{2} \text{Tr}(R^{-1} S + S^{-1} R - 2I).$$

The claim follows from $\text{Tr } AB = \text{Tr } BA$. □

Let K be a root of R_n^τ as above and put

$$(2) \quad T \stackrel{d}{=} K^{-1} S K^{-*}.$$

Then T is symmetric, positive definite, and $T^{-1} = K^* S^{-1} K$, so from claim 2.1,

$$H_n^\tau = \frac{1}{2} \text{Tr}(T + T^{-1} - 2I).$$

In terms of the (positive) eigenvalues $\lambda_i^n = \lambda_i^n(\tau)$, of T :

$$H_n^\tau = \frac{1}{2} \sum_{i=0}^n \frac{(\lambda_i^n - 1)^2}{\lambda_i^n}$$

For $0 < \tau \leq \tau_0$, $H^\tau \leq H \stackrel{d}{=} H^{\tau_0}$, and since $H^\tau = \overline{\lim} H_n^\tau$, it follows that

$$(3) \quad H < \infty \quad \text{only if} \quad \overline{\lim}_n \sum_i (\lambda_i^n - 1)^2 \leq C < \infty,$$

where C is a constant which depends on τ_0 . Note that

$$(4) \quad \sum_{i=0}^n (\lambda_i^n - 1)^2 = \text{Tr} (T - I)^2 = \sum_{i \neq j} t_{ij}^2 + \sum_i (t_{ii} - 1)^2.$$

Let $\delta = \delta_n^\tau = 2\tau/n$ be the mesh of the partition. Returning to the case of $R = t \wedge s$ and a stationary correlation $S = S(t - s)$, we choose

$$K \stackrel{d}{=} \begin{pmatrix} \sqrt{1-\tau} & 0 & 0 & 0 & \dots & 0 \\ \sqrt{1-\tau} & \sqrt{\delta} & 0 & 0 & \dots & 0 \\ \sqrt{1-\tau} & \sqrt{\delta} & \sqrt{\delta} & 0 & \dots & 0 \\ \vdots & \vdots & & \ddots & & \\ \sqrt{1-\tau} & \sqrt{\delta} & \sqrt{\delta} & \dots & \sqrt{\delta} & 0 \\ \sqrt{1-\tau} & \sqrt{\delta} & \sqrt{\delta} & \dots & \sqrt{\delta} & \sqrt{\delta} \end{pmatrix}.$$

It is easy to verify that $R_n^\tau = KK^*$ and that

$$K^{-1} = \begin{pmatrix} \frac{1}{\sqrt{1-\tau}} & 0 & 0 & 0 & \dots & 0 \\ -\frac{1}{\sqrt{\delta}} & \frac{1}{\sqrt{\delta}} & 0 & 0 & \dots & 0 \\ 0 & -\frac{1}{\sqrt{\delta}} & \frac{1}{\sqrt{\delta}} & 0 & \dots & 0 \\ \vdots & & \ddots & & & \\ 0 & 0 & \dots & -\frac{1}{\sqrt{\delta}} & \frac{1}{\sqrt{\delta}} & 0 \\ 0 & 0 & \dots & & -\frac{1}{\sqrt{\delta}} & \frac{1}{\sqrt{\delta}} \end{pmatrix}.$$

It follows that, with $S_k = S(k\delta)$,

$$(5) \quad T = \begin{pmatrix} \frac{S_0}{1-\tau} & \frac{1}{\sqrt{1-\tau}} \frac{S_1-S_0}{\sqrt{\delta}} & \frac{1}{\sqrt{1-\tau}} \frac{S_2-S_1}{\sqrt{\delta}} & \frac{1}{\sqrt{1-\tau}} \frac{S_3-S_2}{\sqrt{\delta}} & \dots & \frac{1}{\sqrt{1-\tau}} \frac{S_n-S_{n-1}}{\sqrt{\delta}} \\ * & 2\frac{S_0-S_1}{\delta} & \frac{2S_1-S_0-S_2}{\delta} & \frac{2S_2-S_1-S_3}{\delta} & \dots & \frac{2S_{n-1}-S_{n-2}-S_n}{\delta} \\ * & * & 2\frac{S_0-S_1}{\delta} & \frac{2S_1-S_0-S_2}{\delta} & \dots & \frac{2S_{n-2}-S_{n-3}-S_{n-1}}{\delta} \\ * & * & * & 2\frac{S_0-S_1}{\delta} & \ddots & \vdots \\ & & & & \ddots & \frac{2S_1-S_0-S_2}{\delta} \\ * & * & * & * & & 2\frac{S_0-S_1}{\delta} \end{pmatrix}$$

where the *'s are filled in according to the symmetry of T (note the Töeplitz $n \times n$ sub-matrix). Using (4) we get:

$$(6) \quad \sum_{i=0}^n (\lambda_i^n - 1)^2 = \left(\frac{S_0}{1-\tau} - 1 \right)^2 + n \left(2\frac{S_0-S_1}{\delta} - 1 \right)^2 \\ + 2\frac{1}{1-\tau} \sum_{k=1}^n \frac{(S_k - S_{k-1})^2}{\delta} \\ + 2 \sum_{k=1}^{n-1} \left(\frac{2S_k - S_{k-1} - S_{k+1}}{\delta} \right)^2 (n-k).$$

Assume now that Q^{τ_0} is absolutely continuous with respect to W^{τ_0} . Using (3) and the last equation, with $\delta = 2\tau_0/n$, we find:

$$(7) \quad M_0 = \sup_{\tau \leq \tau_0} \sup_n n \left(2 \frac{S_0 - S_1}{2\tau/n} - 1 \right)^2 < \infty,$$

$$(8) \quad M_1 = \sup_n \sum_{k=1}^n (S_k - S_{k-1})^2 \frac{1}{\delta} < \infty,$$

and

$$(9) \quad M_2 = \sup_n \sum_{k=1}^{n-1} \left(\frac{2S_k - S_{k-1} - S_{k+1}}{\delta} \right)^2 (n-k) < \infty.$$

It follows from (7) that S is continuous from the right at 0, and since it is symmetric and positive definite, S is also uniformly continuous on its domain (see e.g. [1], p.191).

Claim 2.2. S is an absolutely continuous function.

Proof. Let

$$S_n(r) \stackrel{d}{=} \begin{cases} S(k\delta) & \text{for } k\delta \leq r < (k+1)\delta, \end{cases}$$

and

$$f_n(r) \stackrel{d}{=} \frac{S_n(r+\delta) - S_n(r)}{\delta}.$$

It follows from (8) that, with $\delta = 2\tau_0/n$ as before, $\int_0^{2\tau_0} f_n^2 \leq M_1$, so there exists a subsequence f_{n_k} converging weakly in L^2 to some function f . If for any smooth compactly supported φ on $(0, 2\tau_0)$,

$$(10) \quad \int_0^{2\tau_0} f\varphi = - \int_0^{2\tau_0} S\varphi',$$

then standard Sobolev type arguments show that S is absolutely continuous (and $S' = f$). To prove (10), note that for sufficiently large n (φ being compactly supported),

$$\begin{aligned} \int_0^{2\tau_0} f_{n_k}(t)\varphi(t) dt &= \int_0^{2\tau_0} \frac{S_{n_k}(t+\delta) - S_{n_k}(t)}{\delta} \varphi(t) dt \\ &= - \int_0^{2\tau_0} S_{n_k}(t) \frac{\varphi(t) - \varphi(t-\delta)}{\delta} dt. \end{aligned}$$

Since S is continuous and φ is smooth, by letting $k \rightarrow \infty$ in the last equation we get (10). \square

It is a corollary of the last claim and (7) that $S^+(0) = -\frac{1}{2}$, as in (ii) of the theorem.

Claim 2.3. S' is absolutely continuous on $(0, 2\tau_0)$ and $\int_0^{2\tau_0} S''(t)^2(2\tau_0 - t) dt < \infty$.

Proof. With a slight abuse of notation, define

$$S'_n(r) \stackrel{d}{=} \left\{ \frac{1}{\delta} \int_{k\delta}^{(k+1)\delta} S'(\eta) d\eta \quad \text{for} \quad k\delta \leq r < (k+1)\delta, \right.$$

and let

$$\begin{aligned} g_n(r) &\stackrel{d}{=} \frac{S'_n(r + \delta/2) - S'_n(r - \delta/2)}{\delta} \\ &= \left\{ \frac{S((k+1)\delta) + S((k-1)\delta) - 2S(k\delta)}{\delta^2} \quad \left(k - \frac{1}{2}\right)\delta \leq r < \left(k + \frac{1}{2}\right)\delta. \right. \end{aligned}$$

Since, by (9),

$$\begin{aligned} \int_0^{2\tau_0} g_n(t)^2(2\tau_0 - t) dt &= \\ &\sum \left[\frac{S((k+1)\delta) + S((k-1)\delta) - 2S(k\delta)}{\delta^2} \right]^2 (2\tau_0 - k\delta) \leq M_2, \end{aligned}$$

there exists a subsequence $\{g_{n_k}\}$ that converges weakly to a limit g in $L^2((2\tau_0 - t)dt)$. Hence, for every proper subinterval $I \subset (0, 2\tau_0)$, $\{g_{n_k}\}$ converges weakly in $L^2(I, dt)$. Again,

$$(11) \quad \int_0^{2\tau_0} g\varphi = - \int_0^{2\tau_0} S'\varphi'$$

will show S' is absolutely continuous with $g = S''$. For sufficiently large n ,

$$\int_0^{2\tau_0} g_{n_k}\varphi = - \int_0^{2\tau_0} S'_{n_k}(t) \frac{\varphi(t + \delta/2) - \varphi(t - \delta/2)}{\delta} dt.$$

Since $S'_{n_k} \rightarrow S'$ in L^1 and φ is smooth, we get (11) by letting $k \rightarrow \infty$. Finally, by its definition, $g \in L^2((2\tau_0 - t)dt)$; in particular, we get (i) of theorem 1. \square

This proves half of theorem 1. Assume now that S satisfies (i) and (ii) of theorem 1.

Claim 2.4. For any $\tau \leq \tau_0$,

$$\begin{aligned} \sup_n \sum_i (\lambda_i^n(\tau) - 1)^2 &\leq \left(\frac{S_0}{1-\tau} - 1 \right)^2 + \int_0^{2\tau} S''(t)^2(2\tau - t) dt \\ &\quad + \frac{2}{1-\tau} \int_0^{2\tau} S'(t)^2 dt < \infty. \end{aligned}$$

Proof. Note that, with $T = T_n(\tau) = (t_{ij})_{0 \leq i, j \leq n}$,

$$(12) \quad \sum_{i=1}^n t_{i0}^2 + t_{0i}^2 = \frac{2}{1-\tau} \sum_{i=1}^n (S_i - S_{i-1})^2 \frac{1}{\delta} \leq \frac{2}{1-\tau} \int_0^{2\tau} S'(t)^2 dt.$$

Let

$$\bar{S}(r) \stackrel{d}{=} S(r) - \left(1 - \frac{|r|}{2}\right).$$

Then \bar{S} is an even function, with absolutely continuous derivative \bar{S}' on $(-2\tau_0, 2\tau_0)$, and $\bar{S}'' \equiv S''$ on $(0, 2\tau_0)$. It is not hard to verify that

$$(13) \quad \begin{aligned} t_{ij} &= -\frac{1}{\delta} \int_{(i-1)\delta}^{i\delta} \int_{(j-1)\delta}^{j\delta} \bar{S}''(t-s) dt ds & 1 \leq i \neq j \leq n, \\ t_{ii} - 1 &= -\frac{1}{\delta} \int_{(i-1)\delta}^{i\delta} \int_{(i-1)\delta}^{i\delta} \bar{S}''(t-s) dt ds & 1 \leq i \leq n. \end{aligned}$$

Thus, for $\tau \leq \tau_0$

$$\begin{aligned} \left[\sum_{1 \leq i \neq j \leq n} t_{ij}^2 + \sum_{i=1}^n (t_{ii} - 1)^2 \right] &\leq \int_0^{2\tau} \int_0^{2\tau} \bar{S}''(t-s)^2 dt ds \\ &= \int_0^{2\tau} S''(t)^2 (2\tau - t) dt < \infty, \end{aligned}$$

which together with (12) and (4) completes the proof of claim 2.4. \square

We next show that for $\tau < \tau_0$, $\inf_{i,n} \lambda_i^n(\tau) > 0$. It follows that,

$$H^\tau = \overline{\lim} \sum \frac{(\lambda_i^n - 1)^2}{\lambda_i^n} \leq \frac{1}{\inf_{i,n} \lambda_i^n(\tau)} \overline{\lim} \sum (\lambda_i^n - 1)^2 < \infty,$$

which completes the proof of theorem 1.

Since $\bar{S}'' \in L^2([0, 2\tau] \times [0, 2\tau])$, it defines a compact integral operator on $L^2(0, 2\tau)$:

$$(\bar{S}'' * f)(r) \stackrel{d}{=} \int_0^{2\tau} \bar{S}''(r-t) f(t) dt.$$

Let $\mathcal{H} = \mathcal{H}_\tau \stackrel{d}{=} \mathbb{R} \oplus L^2(0, 2\tau)$. Define a bounded symmetric operator $\mathcal{T}_\tau : (\beta, f) \in \mathcal{H} \mapsto \mathcal{H}$ as follows:

$$\mathcal{T}(\beta, f) \stackrel{d}{=} \left(\beta \frac{S(0)}{1-\tau} + \frac{1}{\sqrt{1-\tau}} \int_0^{2\tau} S'(t) f(t) dt, \frac{\beta}{\sqrt{1-\tau}} S' + f - \bar{S}'' * f \right).$$

Let U_n be the n dimensional subspace of $L^2(0, 2\tau)$ populated by functions which are constant on the intervals $[i\delta, (i+1)\delta)$, $\delta = 2\tau/n$. For $f \in U_n$ and $X \in C[0, 2\tau]$, we define $\int_0^{2\tau} f(t) dX_t$ in the obvious

manner. Let $\hat{\cdot}: U_n \mapsto \mathbb{R}^n$ be the natural isometry between these spaces, i.e., for

$$f(t) = \sum_{i=1}^n f_i 1_{[(i-1)\delta, i\delta)}(t) \in U_n,$$

$$\hat{f} \stackrel{d}{=} (f_1\sqrt{\delta}, f_2\sqrt{\delta}, \dots, f_n\sqrt{\delta}).$$

Let Q be a stationary extension of Q^{τ_0} to $C(\mathbb{R})$ corresponding to a positive-definite extension of S from $[-2\tau_0, 2\tau_0]$ to \mathbb{R} . A straight forward computation shows that for $f \in U_n$ and $\beta \in \mathbb{R}$,

$$\left\langle T(\beta, \hat{f}), (\beta, \hat{f}) \right\rangle_{\mathbb{R}^{n+1}} = E^Q \left[\frac{\beta}{\sqrt{1-\tau}} X_0 + \int_0^{2\tau} f(t) dX_t \right]^2.$$

Remark. The last equation might deserve a second look: on the left hand side we have T which was obtained from S and R , while on the right hand side R does not appear explicitly.

It follows from (13) that,

$$\left\langle T(\beta, \hat{f}), (\beta, \hat{f}) \right\rangle_{\mathbb{R}^{n+1}} = \left\langle \mathcal{T}(\beta, f), (\beta, f) \right\rangle_{\mathcal{H}}.$$

Thus, for $\beta = 0$ and $f \in U_n$,

$$E^Q \left[\int_0^{2\tau} f(t) dX_t \right]^2 = \left\langle \mathcal{T}(0, f), (0, f) \right\rangle_{\mathcal{H}} \leq \|\mathcal{T}\| \|f\|_{L^2}^2.$$

Hence the map

$$f \mapsto \int_0^{2\tau} f(t) dX_t,$$

defined initially on $\bigcup U_n$, can be extended uniquely as a bounded linear map from $L^2(0, 2\tau)$ into $L^2(\Omega, dQ)$, where $\Omega = C_0$ is our probability space. Furthermore, by continuity, for *any* $f \in L^2(0, 2\tau)$ and $\beta \in \mathbb{R}$,

$$(14) \quad \left\langle \mathcal{T}(\beta, f), (\beta, f) \right\rangle_{\mathcal{H}} = E^Q \left[\frac{\beta}{\sqrt{1-\tau}} X_0 + \int_0^{2\tau} f(t) dX_t \right]^2.$$

In particular, \mathcal{T} is positive-definite.

Claim 2.5. Suppose that $\inf_{i,n} \lambda_i^n(\tau) = 0$, then there exists $\beta \in \mathbb{R}$ and $f \in L^2(0, 2\tau)$, not identically 0, with

$$(15) \quad \frac{\beta}{\sqrt{1-\tau}} X_0 + \int_0^{2\tau} f(t) dX_t = 0 \quad \text{a.s. } dQ,$$

Proof. Assuming the eigenvalues of T_n satisfy $\lambda_0^n(\tau) \leq \lambda_1^n \leq \dots \leq \lambda_n^n$, there exists a sub-sequence n_k such that

$$\nu_k \stackrel{d}{=} \lambda_0^{n_k} \longrightarrow 0.$$

Let $\mathbf{v}_k = (\mathbf{v}_k(0), \mathbf{v}_k(1), \dots, \mathbf{v}_k(n_k)) \in \mathbb{R}^{n_k+1}$ be the corresponding normalized eigenvectors, i.e.,

$$T_{n_k} \mathbf{v}_k = \nu_k \mathbf{v}_k \quad , \quad |\mathbf{v}_k| = 1.$$

Let $\beta_k = \mathbf{v}_k(0)$ and let

$$f_k \stackrel{d}{=} (\mathbf{v}_k(1), \widehat{\dots}, \mathbf{v}_k(n_k)).$$

Then $\|(\beta_k, f_k)\|_{\mathcal{H}} = 1$, and without loss of generality you may assume that

$$(\beta_k, f_k) \longrightarrow (\beta, f) \quad \text{weakly in } \mathcal{H}.$$

Let \mathcal{I} be the identity in \mathcal{H} . The operator $\mathcal{T}_0 \stackrel{d}{=} \mathcal{T} - \mathcal{I}$ is compact, by inspection, so

$$\mathcal{T}_0(\beta_k, f_k) \longrightarrow \mathcal{T}_0(\beta, f) \quad \text{strongly in } \mathcal{H}.$$

It follows that

$$\nu_k - 1 = \left\langle \mathcal{T}_0(\beta_k, f_k), (\beta_k, f_k) \right\rangle_{\mathcal{H}} \longrightarrow \left\langle \mathcal{T}_0(\beta, f), (\beta, f) \right\rangle_{\mathcal{H}},$$

or

$$\left\langle \mathcal{T}_0(\beta, f), (\beta, f) \right\rangle_{\mathcal{H}} = -1.$$

But \mathcal{T} is symmetric and positive-definite, so the spectrum $\sigma(\mathcal{T}_0) \subset [-1, \infty)$, whence $\|(\beta, f)\|_{\mathcal{H}} = 1$ and

$$\left\langle \mathcal{T}(\beta, f), (\beta, f) \right\rangle_{\mathcal{H}} = 0.$$

The claim now follows from (14). □

Remarks:

- In hind sight (β_k, f_k) converge strongly to (β, f) .
- S has a unique extension as a symmetric positive-definite function from $[-2\tau, 2\tau]$ to \mathbb{R} . Indeed, let Δ be any spectral distribution function corresponding to such an extension of S . Using the standard isometry between $L^2(\mathbb{R}, \Delta)$ and $L^2(\Omega, Q)$, defined by

$$e^{it\omega} \longleftrightarrow X_t,$$

it follows from (15) that

$$\frac{\beta}{\sqrt{1-\tau}} + i\omega \int_0^{2\tau} f(t)e^{i\omega t} dt = 0,$$

where the equality is in $L^2(\Delta)$. Thus, there exists a non-trivial holomorphic function of exponential type $\leq \tau$, which vanishes in $L^2(\Delta)$. Krein's alternative says that in this case the stationary process governed by any extension of S must be completely predictable from any interval of length 2τ [2][see sec. 4.8]. This implies that there exists only one extension ([4] is helpful).

Claim 2.6. If (15) holds with a non-trivial $f \in L^2(0, 2\tau)$, then there exists $\tau' \in (\tau, \tau_0)$, such that the eigenspace,

$$V \stackrel{d}{=} \left\{ g \in L^2(0, 2\tau') : \int_0^{2\tau'} \bar{S}''(t-r)g(r) dr = g(t) \right\},$$

is infinite-dimensional.

Proof. Since Q is stationary, it follows from (15) that for any $\delta > 0$,

$$(16) \quad \frac{\beta}{\sqrt{1-\tau}} X_{2\delta} + \int_0^{2\tau} f(t) dX_{t+2\delta} = 0 \quad \text{a.s. } dQ.$$

Let

$$g(t) \stackrel{d}{=} \begin{cases} \frac{\beta}{\sqrt{1-\tau}} - f(t) & 0 \leq t < 2\delta \\ f(t-2\delta) - f(t) & 2\delta \leq t < 2\tau \\ f(t-2\delta) & 2\tau \leq t < 2\tau + 2\delta \\ 0 & \text{elsewhere.} \end{cases}$$

Subtracting (15) from (16), we find

$$\int_0^{2(\tau+\delta)} g(t) dX_t = 0 \quad \text{a.s. } dQ.$$

Fix $\tau' \in (\tau, \tau_0)$ and choose $\delta > 0$ so small that $\tau + \delta < \tau'$. By (14),

$$\left\langle \mathcal{T}_{\tau'}(0, g), (0, g) \right\rangle_{\mathcal{H}_{\tau'}} = E^Q \left[\int_0^{2\tau'} g(t) dX_t \right]^2 = 0.$$

For any $\alpha > 0$ such that $\tau + \delta + \alpha < \tau'$, let $g_\alpha(t) \stackrel{d}{=} g(t - \alpha)$. By stationarity,

$$\left\langle \mathcal{T}_{\tau'}(0, g_\alpha), (0, g_\alpha) \right\rangle_{\mathcal{H}_{\tau'}} = E^Q \left[\int_0^{2\tau'} g(t - \alpha) dX_t \right]^2 = 0.$$

But $\mathcal{T}_{\tau'}$ is symmetric and positive-definite, so

$$\mathcal{T}_{\tau'}(0, g_\alpha) = 0,$$

which implies

$$g_\alpha = \bar{S}'' * g_\alpha.$$

These α -translates of g span an infinite dimensional subspace $\subset V$. \square

The next lemma shows that V cannot be continuously embedded in $L^\infty(0, 2\tau')$.

Lemma 2.7. *Let U be an infinite-dimensional subspace of $L^2(0, T)$, with $T < \infty$. Then U cannot be continuously embedded in $L^\infty(0, T)$.*

Remarks:

- Undoubtedly, this lemma is known. However having no citation to provide, we give here a proof.
- $T < \infty$ is crucial here. Assume without loss of generality that $T = 1$.

Proof. Let $\{\varphi_n\}$ be an orthonormal set in U . If $M \stackrel{d}{=} \sup_n \|\varphi_n\|_\infty$ is infinite, we are done, so assume $M < \infty$. Let

$$A_n \stackrel{d}{=} \left\{x \in [0, T] : \varphi_n(x) > \frac{1}{\sqrt{2}}\right\},$$

$$B_n \stackrel{d}{=} \left\{x \in [0, T] : \varphi_n(x) < -\frac{1}{\sqrt{2}}\right\}.$$

Let $\lambda(A)$ be the Lebesgue measure of A . Then

$$\lambda(A_n \cup B_n) \geq \frac{1}{2M^2} = 2\alpha > 0.$$

We can assume, without loss of generality, that $\lambda(A_n) \geq \alpha$ for all n . Let $N \stackrel{d}{=} \lceil n/\alpha + 1 \rceil$ and let

$$\varphi(x) \stackrel{d}{=} \sum_{k=1}^N 1_{A_k}(x).$$

Then

$$\int_0^1 \varphi(x) dx \geq N\alpha.$$

Necessarily,

$$\lambda\left(\left\{x : \sum_{k=1}^N 1_{A_k} \geq n\right\}\right) > 0,$$

whence there are n indices, k_1, \dots, k_n , such that

$$\lambda(A_{k_1} \cap A_{k_2} \cap \dots \cap A_{k_n}) > 0.$$

Now

$$\left\| \sum_{j=1}^n \frac{1}{\sqrt{n}} \varphi_{k_j} \right\|_{\infty} > \sqrt{\frac{n}{2}} \quad \text{but} \quad \left\| \sum_{j=1}^n \frac{1}{\sqrt{n}} \varphi_{k_j} \right\|_2 = 1,$$

which completes the proof. \square

The proof that $\inf_{i,n} \lambda_i^n(\tau) > 0$ is now complete: else, by claims 2.5 and 2.6 there exists $\tau' \in (\tau, \tau_0)$ and an infinite dimensional eigenspace V as described in claim 2.6. Hence by the last lemma, there exist a sequence $f_n \in V$ such that,

$$\|f_n\|_{L^2} = 1, \quad f_n = \bar{S}'' * f_n \quad \text{and} \quad \|f_n\|_{\infty} > n.$$

But this implies $\bar{S}'' \notin L^2(0, 2\tau')$, contradicting the assumption that (i) holds.

REFERENCES

- [1] Akhiezer, N.I. (1965). The Classical Moment Problem and Some Related Questions in Analysis. Hafner Publishing Company.
- [2] Dym, H.; McKean, H. P. (1976). Gaussian Processes, Function Theory, and the Inverse Spectral Problem. Academic Press.
- [3] Hida, T.; Hitsuda, M. (1993). Gaussian Processes. AMS.
- [4] Krein, M.G. (1940). On the continuation problem for Hermitian-positive continuous functions. Dokl. Akad. Nauk SSSR **26** 17-21.
- [5] Shepp, L. A. (1966). Radon-Nikodym derivatives of Gaussian measures. Ann. Math. Stat. **37** 321-354.

APPLIED MATH, CALTECH 217-50, CA 91125
E-mail address: keich@ama.caltech.edu