

What's It All About?

- Continuous mathematics—*calculus*—considers objects that vary continuously
 - distance from the wall
- Discrete mathematics considers *discrete* objects, that come in *discrete* bundles
 - number of babies: can't have 1.2

The mathematical techniques for discrete mathematics differ from those for continuous mathematics:

- counting/combinatorics
- number theory
- probability
- logic

We'll be studying these techniques in this course.

Why is it computer science?

This is basically a mathematics course:

- no programming
- lots of theorems to prove

So why is it computer science?

Discrete mathematics is the mathematics underlying almost all of computer science:

- Designing high-speed networks
- Finding good algorithms for sorting
- Doing good web searches
- Analysis of algorithms
- Proving algorithms correct

This Course

We will be focusing on:

- Tools for discrete mathematics:
 - computational number theory (handouts)
 - * the mathematics behind the RSA cryptosystems
 - a little graph theory (Chapter 3)
 - counting/combinatorics (Chapter 4)
 - probability (Chapter 6)
 - * randomized algorithms for primality testing, routing
 - logic (Chapter 7)
 - * how do you *prove* a program is correct
- Tools for proving things:
 - induction (Chapter 2)
 - (to a lesser extent) recursion

First, some background you'll need but may not have . . .

Sets

You need to be comfortable with set notation:

$$S = \{m \mid 2 \leq m \leq 100, m \text{ is an integer}\}$$

S is

the set of

all m

such that

m is between 2 and 100

and

m is an integer.

Important Sets

(More notation you need to know and love ...)

- N (occasionally \mathbb{N}): the nonnegative integers $\{0, 1, 2, 3, \dots\}$
- N^+ : the positive integers $\{1, 2, 3, \dots\}$
- Z : all integers $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
- Q : the rational numbers $\{a/b : a, b \in Z, b \neq 0\}$
- R : the real numbers
- Q^+, R^+ : the positive rationals/reals

Set Notation

- $|S|$ = *cardinality of* (number of elements in) S
 - $|\{a, b, c\}| = 3$
- **Subset:** $A \subset B$ if every element of A is an element of B
 - Note: Lots of people (including me, but not the authors of the text) usually write $A \subset B$ only if A is a *strict* or *proper* subset of B (i.e., $A \neq B$). I write $A \subseteq B$ if $A = B$ is possible.
- Power set: $\mathcal{P}(S)$ is the set of all subsets of S (sometimes denoted 2^S).
 - E.g., $\mathcal{P}(\{1, 2, 3\}) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$
 - $|\mathcal{P}(S)| = 2^{|S|}$

Set Operations

- **Union:** $S \cup T$ is the set of all elements in S or T
 - $S \cup T = \{x | x \in S \text{ or } x \in T\}$
 - $\{1, 2, 3\} \cup \{3, 4, 5\} = \{1, 2, 3, 4, 5\}$
- **Intersection:** $S \cap T$ is the set of all elements in both S and T
 - $S \cap T = \{x | x \in S, x \in T\}$
 - $\{1, 2, 3\} \cap \{3, 4, 5\} = \{3\}$
- **Set Difference:** $S - T$ is the set of all elements in S not in T
 - $S - T = \{x | x \in S, x \notin T\}$
 - $\{3, 4, 5\} - \{1, 2, 3\} = \{4, 5\}$
- **Complementation:** \overline{S} is the set of elements not in S
 - What is $\overline{\{1, 2, 3\}}$?
 - Complementation doesn't make sense unless there is a *universe*, the set of elements we want to consider.
 - If U is the universe, $\overline{S} = \{x | x \in U, x \notin S\}$
 - $\overline{\overline{S}} = U - S$.

Venn Diagrams

Sometimes a picture is worth a thousand words (at least if we don't have too many sets involved).

A Connection

Lemma: For all sets S and T , we have

$$S = (S \cap T) \cup (S - T)$$

Proof: We'll show (1) $S \subset (S \cap T) \cup (S - T)$ and (2) $(S \cap T) \cup (S - T) \subset S$.

For (1), suppose $x \in S$. Either

(a) $x \in T$ or (b) $x \notin T$.

If (a) holds, then $x \in S \cap T$.

If (b) holds, then $x \in S - T$.

In either case, $x \in (S \cap T) \cup (S - T)$.

Since this is true for all $x \in S$, we have (1).

For (2), suppose $x \in (S \cap T) \cup (S - T)$. Thus, either (a) $x \in (S \cap T)$ or $x \in (S - T)$. Either way, $x \in S$.

Since this is true for all $x \in (S \cap T) \cup (S - T)$, we have (2).

Two Important Morals

1. One way to show $S = T$ is to show $S \subset T$ and $T \subset S$.
2. One way to show $S \subset T$ is to show that for every $x \in S$, x is also in T .

Relations

- **Cartesian product:**

$$S \times T = \{(s, t) : s \in S, t \in T\}$$

- $\{1, 2, 3\} \times \{3, 4\} = \{(1, 3), (2, 3), (3, 3), (1, 4), (2, 4), (3, 4)\}$
- $|S \times T| = |S| \times |T|$.

- A *relation* on S and T (or, on $S \times T$) is a subset of $S \times T$

- A *relation* on S is a subset of $S \times S$

- *Taller than* is a relation on people: (Joe, Sam) is in the Taller than relation if Joe is Taller than Sam
- *Larger than* is a relation on R :

$$L = \{(x, y) | x, y \in R, x > y\}$$

- *Divisibility* is a relation on N :

$$D = \{(x, y) | x, y \in N, x|y\}$$

Reflexivity, Symmetry, Transitivity

- A relation R on S is *reflexive* if $(x, x) \in R$ for all $x \in S$.
 - \leq is reflexive; $<$ is not
- A relation R on S is *symmetric* if $(x, y) \in R$ implies $(y, x) \in R$.
 - “sibling-of” is symmetric (what about “sister of”)
 - \leq is not symmetric
- A relation R on S is *transitive* if $(x, y) \in R$ and $(y, z) \in R$ implies $(x, z) \in R$.
 - $\leq, <, \geq, >$ are all transitive;
 - “parent-of” is not transitive; “ancestor-of” is

Pictorially, we have:

Transitive Closure

[[NOT DISCUSSED ENOUGH IN THE TEXT]]

The *transitive closure* of a relation R is the least relation R^* such that

1. $R \subset R^*$
2. R^* is transitive (so that if $(u, v), (v, w) \in R^*$, then so is (u, w)).

Example: Suppose $R = \{(1, 2), (2, 3), (1, 4)\}$.

- $R^* = \{(1, 2), (1, 3), (2, 3), (1, 4)\}$
- we need to add $(1, 3)$, because $(1, 2), (2, 3) \in R$

Note that we don't need to add $(2, 4)$.

- If $(2, 1), (1, 4)$ were in R , then we'd need $(2, 4)$
- $(1, 2), (1, 4)$ doesn't force us to add anything (it doesn't fit the "pattern" of transitivity).

Note that if R is already transitive, then $R^* = R$.

Equivalence Relations

- A relation R is an *equivalence relation* if it is reflexive, symmetric, and transitive
 - $=$ is an equivalence relation
 - *Parity* is an equivalence relation on N ;
 $(x, y) \in \text{Parity}$ if $x - y$ is even

Functions

We think of a function $f : S \rightarrow T$ as providing a mapping from S to T . But ...

Formally, a *function* is a relation R on $S \times T$ such that for each $s \in S$, there is a unique $t \in T$ such that $(s, t) \in R$.

If $f : S \rightarrow T$, then S is the *domain* of f , T is the *range*; $\{y : f(x) = y \text{ for some } x \in S\}$ is the *image*.

We often think of a function as being characterized by an algebraic formula

- $y = 3x - 2$ characterizes $f(x) = 3x - 2$.

It ain't necessarily so.

- Some formulas don't characterize functions:
 - $x^2 + y^2 = 1$ defines a circle; no unique y for each x
- Some functions can't be characterized by algebraic formulas
 - $f(n) = \begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \text{ is odd} \end{cases}$

Function Terminology

Suppose $f : S \rightarrow T$

- f is *onto* (or *surjective*) if, for each $t \in T$, there is some $s \in S$ such that $f(s) = t$.

- if $f : R^+ \rightarrow R^+$, $f(x) = x^2$, then f is onto
- if $f : R \rightarrow R$, $f(x) = x^2$, then f is *not* onto
- f is *one-to-one* (1-1, *injective*) if it is not the case that $s \neq s'$ and $f(s) = f(s')$.

- if $f : R^+ \rightarrow R^+$, $f(x) = x^2$, then f is 1-1
- if $f : R \rightarrow R$, $f(x) = x^2$, then f is *not* 1-1.

17

- a function is *bijective* if it is 1-1 and onto.

- if $f : R^+ \rightarrow R^+$, $f(x) = x^2$, then f is bijective
- if $f : R \rightarrow R$, $f(x) = x^2$, then f is *not* bijective.

If $f : S \rightarrow T$ is bijective, then $|S| = |T|$.

18

Inverse Functions

If $f : S \rightarrow T$, then f^{-1} maps an element in the range of f to all the elements that are mapped to it by f .

$$f^{-1}(t) = \{s | f(s) = t\}$$

- if $f(2) = 3$, then $2 \in f^{-1}(3)$.

f^{-1} is not a function from $\text{range}(f)$ to S .

It is a function if f is one-to-one.

- In this case, $f^{-1}(f(x)) = x$.

19

Functions You Should Know (and Love)

- *Absolute value*: Domain = R ; Range = $\{0\} \cup R^+$

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

- $|3| = |-3| = 3$

- *Floor function*: Domain = R ; Range = Z

$$\lfloor x \rfloor = \text{largest integer not greater than } x$$

- $\lfloor 3.2 \rfloor = 3$; $\lfloor \sqrt{3} \rfloor = 1$; $\lfloor -2.5 \rfloor = -3$

- *Ceiling function*: Domain = R ; Range = Z

$$\lceil x \rceil = \text{smallest integer not less than } x$$

- $\lceil 3.2 \rceil = 4$; $\lceil \sqrt{3} \rceil = 2$; $\lceil -2.5 \rceil = -2$

- *Factorial function*: Domain = Range = N

$$n! = n(n-1)(n-2)\dots 3 \times 2 \times 1$$

- $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$

- By convention, $0! = 1$

20

Exponents

Exponential with base a : Domain = R , Range = R^+

$$f(x) = a^x$$

- Note: a , the *base*, is fixed; x varies
- You probably know: $a^n = a \times \cdots \times a$ (n times)

How do we define $f(x)$ if x is not a positive integer?

- **Want:** (1) $a^{x+y} = a^x a^y$; (2) $a^1 = a$

This means

- $a^2 = a^{1+1} = a^1 a^1 = a \times a$
- $a^3 = a^{2+1} = a^2 a^1 = a \times a \times a$
- \dots
- $a^n = a \times \cdots \times a$ (n times)

We get more:

- $a = a^1 = a^{1+0} = a \times a^0$
 - Therefore $a^0 = 1$
- $1 = a^0 = a^{b+(-b)} = a^b \times a^{-b}$
 - Therefore $a^{-b} = 1/a^b$

21

- $a = a^1 = a^{\frac{1}{2}+\frac{1}{2}} = a^{\frac{1}{2}} \times a^{\frac{1}{2}} = (a^{\frac{1}{2}})^2$
 - Therefore $a^{\frac{1}{2}} = \sqrt{a}$
- Similar arguments show that $a^{\frac{1}{k}} = \sqrt[k]{a}$
- $a^{mx} = a^x \times \cdots \times a^x$ (m times) $= (a^x)^m$
 - Thus, $a^{\frac{m}{n}} = (a^{\frac{1}{n}})^m = (\sqrt[n]{a})^m$.

This determines a^x for all x rational. The rest follows by continuity.

22

Computing a^n quickly

What's the best way to compute a^{1000} ?

One way: multiply $a \times a \times a \times a \dots$

- This requires 999 multiplications.

Can we do better?

How many multiplications are needed to compute:

- a^2
- a^4
- a^8
- a^{16}
- \dots

Write 1000 in binary: 1111101000

- How many multiplications are needed to calculate a^{1000} ?

23

Logarithms

Logarithm base a : Domain = R^+ ; Range = R

$$y = \log_a(x) \Leftrightarrow a^y = x$$

- $\log_2(8) = 3$; $\log_2(16) = 4$; $3 < \log_2(15) < 4$

The key properties of the log function follow from those for the exponential:

1. $\log_a(1) = 0$ (because $a^0 = 1$)
2. $\log_a(a) = 1$ (because $a^1 = a$)
3. $\log_a(xy) = \log_a(x) + \log_a(y)$

Proof: Suppose $\log_a(x) = z_1$ and $\log_a(y) = z_2$.

Then $a^{z_1} = x$ and $a^{z_2} = y$.

Therefore $xy = a^{z_1} \times a^{z_2} = a^{z_1+z_2}$.

Thus $\log_a(xy) = z_1 + z_2 = \log_a(x) + \log_a(y)$.

4. $\log_a(x^r) = r \log_a(x)$
5. $\log_a(1/x) = -\log_a(x)$ (because $a^{-y} = 1/a^y$)
6. $\log_b(x) = \log_a(x) / \log_a(b)$

24

Examples:

- $\log_2(1/4) = -\log_2(4) = -2$.
- $\log_2(-4)$ undefined
- - $\log_2(2^{10}3^5)$
 - $= \log_2(2^{10}) + \log_2(3^5)$
 - $= 10 \log_2(2) + 5 \log_2(3)$
 - $= 10 + 5 \log_2(3)$

25

Limit Properties of the Log Function

$$\lim_{x \rightarrow \infty} \log(x) = \infty$$
$$\lim_{x \rightarrow \infty} \frac{\log(x)}{x} = 0$$

As x gets large $\log(x)$ grows without bound.

But x grows MUCH faster than $\log(x)$.

In fact, $\lim_{x \rightarrow \infty} (\log(x)^m)/x = 0$

26

Polynomials

$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_kx^k$ is a *polynomial* function.

- a_0, \dots, a_k are the *coefficients*

You need to know how to multiply polynomials:

$$\begin{aligned} & (2x^3 + 3x)(x^2 + 3x + 1) \\ &= 2x^3(x^2 + 3x + 1) + 3x(x^2 + 3x + 1) \\ &= 2x^5 + 6x^4 + 2x^3 + 3x^3 + 9x^2 + 3x \\ &= 2x^5 + 6x^4 + 5x^3 + 9x^2 + 3x \end{aligned}$$

Exponentials grow MUCH faster than polynomials:

$$\lim_{x \rightarrow \infty} \frac{a_0 + \cdots + a_kx^k}{b^x} = 0 \text{ if } b > 1$$

27

Why Rates of Growth Matter

Suppose you want to design an algorithm to do sorting.

- The naive algorithm takes time $n^2/4$ on average to sort n items
- A more sophisticated algorithm takes time $2n \log(n)$

Which is better?

$$\lim_{n \rightarrow \infty} (2n \log(n)/(n^2/4)) = \lim_{n \rightarrow \infty} (8 \log(n)/n) = 0$$

For example,

- if $n = 1,000,000$, $2n \log(n) = 40,000,000$ — this is doable
- $n^2/4 = 250,000,000,000$ — this is not doable

Algorithms that take exponential time are hopeless on large datasets.

28

Sum and Product Notation

$$\sum_{i=0}^k a_i x^i = a_0 + a_1 x + a_2 x^2 + \cdots + a_k x^k$$

$$\sum_{i=2}^5 i^2 = 2^2 + 3^2 + 4^2 + 5^2 = 54$$

Can limit the set of values taken on by the *index* i :

$$\sum_{\{i: 2 \leq i \leq 8 | i \text{ even}\}} a_i = a_2 + a_4 + a_6 + a_8$$

Can have double sums:

$$\begin{aligned} & \sum_{i=1}^2 \sum_{j=0}^3 a_{ij} \\ &= \sum_{i=1}^2 (\sum_{j=0}^3 a_{ij}) \\ &= \sum_{j=0}^3 a_{1j} + \sum_{j=0}^3 a_{2j} \\ &= a_{10} + a_{11} + a_{12} + a_{13} + a_{20} + a_{21} + a_{22} + a_{23} \end{aligned}$$

Product notation similar:

$$\prod_{i=0}^k a_i = a_0 a_1 \cdots a_k$$

Changing the Limits of Summation

This is like changing the limits of integration.

$$\bullet \sum_{i=1}^{n+1} a_i = \sum_{i=0}^n a_{i+1} = a_1 + \cdots + a_{n+1}$$

Steps:

- Start with $\sum_{i=1}^{n+1} a_i$.
- Let $j = i - 1$. Thus, $i = j + 1$.
- Rewrite limits in terms of j : $i = 1 \rightarrow j = 0$; $i = n + 1 \rightarrow j = n$
- Rewrite body in terms of $a_i \rightarrow a_{j+1}$
- Get $\sum_{j=0}^n a_{j+1}$
- Now replace j by i (j is a dummy variable). Get

$$\sum_{i=0}^n a_{i+1}$$

Matrix Algebra

An $m \times n$ *matrix* is a two-dimensional array of numbers, with m rows and n columns:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- A $1 \times n$ matrix $[a_1 \dots a_n]$ is a *row vector*.
- An $m \times 1$ matrix is a *column vector*.

We can add two $m \times n$ matrices:

- If $A = [a_{ij}]$ and $B = [b_{ij}]$ then $A + B = [a_{ij} + b_{ij}]$.

$$\begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} + \begin{bmatrix} 3 & 7 \\ 4 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 10 \\ 9 & 9 \end{bmatrix}$$

Another important operation: *transposition*.

- If we transpose an $m \times n$ matrix, we get an $n \times m$ matrix by switching the rows and columns.

$$\begin{bmatrix} 2 & 3 & 9 \\ 5 & 7 & 12 \end{bmatrix}^T = \begin{bmatrix} 2 & 5 \\ 3 & 7 \\ 9 & 12 \end{bmatrix}$$