

How to Guess What to Prove

Sometimes formulating $P(n)$ is straightforward; sometimes it's not. This is what to do:

- Compute the result in some specific cases
- Conjecture a generalization based on these cases
- Prove the correctness of your conjecture (by induction)

Example

Suppose $a_1 = 1$ and $a_n = a_{\lceil n/2 \rceil} + a_{\lfloor n/2 \rfloor}$ for $n > 1$. Find an explicit formula for a_n .

Try to see the pattern:

- $a_1 = 1$
- $a_2 = a_1 + a_1 = 1 + 1 = 2$
- $a_3 = a_2 + a_1 = 2 + 1 = 3$
- $a_4 = a_2 + a_2 = 2 + 2 = 4$

Suppose we modify the example. Now $a_1 = 3$ and $a_n = a_{\lceil n/2 \rceil} + a_{\lfloor n/2 \rfloor}$ for $n > 1$. What's the pattern?

- $a_1 = 3$
- $a_2 = a_1 + a_1 = 3 + 3 = 6$
- $a_3 = a_2 + a_1 = 6 + 3 = 9$
- $a_4 = a_2 + a_2 = 6 + 6 = 12$

$$a_n = 3n!$$

Theorem: If $a_1 = k$ and $a_n = a_{\lceil n/2 \rceil} + a_{\lfloor n/2 \rfloor}$ for $n > 1$, then $a_n = kn$ for $n \geq 1$.

Proof: By strong induction. Let $P(n)$ be the statement that $a_n = kn$.

Basis: $P(1)$ says that $a_1 = k$, which is true by hypothesis.

Inductive step: Assume $P(1), \dots, P(n)$; prove $P(n+1)$.

$$\begin{aligned} a_{n+1} &= a_{\lceil (n+1)/2 \rceil} + a_{\lfloor (n+1)/2 \rfloor} \\ &= k\lceil (n+1)/2 \rceil + k\lfloor (n+1)/2 \rfloor \text{ [Induction hypothesis]} \\ &= k(\lceil (n+1)/2 \rceil + \lfloor (n+1)/2 \rfloor) \\ &= k(n+1) \end{aligned}$$

We used the fact that $\lceil n/2 \rceil + \lfloor n/2 \rfloor = n$ for all n (in particular, for $n+1$). To see this, consider two cases: n is odd and n is even.

- if n is even, $\lceil n/2 \rceil + \lfloor n/2 \rfloor = n/2 + n/2 = n$
- if n is odd, suppose $n = 2k + 1$
 - $\lceil n/2 \rceil + \lfloor n/2 \rfloor = (k+1) + k = 2k + 1 = n$

This proof has a (small) gap:

- We should check that $\lceil (n+1)/2 \rceil \leq n$

In general, there is no rule for guessing the right inductive hypothesis. However, if you have a sequence of numbers

$$r_1, r_2, r_3, \dots$$

and want to guess a general expression, here are some guidelines for trying to find the *type* of the expression (exponential, polynomial):

- Compute $\lim_{n \rightarrow \infty} r_{n+1}/r_n$
 - if it looks like $\lim_{n \rightarrow \infty} r_{n+1}/r_n = b \notin \{0, 1\}$, then r_n probably has the form $Ab^n + \dots$
 - You can compute A by computing $\lim_{n \rightarrow \infty} r_n/b^n$
 - Try to compute the form of \dots by considering the sequence $r_n - Ab^n$; that is,

$$r_1 - Ab, r_2 - Ab^2, r_3 - Ab^3, \dots$$

- $\lim_{n \rightarrow \infty} r_{n+1}/r_n = 1$, then r_n is most likely a polynomial.
- $\lim_{n \rightarrow \infty} r_{n+1}/r_n = 0$, then r_n may have the form $A/b^{f(n)}$, where $f(n)/n \rightarrow \infty$
 - $f(n)$ could be $n \log n$ or n^2 , for example

Once you have guessed the form of r_n , prove that your guess is right by induction.

More examples

Come up with a simple formula for the sequence

$$1, 5, 13, 41, 121, 365, 1093, 3281, 9841, 29525$$

Compute limit of r_{n+1}/r_n :

$$5/1 = 5, \quad 13/5 \approx 2.6, \quad 41/13 \approx 3.2, \quad 121/41 \approx 2.95, \\ \dots, 29525/9841 \approx 3.000$$

Guess: limit is 3 ($\Rightarrow r_n = A3^n + \cdot$)

Compute limit of $r_n/3^n$:

$$1/3 \approx .33, \quad 5/9 \approx .56, \quad 13/27 \approx .5, \quad 41/81 \approx .5, \\ \dots, 29525/3^{10} \approx .5000$$

Guess: limit is $1/2$ ($\Rightarrow r_n = \frac{1}{2}3^n + \dots$) +

Compute $r_n - 3^n/2$:

$$(1 - 3/2), (5 - 9/2), (13 - 27/2), (41 - 81/2), \dots \\ = -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \dots$$

Guess: general term is $3^n/2 + (-1)^n/2$

Verify (by induction ...)

One more example

Find a formula for

$$\frac{1}{1 \cdot 4} + \frac{1}{4 \cdot 7} + \frac{1}{7 \cdot 10} + \cdots + \frac{1}{(3n-2)(3n+1)}$$

Some values:

- $r_1 = 1/4$
- $r_2 = 1/4 + 1/28 = 8/28 = 2/7$
- $r_3 = 1/4 + 1/28 + 1/70 = (70 + 10 + 4)/280 = 84/280 = 3/10$

Conjecture: $r_n = n/(3n+1)$. Let this be $P(n)$.

Basis: $P(1)$ says that $r_1 = 1/4$.

Inductive step:

$$\begin{aligned} r_{n+1} &= r_n + \frac{1}{(3n+1)(3n+4)} \\ &= \frac{n}{3n+1} + \frac{1}{(3n+1)(3n+4)} \\ &= \frac{n(3n+4)+1}{(3n+1)(3n+4)} \\ &= \frac{3n^2+4n+1}{(3n+1)(3n+4)} \\ &= \frac{(n+1)(3n+1)}{(3n+1)(3n+4)} \\ &= \frac{n+1}{3n+4} \end{aligned}$$

Faulty Inductions

Part of why I want you to write out your assumptions carefully is so that you don't get led into some standard errors.

Theorem: All women are blondes.

Proof by induction: Let $P(n)$ be the statement: For any set of n women, if at least one of them is a blonde, then all of them are.

Basis: Clearly OK.

Inductive step: Assume $P(n)$. Let's prove $P(n + 1)$.

Given a set W of $n + 1$ women, one of which is blonde. Let A and B be two subsets of W of size n , each of which contains the known blonde, whose union is W .

By the induction hypothesis, each of A and B consists of all blondes. Thus, so does W . This proves $P(n) \Rightarrow P(n + 1)$.

Take W to be the set of women in the world, and let $n = |W|$. Since there is clearly at least one blonde in the world, it follows that all women are blonde!

Where's the bug?

Theorem: Every integer > 1 has a unique prime factorization.

[The result is true, but the following proof is not:]

Proof: By strong induction. Let $P(n)$ be the statement that n has a unique factorization. We prove $P(n)$ for $n > 1$.

Basis: $P(2)$ is clearly true.

Induction step: Assume $P(2), \dots, P(n)$. We prove $P(n+1)$. If $n+1$ is prime, we are done. If not, it factors somehow. Suppose $n+1 = rs$ $r, s > 1$. By the induction hypothesis, r has a unique factorization $\prod_i p_i$ and s has a unique prime factorization $\prod_j q_j$. Thus, $\prod_i p_i \prod_j q_j$ is a prime factorization of $n+1$, and since none of the factors of either piece can be changed, it must be unique.

What's the flaw??

Problem: Suppose $n + 1 = 36$. That is, you've proved that every number up to 36 has a unique factorization. Now you need to prove it for 36.

36 isn't prime, but $36 = 3 \times 12$. By the induction hypothesis, 12 has a unique prime factorization, say $p_1 p_2 p_3$. Thus, $36 = 3 p_1 p_2 p_3$.

However, 36 is also 4×9 . By the induction hypothesis, $4 = q_1 q_2$ and $9 = r_1 r_2$. Thus, $36 = q_1 q_2 r_1 r_2$.

How do you know that $3 p_1 p_2 p_3 = q_1 q_2 r_1 r_2$. (They do, but it doesn't follow from the induction hypothesis.)

This is a *breakdown error*. If you're trying to show something is unique, and you break it down (as we broke down $n+1$ into r and s) you have to argue that nothing changes if we break it down a different way. What if $n + 1 = tu$?

- The actual proof of this result is quite subtle

Theorem: The sum of the internal angles of a regular n -gon is $180(n - 2)$ for $n \geq 3$.

Proof: By induction. Let $P(n)$ be “the sum of the internal angles of a regular n -gon is $180(n - 2)$.” For $n = 3$, the result was shown in high school. Assume $P(n)$; let’s prove $P(n + 1)$. Given a regular $(n + 1)$ -gon, we can lop off one of the corners:

By the induction hypothesis, the sum of the internal angles of the regular n -gon is $180(n - 2)$ degrees; the sum of the internal angles of the triangle is 180 degrees. Thus, the internal angles of the original $(n + 1)$ -gon is $180(n - 1)$. What’s wrong??

- When you lop off a corner, you don’t get a *regular* n -gon.

The fix: **Strengthen the induction hypothesis.**

- Let $P(n)$ say that the sum of the internal angles of *any* n -gon is $180(n - 2)$.

Consider 0-1 sequences in which 1's may not appear consecutively, except in the rightmost two positions.

- 010110 is not allowed, but 010011 is

Prove that there are 2^n allowed sequences of length n for $n \geq 1$

Why can't this be right?

“Proof” Let $P(n)$ be the statement of the theorem.

Basis: There are 2 sequences of length 1—0 and 1—and they're both allowed.

Inductive step: Assume $P(n)$. Let's prove $P(n + 1)$. Take any allowed sequence x of length n . We get a sequence of length $n + 1$ by appending either a 0 or 1 at the end. In either case, it's allowed.

- If x ends with a 1, it's OK, because $x1$ is allowed to end with 2 1's.

Thus, $s_{n+1} = 2s_n = 22^n = 2^{n+1}$.

Where's the flaw?

- What if x already ends with 2 1's?

Correct expression involves separating out sequences which end in 0 and 1 (it's done in Chapter 5, but I'm not sure we'll get to it)

The Sorites Paradox

If a pile of sand has 1,000,000 grains of sand, it's a heap.

Removing one grain of sand from a heap leaves 1 heap.

Therefore, by induction, if a pile of sand has only one grain, it's also a heap.

Prove by induction on n that if a pile of sand has 1,000,000— n grains of sand, it's a heap.

Where's the bug?

- This leads to a whole topic in the philosophy of language called “vagueness”

Inductive Definitions

Example: Define $\sum_{k=1}^n a_k$ inductively (i.e., by induction on n):

- $\sum_{k=1}^1 a_k = a_1$
- $\sum_{k=1}^{n+1} a_k = \sum_{k=1}^n a_k + a_{n+1}$

The inductive definition avoids the use of \dots , and thus is less ambiguous.

Example: An inductive definition of $n!$:

- $1! = 1$
- $(n+1)! = (n+1)n!$

Could even start with $0! = 1$.

Inductive Definitions of Sets

A *palindrome* is an expression that reads the same backwards and forwards:

- Madam I'm Adam
- Able was I ere I saw Elba

What is the set of palindromes over $\{a, b, c, d\}$? Two approaches:

1. The smallest set P such that
 - (a) P contains $a, b, c, d, aa, bb, cc, dd$
 - (b) if x is in P , then so is axa, bxb, cxc , and dxd

Things to think about:

- How do you know that there is a smallest set (one which is a subset of all others)
- How do you know that it doesn't contain ab

2. Define P_n , the palindromes of length n , inductively:

- $P_1 = \{a, b, c, d\}$
- $P_2 = \{aa, bb, cc, dd\}$
- $P_{n+1} = \{axa, bxb, cxc, dxd \mid x \in P_{n-1}\}$ for $n \geq 2$

Let $P' = \cup_n P_n$.

Theorem: $P = P'$. (The two approaches define the same set.)

Proof: Show $P \subseteq P'$ and $P' \subseteq P$.

To see that $P \subseteq P'$, it suffices to show that

(a) P' contains $a, b, c, d, aa, bb, cc, dd$

(b) if x is in P' , then so is axa, bxb, cxc , and dxd

(since P is the smallest set with these properties).

Clearly $P_1 \cup P_2$ satisfies (1), so P' does. And if $x \in P'$, then $x \in P_n$ for some n , in which case axa, bxb, cxc , and dxd are all in P_{n+2} and hence in P' . Thus, $P \subseteq P'$.

To see that $P' \subseteq P$, we prove by strong induction that $P_n \subseteq P$ for all n . Let $P(n)$ be the statement “ $P_n \subseteq P$.”

Basis: $P_1, P_2 \subseteq P$: Obvious.

Suppose $P_1, \dots, P_n \subseteq P$. If $n \geq 2$, the fact that $P_{n+1} \subseteq P$ follows immediately from (b). (Actually, all we need is the fact that $P_{n-1} \subseteq P$, which follows from the (strong) induction hypothesis.)

Thus, $P' = \cup_n P_n \subseteq P$.

Recall that the set of palindromes is the smallest set P such that

- (a) P contains $a, b, c, d, aa, bb, cc, dd$
- (b) if x is in P , then so is axa, bxb, cxc , and dxd

“Smallest” is not in terms of cardinality.

- P is guaranteed to be infinite

“Smallest” is in terms of the subset relation.

Here’s a set that satisfies (a) and (b) and isn’t the smallest:

Define Q_n inductively:

- $Q_1 = \{a, b, c, d\}$
- $Q_2 = \{aa, bb, cc, dd, ab\}$
- $Q_{n+1} = \{axa, bxb, cxc, dxd \mid x \in Q_{n-1}\}, n \geq 2$

Let $Q = \cup_n Q_n$.

It’s easy to see that Q satisfies (a) and (b), but it isn’t the smallest set to do so.

Fibonacci Numbers

[Leonardi of Pisa, 12th century:] Suppose you start with two rabbits, one of each gender. After two months, they produce two rabbits (one of each gender) as offspring. Each subsequent pair of offspring behaves the same way, producing another pair in two months. Rabbits never die. How many rabbits do you have after n months?

Let f_n be the number of pairs after n months.

By assumption, $f_1 = f_2 = 1$

For $n > 2$, $f_{n+1} = f_n + f_{n-1}$

- In month $n + 1$, each pair of rabbits that have been around for at least two months (f_{n-1}) produces another pair. So you have f_{n-1} new pairs on top of the f_n you had after n months.
- This is an *inductive definition* of a sequence

The Fibonacci sequence has the form 1, 1, 2, 3, 5, 8, ...

Fibonacci numbers grow exponentially

The Fibonacci sequence has lots of nice properties; we'll prove one.

Let $r = (1 + \sqrt{5})/2 \approx 1.62$.

Claim: $f_n \geq r^{n-2}$ for all n .

Where did this weird r come from?

- It's a solution to the equation $r^2 = r + 1$.
 - The other solution is $(1 - \sqrt{5})/2$

We can prove the claim by induction.

Base case: $f_1 = 1$; $r^{-1} = 1/r < 1$; so $f_1 > r^{-1}$

$f_2 = 1$; $r^0 = 1$; so $f_2 \geq r^0$.

Inductive step: If $n \geq 2$

$$\begin{aligned} f_{n+1} &= f_n + f_{n-1} \\ &\geq r^{n-2} + r^{n-3} \\ &= r^{n-3}(r + 1) \\ &= r^{n-3}r^2 && [\text{since } r + 1 = r^2] \\ &= r^{n-1} \end{aligned}$$

That's it!

It can be shown [see Chapter 5] that

$$f_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right]$$

The Trust Game

Consider a game where, after n steps, there are piles of money on the table:

- The big one has $\$2^{n+1}$; the small one has $\$2^{n-1}$

There are two players, Alice and Bob. Initially Alice is in charge. She can either quit the game or continue

- If she quits, she gets the money in the bigger pile (\$4) and Bob gets the money in the smaller pile (\$1)
- If she continues, Bob is in charge
- If he quits, he gets the money in the bigger pile (\$8), Alice gets the money in the smaller pile (\$2).
- If he continues, Alice is in charge, and so on.
- The game goes on for 20 steps;
 - if they're still playing then, Bob gets $\$2^{21}$ ($> \$2,000,000$); Alice gets $\$2^{19}$ ($\approx \$500,000$)

What should you do?

- Should you trust the other player to keep playing, or take your money and run?

In the game theory literature, this is called the *centipede game*.

What should Alice do if they're still playing at step 19?

- If she quits, she gets $\$2^{20}$ (about \$1,000,000); if she continues she gets only $\$2^{19}$.
- So Alice will quit, which means Bob will get $\$2^{18}$

So what should Bob do if they're still playing at step 18?

- If he quits, he gets $\$2^{19}$; if he continues, most likely he'll get $\$2^{18}$, since Alice will quit at step 19.
- So Bob quits, which means Alice will get $\$2^{16}$.

Continuing this way (by *backwards induction*), Alice quits at step 1 and gets \$4!

Under a specific model of *rationality*, quitting at the first step is the only right thing to do.

- It's the only *Nash equilibrium*

In practice (with smaller amounts of money), people play for a little while before quitting.

The muddy children puzzle

We can prove by induction on k that if k children have muddy foreheads, they say “yes” on the k^{th} question. It appears as if the father didn’t tell the children anything they didn’t already know. Yet without the father’s statement, they could not have deduced anything. So what was the role of the father’s statement?