Mathematical Induction

CS 2800: Discrete Structures, Fall 2014

Sid Chaudhuri

 A prime number is a positive integer with exactly two divisors: 1 and itself


- 2, 3, 5, 7, 11, 13, 17, ...


 A prime number is a positive integer with exactly two divisors: 1 and itself


• Claim: Every natural number ≥ 2 can be expressed as a finite product of prime numbers


- E.g.
$$3 = 3$$

 $15 = 3 \times 5$
 $16 = 2 \times 2 \times 2 \times 2$













(And there was


much rejoicing)
































• Claim: Every natural number ≥ 2 can be expressed as a finite product of prime numbers

• Claim: Every natural number ≥ 2 can be expressed as a finite product of prime numbers

Proof:

 Let's start with 2. This is trivially the product of one prime number. So the claim is true for 2.

• Claim: Every natural number ≥ 2 can be expressed as a finite product of prime numbers

Proof:

- Let's start with 2. This is trivially the product of one prime number. So the claim is true for 2.
- Now assume the claim is true for all natural numbers
 from 2 to n

• Claim: Every natural number ≥ 2 can be expressed as a finite product of prime numbers

Proof:

- Let's start with 2. This is trivially the product of one prime number. So the claim is true for 2.
- Now assume the claim is true for all natural numbers
 from 2 to n

which n? we'll assume n is arbitrary, but ≥ 2

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)
 - Consider n+1

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)
 - Consider n+1
 - If it is prime, then the claim is trivially true for n + 1

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)
 - Consider n+1
 - If it is prime, then the claim is trivially true for n + 1
 - **If it is composite**, then it is (by definition), the product of two natural numbers a and b, both > 1 but < n + 1

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)
 - Consider n+1
 - If it is prime, then the claim is trivially true for n + 1
 - **If it is composite**, then it is (by definition), the product of two natural numbers a and b, both > 1 but < n + 1
 - Since the claim is assumed to be true for all natural numbers from 2 to n, it is also true for a and b

- Proof (contd):
 - (We've assumed the claim is true for all natural numbers from 2 to n)
 - Consider n+1
 - If it is prime, then the claim is trivially true for n + 1
 - **If it is composite**, then it is (by definition), the product of two natural numbers a and b, both > 1 but < n + 1
 - Since the claim is assumed to be true for all natural numbers from 2 to n, it is also true for a and b
 - So $a = p_1 p_2 p_3 \dots p_m$ and $b = q_1 q_2 q_3 \dots q_k$, where all p_i , q_i are prime

Proof (contd):

- (We've assumed the claim is true for all natural numbers from 2 to n)
- Consider n+1
 - If it is prime, then the claim is trivially true for n + 1
 - **If it is composite**, then it is (by definition), the product of two natural numbers a and b, both > 1 but < n + 1
 - Since the claim is assumed to be true for all natural numbers from 2 to n, it is also true for a and b
 - So $a = p_1 p_2 p_3 \dots p_m$ and $b = q_1 q_2 q_3 \dots q_k$, where all p_i , q_i are prime
 - So $n + 1 = p_1 p_2 ... p_m q_1 q_2 ... q_k$, i.e. the claim is true for n + 1

• What have we shown?

- What have we shown?
 - 2 is a finite product of prime numbers

What have we shown?

- 2 is a finite product of prime numbers
- For any $n \ge 2$, if all natural numbers from 2 to n are finite products of prime numbers, then so is n+1

- What have we shown?
 - 2 is a finite product of prime numbers
 - For any $n \ge 2$, if all natural numbers from 2 to n are finite products of prime numbers, then so is n+1
- Can we conclude that all natural numbers ≥ 2 are finite products of prime numbers?

- What have we shown?
 - 2 is a finite product of prime numbers
 - For any $n \ge 2$, if all natural numbers from 2 to n are finite products of prime numbers, then so is n+1
- Can we conclude that all natural numbers ≥ 2 are finite products of prime numbers?
 - Yes!

- What have we shown?
 - 2 is a finite product of prime numbers
 - For any $n \ge 2$, if all natural numbers from 2 to n are finite products of prime numbers, then so is n+1
- Can we conclude that all natural numbers ≥ 2 are finite products of prime numbers?
 - Yes!
 - If it's true for 2, it must be true for 3. If it's true for 3, it must be true for 4. If it's true for 4...

- What have we shown?
 - 2 is a finite product of prime numbers
 - For any $n \ge 2$, if all natural numbers from 2 to n are finite products of prime numbers, then so is n+1
- Can we conclude that all natural numbers ≥ 2 are finite products of prime numbers?
 - Yes!
 - If it's true for 2, it must be true for 3. If it's true for 3, it must be true for 4. If it's true for 4...
 - We've applied the Principle of Mathematical Induction

• We need to prove statement S(n) about natural numbers n

- We need to prove statement S(n) about natural numbers n
- If we can show that
 - Base case: S(1) is true, and

- We need to prove statement S(n) about natural numbers n
- If we can show that
 - Base case: S(1) is true, and
 - *Inductive step*: Assumption that S(k) is true for all natural numbers $k \le n$ implies that S(n+1) is true

• We need to prove statement S(n) about natural numbers n

Inductive hypothesis

- If we can show that
 - Base case: S(1) is true, and
 - *Inductive step*: Assumption that S(k) is true for all natural numbers $k \le n$ implies that S(n+1) is true

Inductive hypothesis

- We need to prove statement S(n) about natural numbers n
- If we can show that
 - Base case: S(1) is true, and
 - *Inductive step*: Assumption that S(k) is true for all natural numbers $k \le n$ implies that S(n+1) is true
- ... then S(n) is true for all natural numbers n

- Can start from an integer k which is not 1. The statement is then proved for all integers $\geq k$
 - Starting from a negative number doesn't change the applicability of induction, but be careful when stating the inductive hypothesis and when proving the inductive step!

- Can start from an integer k which is not 1. The statement is then proved for all integers $\geq k$
 - Starting from a negative number doesn't change the applicability of induction, but be careful when stating the inductive hypothesis and when proving the inductive step!
- Can start with multiple base cases, i.e. directly prove S(1), S(2), ..., S(m) and use induction for all integers greater than m

- Can start from an integer k which is not 1. The statement is then proved for all integers $\geq k$
 - Starting from a negative number doesn't change the applicability of induction, but be careful when stating the inductive hypothesis and when proving the inductive step!
- Can start with multiple base cases, i.e. directly prove S(1), S(2), ..., S(m) and use induction for all integers greater than m
- Can apply to any countable set (prove!)

•
$$S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"$$

- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true


- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true
- Inductive step:
 - Assume 1 + 2 + ... + k = k(k+1)/2 for all natural numbers k < n

- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true
- Inductive step:
 - Assume $1+2+\ldots+k=k\left(k+1\right)/2$ for all natural numbers $k\leq n$
 - Then 1 + 2 + ... + n + (n + 1) = n(n + 1) / 2 + (n + 1)


- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true
- Inductive step:
 - Assume 1 + 2 + ... + k = k(k+1)/2 for all natural numbers k < n
 - Then 1 + 2 + ... + n + (n + 1) = n(n + 1) / 2 + (n + 1)= (n + 1)(n + 2) / 2

- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true
- Inductive step:
 - Assume $1+2+\ldots+k=k\left(k+1\right)/2$ for all natural numbers $k\leq n$
 - Then 1 + 2 + ... + n + (n + 1) = n(n + 1) / 2 + (n + 1)= (n + 1)(n + 2) / 2
 - Hence S(n + 1) is true

- S(n) = "1 + 2 + 3 + ... + n = n(n + 1) / 2"
- Base case: 1 = 1(1 + 1)/2, hence S(1) is true
- Inductive step:
 - Assume $1+2+\ldots+k=k\left(k+1\right)/2$ for all natural numbers $k\leq n$
 - Then 1 + 2 + ... + n + (n + 1) = n(n + 1) / 2 + (n + 1)= (n + 1)(n + 2) / 2
 - Hence S(n + 1) is true
- Hence by induction, S(n) is true for all $n \in \mathbb{N}$





 $S(n) = \text{``A } 2^n \times 2^n \text{ chessboard with one corner missing can be tiled with triominoes''}$



Base case: A 2×2 chessboard with one corner missing is just a single triomino, so S(1) is true

- Assume a $2^k \times 2^k$ chessboard with a corner missing can be tiled with triominoes, for all natural numbers $k \le n$
- Consider a $2^{n+1} \times 2^{n+1}$ board (with a corner missing)





- Assume a $2^k \times 2^k$ chessboard with a corner missing can be tiled with triominoes, for all natural numbers $k \le n$
- Consider a $2^{n+1} \times 2^{n+1}$ board (with a corner missing)
- It is just four $2^n \times 2^n$ boards, plus one triomino

- Assume a $2^k \times 2^k$ chessboard with a corner missing can be tiled with triominoes, for all natural numbers $k \le n$
- Consider a $2^{n+1} \times 2^{n+1}$ board (with a corner missing)
- It is just four $2^n \times 2^n$ boards, plus one triomino
- From the inductive hypothesis, we know each of these boards can be tiled with triominoes

- Assume a $2^k \times 2^k$ chessboard with a corner missing can be tiled with triominoes, for all natural numbers $k \le n$
- Consider a $2^{n+1} \times 2^{n+1}$ board (with a corner missing)
- It is just four $2^n \times 2^n$ boards, plus one triomino
- From the inductive hypothesis, we know each of these boards can be tiled with triominoes
- Hence S(n + 1) is true

- Assume a $2^k \times 2^k$ chessboard with a corner missing can be tiled with triominoes, for all natural numbers $k \le n$
- Consider a $2^{n+1} \times 2^{n+1}$ board (with a corner missing)
- It is just four $2^n \times 2^n$ boards, plus one triomino
- From the inductive hypothesis, we know each of these boards can be tiled with triominoes
- Hence S(n + 1) is true
- Hence by induction, S(n) is true for all $n \in \mathbb{N}$

S(n) = "The following program returns the n^{th} Fibonacci number, given input n"

```
int fibonacci(int n)
{
  if (n <= 2)
    return 1;
  else
    return fibonacci(n - 1) + fibonacci(n - 2);
}</pre>
```

```
S(n) = "The following program returns the n^{th}
Fibonacci number, given input n"
            1, 1, 2, 3, 5, 8, 13, 21, 34, ...
int fibonacci(int n)
  if (n \le 2)
    return 1;
  else
    return fibonacci(n -1) + fibonacci(n -2);
```

• Base cases: fibonacci(1) and fibonacci(2) return the first two Fibonacci numbers (1 and 1, easily verified), so S(1) and S(2) are true

```
int fibonacci(int n)
{
  if (n <= 2)
    return 1;
  else
    return fibonacci(n - 1) + fibonacci(n - 2);
}</pre>
```

- Inductive step:
 - Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$

- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$
- Consider fibonacci(n + 1)


- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all k < n
- Consider fibonacci(n + 1)
 - From the program, it returns
 fibonacci(n) + fibonacci(n 1)

- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$
- Consider fibonacci(n + 1)
 - From the program, it returns
 fibonacci(n) + fibonacci(n 1)
 - From the inductive hypothesis, these are the $n^{\rm th}$ and $(n-1)^{\rm th}$ Fibonacci numbers, respectively

- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$
- Consider fibonacci(n + 1)
 - From the program, it returns
 fibonacci(n) + fibonacci(n 1)
 - From the inductive hypothesis, these are the $n^{\rm th}$ and $(n-1)^{\rm th}$ Fibonacci numbers, respectively
 - Hence by definition of Fibonacci numbers, fibonacci (n + 1) returns the $(n+1)^{\text{th}}$ Fibonacci number, i.e. S(n+1) is true

- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$
- Consider fibonacci(n + 1)
 - From the program, it returns
 fibonacci(n) + fibonacci(n 1)
 - From the inductive hypothesis, these are the $n^{\rm th}$ and $(n-1)^{\rm th}$ Fibonacci numbers, respectively
 - Hence by definition of Fibonacci numbers, fibonacci (n + 1) returns the $(n + 1)^{th}$ Fibonacci number, i.e. S(n + 1) is true
- Hence by induction, S(n) is true for all $n \in \mathbb{N}$

- Assume, for $n \ge 2$, that fibonacci(k) returns the k^{th} Fibonacci number for all $k \le n$
- Consider fibonacci(n + 1)
 - From the program, it returns
 fibonacci(n) + fibonacci(n 1)
 - From the inductive hypothesis, these are the $n^{\rm th}$ and $(n-1)^{\rm th}$ Fibonacci numbers, respectively
 - Hence by definition of Fibonacci numbers, fibonacci (n + 1) returns the $(n + 1)^{th}$ Fibonacci number, i.e. S(n + 1) is true
- Hence by induction, S(n) is true for all $n \in \mathbb{N}$



• S(n) = "Any group of n horses is the same color"

• S(n) = "Any group of n horses is the same color"

Doesn't say anything about whether different groups of n horses have different colors or not

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n+1 horses can be expressed as the union of two groups of n horses each

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n+1 horses can be expressed as the union of two groups of n horses each
 - These two groups are individually the same color, by hypothesis

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n+1 horses can be expressed as the union of two groups of n horses each
 - These two groups are individually the same color, by hypothesis
 - ... and they overlap

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n+1 horses can be expressed as the union of two groups of n horses each
 - These two groups are individually the same color, by hypothesis
 - ... and they overlap
 - So the group of n + 1 horses is the same color

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n + 1 horses can be expressed as the union of two groups of n horses each
 - These two groups are individually the same color, by hypothesis
 - ... and they overlap
 - So the group of n + 1 horses is the same color
- Hence by induction, all horses are the same color

- S(n) = "Any group of n horses is the same color"
- Base case: A single horse is obviously the same color as itself, so S(1) is true
- Inductive step:
 - Assume any group of $k \le n$ horses is the same color
 - A group of n+1 horses can be expressed as the union of two groups of n horses each
 - These two groups are individually the same color, by hypothesis
 - ... and they overlap \longrightarrow Not for n = 1!
 - So the group of n + 1 horses is the same color
- Hence by induction, all horses are the same color