

# CS5670: Computer Vision Noah Snavely

#### Light & Perception






# Announcements

- Project 4
  - To be released on Wednesday
  - Groups of 2
- Please submit midterm regrade requests by tomorrow (Tuesday, 4/9) by 11:59pm
- Project 3 artifact voting open until Friday
- Quiz 3 this Wednesday, April 10
  - Will cover topics since the midterm was released (single-view modeling, stereo, two-view geometry / Fundamental matrices, and today's lecture on light)

# Road map

- Finishing up geometry
- Next: deep learning and hot topics in computer vision

#### Can we determine shape from lighting?



- Are these spheres?
  - Or just flat discs painted with varying albedo?
  - There is ambiguity between *shading* and *reflectance*
  - But still, as humans we can understand the shapes of these objects

# What we know: Stereo





Key Idea: use feature motion to understand shape

# Next: Photometric Stereo





Key Idea: use pixel brightness to understand shape

# Next: Photometric Stereo








Key Idea: use pixel brightness to understand shape

# **Photometric Stereo**

What results can you get?



Input (1 of 12)

Normals (RGB colormap)


Normals (vectors)

Shaded 3D rendering

Textured 3D rendering



# Light




by Ted Adelson

#### Readings

• Szeliski, 2.2, 2.3.2

# Light



by Ted Adelson

#### Readings


• Szeliski, 2.2, 2.3.2

# Properties of light


Today

- What is light?
- How do we measure it?
- How does light propagate?
- How does light interact with matter?

#### What determines the brightness of a pixel?



#### What determines the brightness of a pixel?




@robertwestonbreshears https://www.instagram.com/p/BtgX55ZBhU-/

#### What determines the brightness of a pixel?



What determines the brightness of an image pixel?



Slide by L. Fei-Fei

# What is light?


Electromagnetic radiation (EMR) moving along rays in space

- $R(\lambda)$  is EMR, measured in units of power (watts)
  - $-\lambda$  is wavelength



Light field

 We can describe all of the light in the scene by specifying the radiation (or "radiance" along all light rays) arriving at every point in space and from every direction



 $R(X,Y,Z,\theta,\phi,\lambda,t)$ 

# **Color perception**

Electromagnetic radiation (EMR) moving along rays in space

- $R(\lambda)$  is EMR, measured in units of power (watts)
  - $-\lambda$  is wavelength




Perceiving light

- How do we convert radiation into "color"?
- What part of the spectrum do we see?

# Visible light


We "see" electromagnetic radiation in a range of wavelengths



# Light spectrum

The appearance of light depends on its power **spectrum** 


How much power (or energy) at each wavelength



Our visual system converts a light spectrum into "color"

• This is a rather complex transformation

## The human visual system




Color perception

- Light hits the retina, which contains photosensitive cells

   rods and cones
- These cells convert the spectrum into a few discrete values


### Density of rods and cones



Rods and cones are non-uniformly distributed on the retina

- Rods responsible for intensity, cones responsible for color
- Fovea Small region (1 or 2°) at the center of the visual field containing the highest density of cones (and no rods).
- Less visual acuity in the periphery-many rods wired to the same neuron

#### Demonstrations of visual acuity



With one eye shut, at the right distance, all of these letters should appear equally legible (Glassner, 1.7).

#### Demonstrations of visual acuity

With left eye shut, look at the cross on the left. At the right distance, the circle on the right should disappear (Glassner, 1.8).

# Brightness contrast and constancy

The apparent brightness depends on the surrounding region

• **brightness contrast**: a constant colored region seems lighter or darker depending on the surrounding intensity:



- <u>http://www.sandlotscience.com/Contrast/Checker Board 2.htm</u>
- **brightness constancy**: a surface looks the same under widely varying lighting conditions.

# Light response is nonlinear

Our visual system has a large *dynamic range* 

- We can resolve both light and dark things at the same time
- One mechanism for achieving this is that we sense light intensity on a *logarithmic scale* 
  - an exponential intensity ramp will be seen as a linear ramp
- Another mechanism is *adaptation* 
  - rods and cones adapt to be more sensitive in low light, less sensitive in bright light.

# Visual dynamic range

Background	Luminance (candelas per square meter)
Horizon sky	
Moonless overcast night	0.00003
Moonless clear night	0.0003
Moonlit overcast night	0.003
Moonlit clear night	0.03
Deep twilight	0.3
Twilight	3
Very dark day	30
Overcast day	300
Clear day	3,000
Day with sunlit clouds	30,000
Daylight fog	
Dull	300-1,000
Typical	1,000-3,000
Bright	3,000-16,000
Ground	
Overcast day	30-100
Sunny day	300
Snow in full sunlight	16,000

A piece of white paper can be 1,000,000,000 times brighter in outdoor sunlight than in a moonless night.

BUT in a given lighting condition, light ranges over only about two orders of magnitude.

#### FIGURE 1.13

Luminance of everyday backgrounds. Source: Data from Rea, ed., Lighting Handbook 1984 Reference and Application, fig. 3-44, p. 3-24.

#### Learning to See in the Dark

Chen Chen, Qifeng Chen, Jia Xu and Vladlen Koltun

IEEE Conference on Computer Vision and Pattern Recognition (CVPR), 2018




- (a) Camera output with ISO 8,000
- (b) Camera output with ISO 409,600
- (c) Our result from the raw data of (a)

Figure. Extreme low-light imaging by a Sony a7S II camera using ISO 8000, f/5.6, 1/30 second. Dark indoor environment. The illuminance at the camera is <0.1 lux.


http://cchen156.web.engr.illinois.edu/SID.html

## Visual dynamic range




If we were sensitive to this whole range all the time, we wouldn't be able to discriminate lightness levels in a typical scene.

The visual system solves this problem by restricting the 'dynamic range' of its response to match the current overall or 'ambient' light level.




# **Color perception**



#### Three types of cones

- Each is sensitive in a different region of the spectrum
  - but regions overlap
  - Short (S) corresponds to blue
  - Medium (M) corresponds to green
  - Long (L) corresponds to red
- Different sensitivities: we are more sensitive to green than red
  - varies from person to person (and with age)
- Colorblindness—deficiency in at least one type of cone

# **Color perception**



Rods and cones act as filters on the spectrum

- To get the output of a filter, multiply its response curve by the spectrum, integrate over all wavelengths
  - Each cone yields one number
- Q: How can we represent an entire spectrum with 3 numbers?
- A: We can't! Most of the information is lost.
  - As a result, two different spectra may appear indistinguishable
    - » such spectra are known as **metamers**
    - » <u>http://www.cs.brown.edu/exploratories/freeSoftware/repository/edu/brown/cs/explo ratories/applets/spectrum/metamers\_guide.html</u>

#### What kind of bulb is it?



#### http://www.chemistryland.com/CHM107Lab/Exp7/Spectroscope/Spectroscope.html

## Perception summary


The mapping from radiance to perceived color is quite complex!

- We throw away most of the data
- We apply a logarithm
- Brightness affected by pupil size
- Brightness contrast and constancy effects

The same is true for cameras

- But we have tools to correct for these effects
  - (Computational Photography)

## Light transport



# Light sources

#### Basic types


- point source
- directional source
  - a point source that is infinitely far away
- area source
  - a union of point sources

More generally

• a light field can describe \*any\* distribution of light sources

What happens when light hits an object?

# **Modeling Image Formation**




We need to reason about:

- How light interacts with the scene
- How a pixel value is related to light energy in the world


Track a "ray" of light all the way from light source to the sensor


## **Directional Lighting**



- Key property: all rays are parallel
- Equivalent to an infinitely distant point source


#### Lambertian Reflectance






## $I = N \cdot L$

Image \_\_\_\_\_ Surface Light intensity \_\_\_\_\_ Image \_\_\_\_\_ cos(angle between N and L)




© Kavita Bala, Computer Science, Cornell University



© Kavita Bala, Computer Science, Cornell University


## Ideal Diffuse Reflection

- Characteristic of multiple scattering materials
- An idealization but reasonable for matte surfaces



© Kavita Bala, Computer Science, Cornell University


### Lambertian Reflectance



- 1. Reflected energy is proportional to cosine of angle between L and N (incoming)
- 2. Measured intensity is viewpoint-independent (outgoing)


### Lambertian Reflectance: Incoming

1. Reflected energy is proportional to cosine of angle between L and N




### Lambertian Reflectance: Incoming

1. Reflected energy is proportional to cosine of angle between L and N




### Lambertian Reflectance: Incoming

1. Reflected energy is proportional to cosine of angle between L and N




Light hitting surface is proportional to the cosine


1. Radiance (what we see) is viewpoint-independent




1. Radiance (what the eye sees) is viewpoint-independent




1. Measured intensity is viewpoint-independent



1. Measured intensity is viewpoint-independent




# Image Formation Model: Final



- 1. Diffuse albedo: what fraction of incoming light is reflected?
  - Introduce scale factor  $k_d$
- 2. Light intensity: how much light is arriving?
  - Compensate with camera exposure (global scale factor)
- 3. Camera response function
  - Assume pixel value is linearly proportional to incoming energy (perform radiometric calibration if not)

## A Single Image: Shape from Shading



 $I = k_d \mathbf{N} \cdot \mathbf{L}$ 

Assume  $k_d$  is 1 for now. What can we measure from one image?

- $\cos^{-1}(I)$  is the angle between N and L
- Add assumptions:
  - Constant albedo
  - A few known normals (e.g. silhouettes)
  - Smoothness of normals

In practice, SFS doesn't work very well: assumptions are too restrictive, too much ambiguity in nontrivial scenes.

#### Application: Detecting composite photos

Real photo

#### Fake photo





#### Questions?