


# CS5670: Computer Vision

## Transformations and warping



# Reading

- Szeliski: Chapter 3.6

# Announcements

- Project 2 out, due Monday, March 4, by 11:59pm on CMSX
  - Please form teams of 2, and create your team on CMSX
  - Please declare your group on CMSX by this Wednesday, 2/19
  - After Wednesday, we will randomly assign ungrouped students to groups
  - Feel free to use Piazza to form groups
  - Project demo last 10 minutes of class

# Alternate Harris score

- For project 2, you will use an alternate definition of the Harris score:


$$R = \lambda_1 \lambda_2 - k \cdot (\lambda_1 + \lambda_2)^2 = \det(M) - k \cdot \text{tr}(M)^2$$

# Image alignment




Why don't these image line up exactly?

# What is the geometric relationship between these two images?




**Answer: Similarity transformation (translation, rotation, uniform scale)**

**What is the geometric relationship between these two images?**





# What is the geometric relationship between these two images?



**Very important for creating mosaics!**

First, we need to know what this transformation is.


Second, we need to figure out how to compute it using feature matches.



# Image Warping


- image filtering: change *range* of image

- $g(x) = h(f(x))$



- image warping: change *domain* of image


- $g(x) = f(h(x))$



# Image Warping


- image filtering: change *range* of image

- $g(x) = h(f(x))$



- image warping: change *domain* of image

- $g(x) = f(h(x))$



# Parametric (global) warping

- Examples of parametric warps:



translation



rotation




aspect

# Parametric (global) warping



$\mathbf{p} = (x, y)$



$\mathbf{p}' = (x', y')$

- Transformation  $T$  is a coordinate-changing machine:

$$\mathbf{p}' = T(\mathbf{p})$$

- What does it mean that  $T$  is global?
  - Is the same for any point  $\mathbf{p}$
  - can be described by just a few numbers (parameters)
- Let's consider *linear* xforms (can be represented by a 2x2 matrix):

$$\mathbf{p}' = \mathbf{T}\mathbf{p} \quad \begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{T} \begin{bmatrix} x \\ y \end{bmatrix}$$

# Common linear transformations

- Uniform scaling by  $s$ :




$$\mathbf{S} = \begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix}$$

What is the inverse?

# Common linear transformations

- Rotation by angle  $\theta$  (about the origin)



$$\mathbf{R} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

What is the inverse?

For rotations:

$$\mathbf{R}^{-1} = \mathbf{R}^T$$



# 2x2 Matrices

- What types of transformations can be represented with a 2x2 matrix?

2D mirror about Y axis?

$$\begin{aligned}x' &= -x \\ y' &= y\end{aligned} \quad \mathbf{T} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

2D mirror across line  $y = x$ ?

$$\begin{aligned}x' &= y \\ y' &= x\end{aligned} \quad \mathbf{T} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

# 2x2 Matrices

- What types of transformations can be represented with a 2x2 matrix?

2D Translation?

$$x' = x + t_x \quad \text{NO!}$$

$$y' = y + t_y$$

Translation is not a linear operation on 2D coordinates

# All 2D Linear Transformations

- Linear transformations are combinations of ...

- Scale,
- Rotation,
- Shear, and
- Mirror

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Properties of linear transformations:

- Origin maps to origin
- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} \begin{bmatrix} i & j \\ k & l \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

# Homogeneous coordinates

Trick: add one more coordinate:

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image  
coordinates



Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

# Translation


- Solution: homogeneous coordinates to the rescue

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$

# Affine transformations

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$



any transformation represented by a 3x3 matrix with last row  $[0 \ 0 \ 1]$  we call an *affine* transformation

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$



# Basic affine transformations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Translate

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Scale

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D *in-plane* rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & sh_x & 0 \\ sh_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Shear

# Affine transformations


- Affine transformations are combinations of ...

- Linear transformations, and
- Translations

$$\begin{bmatrix} x' \\ y' \\ w \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$


- Properties of affine transformations:
  - Origin does not necessarily map to origin
  - Lines map to lines
  - Parallel lines remain parallel
  - Ratios are preserved
  - Closed under composition

# Is this an affine transformation?



# Where do we go from here?

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$




what happens when we  
mess with this row?

affine transformation

# Projective Transformations *aka* Homographies *aka* Planar Perspective Maps

$$\mathbf{H} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix}$$

Called a *homography*  
(or *planar perspective map*)



# Homographies



$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

What happens when  
the denominator is 0?


$$\sim \begin{bmatrix} \frac{ax+by+c}{gx+hy+1} \\ \frac{dx+ey+f}{gx+hy+1} \\ 1 \end{bmatrix}$$




# Points at infinity



# Image warping with homographies



# Homographies



# Homographies

- Homographies ...

- Affine transformations, and
- Projective warps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

- Properties of projective transformations:


- Origin does not necessarily map to origin
- Lines map to lines
- Parallel lines do not necessarily remain parallel
- Ratios are not preserved
- Closed under composition






# Alternate formulation for homographies

$$\begin{bmatrix} x'_i \\ y'_i \\ 1 \end{bmatrix} \cong \begin{bmatrix} h_{00} & h_{01} & h_{02} \\ h_{10} & h_{11} & h_{12} \\ h_{20} & h_{21} & h_{22} \end{bmatrix} \begin{bmatrix} x_i \\ y_i \\ 1 \end{bmatrix}$$

where the length of the vector  $[h_{00} \ h_{01} \ \dots \ h_{22}]$  is 1

# 2D image transformations




Name	Matrix	# D.O.F.	Preserves:	Icon
translation	$\begin{bmatrix} \mathbf{I} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	2	orientation + ...	
rigid (Euclidean)	$\begin{bmatrix} \mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	3	lengths + ...	
similarity	$\begin{bmatrix} s\mathbf{R} & \mathbf{t} \end{bmatrix}_{2 \times 3}$	4	angles + ...	
affine	$\begin{bmatrix} \mathbf{A} \end{bmatrix}_{2 \times 3}$	6	parallelism + ...	
projective	$\begin{bmatrix} \tilde{\mathbf{H}} \end{bmatrix}_{3 \times 3}$	8	straight lines	

These transformations are a nested set of groups

- Closed under composition and inverse is a member

# Implementing image warping


- Given a coordinate xform  $(x',y') = T(x,y)$  and a source image  $f(x,y)$ , how do we compute a transformed image  $g(x',y') = f(T(x,y))$ ?





# Forward Warping


- Send each pixel  $f(x)$  to its corresponding location  $(x',y') = T(x,y)$  in  $g(x',y')$ 
  - What if pixel lands “between” two pixels?






# Forward Warping

- Send each pixel  $f(x,y)$  to its corresponding location  $x' = h(x,y)$  in  $g(x',y')$ 
  - What if pixel lands “between” two pixels?
  - Answer: add “contribution” to several pixels, normalize later (*splattling*)
  - Can still result in holes




# Inverse Warping

- Get each pixel  $g(x',y')$  from its corresponding location  $(x,y) = T^{-1}(x',y')$  in  $f(x,y)$ 
  - Requires taking the inverse of the transform
  - What if pixel comes from “between” two pixels?



# Inverse Warping

- Get each pixel  $g(\mathbf{x}')$  from its corresponding location  $\mathbf{x}' = \mathbf{h}(\mathbf{x})$  in  $f(\mathbf{x})$ 
  - What if pixel comes from “between” two pixels?
  - Answer: *resample* color value from *interpolated (prefiltered)* source image



# Interpolation

- Possible interpolation filters:
  - nearest neighbor
  - bilinear
  - bicubic
  - sinc
- Needed to prevent “jaggies” and “texture crawl”  
(with prefiltering)



**Questions?**