


Lecture 23: Light, shading, and color

CS 5670: Introduction to Computer Vision





What can we infer from intensity changes?







(a) an image









(b) a likely explanation



(a) an image





(b) a likely explanation




(c) painter's explanation


E. Adelson and A. Pentland, "The perception of shading and reflectance," *Perception as Bayesian inference*, 1996.








(b) a likely explanation





(c) painter's explanation



(d) sculptor's explanation

E. Adelson and A. Pentland, "The perception of shading and reflectance," *Perception as Bayesian inference*, 1996.






E. Adelson and A. Pentland, "The perception of shading and reflectance," *Perception as Bayesian inference*, 1996.

Today


- Shape from shading
- Intrinsic image decomposition
- Color perception

Shape perception





Interaction of light and surfaces



Bidirectional reflectance distribution function (BRDF)

 $BRDF = f(\theta_i, \phi_i, \theta_e, \phi_e, \lambda) = \frac{L(\theta_e, \phi_e, \lambda)}{E(\theta_i, \phi_i, \lambda)}$

Spectral irradiance: incident power per unit area, per unit wavelength.

[Horn, 1986]


Source: W. Freeman


Effect of BRDF on sphere rendering




Diffuse reflection

For now, ignore specular reflection





And refraction...






And interreflections...







Source: Photometric Methods for 3D Modeling, Matsushita, Wilburn, Ben-Ezra. Changes by N. Snavely

Diffuse reflection








Diffuse reflection

- Dull, matte surfaces like chalk or latex paint
- Microfacets scatter incoming light randomly
- Effect is that light is reflected equally in all directions


Directional lighting




- All rays are parallel
- Equivalent to an infinitely distant point source

Source: N. Snavely

Diffuse reflection






$$R_e = k_d \mathbf{N} \cdot \mathbf{L} R_i$$
image intensity of $\mathbf{P} \longrightarrow I = k_d \mathbf{N} \cdot \mathbf{L}$


Simplifying assumptions we'll often make:

- $I = R_e$: "camera response function" is the identity
 - In actual cameras it is a nonlinear function
 - Can always achieve this in practice by inverting it
- $R_i = 1$: light source intensity is 1
 - Can achieve this by dividing each pixel in the image by R_i




Source: [Debevec & Malik 1997]


Other BRDFs



Non-smooth-surfaced materials



Shape from shading



$$I = \kappa_d N \cdot L$$

Assume k_d is 1 for now.

What can we measure from one image?

• $\cos^{-1}(I)$ is the angle between N and L

Add assumptions:

- Constant albedo
- A few known normals (e.g. silhouettes)
- Smoothness of normals

In practice, this doesn't work well: assumptions are too restrictive, too much ambiguity in nontrivial scenes.

An ambiguity that artists exploit



[Belhumeur et al. "The Bas-Relief Ambiguity", 1999


Contours provide extra shape information

Consider points on the occluding contour: $N_z = 0$ N_z positive N_z negative Image Projection direction (z)

Application: finding the direction of the light source

$$I(x,y) = \mathbf{N}(x,y) \cdot \mathbf{S}$$

Full 3D case:



For points on the occluding contour, $N_z = 0$:

$$\begin{pmatrix} N_{x}(x_{1}, y_{1}) & N_{y}(x_{1}, y_{1}) \\ N_{x}(x_{2}, y_{2}) & N_{y}(x_{2}, y_{2}) \\ \vdots & \vdots & \\ N_{x}(x_{n}, y_{n}) & N_{y}(x_{n}, y_{n}) \end{pmatrix} \begin{pmatrix} S_{x} \\ S_{y} \end{pmatrix} = \begin{pmatrix} I(x_{1}, y_{1}) \\ I(x_{2}, y_{2}) \\ \vdots \\ I(x_{n}, y_{n}) \end{pmatrix}$$

Finding the direction of the light source



Application: detecting image splices

Real photo


Fake photo





Can we make things easier?

Photometric stereo










27



Source: N. Snavely

Photometric stereo



Can write this as a linear system and solve:


$$\begin{bmatrix} I_1 \\ I_2 \\ I_3 \end{bmatrix} = k_d \begin{bmatrix} \mathbf{L_1}^T \\ \mathbf{L_2}^T \\ \mathbf{L_3}^T \end{bmatrix} \mathbf{N}$$

Photometric Stereo


Input

Recovered Recovered surface Recovered normal field albedo model 0.5 X


Photometric Stereo



Video photometric stereo




What if we don't know the BRDF?



Clear Elastomer


[Johnson and Adelson, 2009]









Photometric stereo reconstruction







e.g., GelSight [Johnson and Adelson, 2009], [Yuan et al., 2017]



GelSight [Johnson et al. 2009]




Vision-based touch sensing




GelSight [Johnson and Adelson, 2009], [Yuan et al., 2017] Example from [Yuan, Zhu, Owens, Srinivasan, Adelson, "Shape-independent Hardness Estimation...", ICRA 2017]

Robotic touch







Today

- Shape from shading
- Intrinsic image decomposition
- Color perception

What about paint?



$$I = k_d \mathbf{N} \cdot \mathbf{L}$$

 k_d is reflectance or albedo

Intrinsic image decomposition




Reflectance




Shading

Intrinsic image decomposition



44

Intrinsic image decomposition



45

Retinex: keep only large magnitude edges




Input



Estimated reflectance

CNN-based reflectance estimation

Input



Shading

[Bell et al., "Intrinsic images in the wild", 2014]


Applications of intrinsic image decomposition




Today


- Shape from shading
- Intrinsic image decomposition
- Color perception

Color names for cartoon spectra




Additive color mixing






When colors combine by adding the color spectra. Example color displays that follow this mixing rule: CRT phosphors, multiple projectors aimed at a screen.

Red and green make...



Yellow!

Real-world spectra



Source: Torralba, Isola, Freeman, FCV book


How is color perceived in the eye?

Cones


cone-shaped less sensitive operate in high light color vision

Rods

rod-shaped
highly sensitive
operate at night
gray-scale vision




Distribution of Rods and Cones




54


Source: A. Efros ?

Human eye photoreceptor spectral sensitivities






How we sense light spectra




<u>Biophysics:</u> integrate the response over all wavelengths, weighted by the photosensor's sensitivity at each wavelength.


Mathematically: take dot product of input spectrum with the cone sensitivity basis vectors. Project the highdimensional test light into a 3D space.




Cone response curves as basis vectors in a 3D subspace of light power spectra





57








4.10 THE COLOR-MATCHING EXPERIMENT. The observer views a bipartite field and adjusts the intensities of the three primary lights to match the appearance of the test light. (A) A top view of the experimental apparatus. (B) The appearance of the stimuli to the observer. After Judd and Wyszecki, 1975.
















Metameric lights



4.11 METAMERIC LIGHTS. Two lights with these spectral power distributions appear identical to most observers and are called metamers. (A) An approximation to the spectral power distribution of a tungsten bulb. (B) The spectral power distribution of light emitted from a conventional television monitor whose three phosphor intensities were set to match the light in panel A in appearance.


Color blindness

- Classical case: 1 type of cone is missing (e.g. red)
- Makes it impossible to distinguish some spectra.
- There are also tetrachromats, who have 4 cones.




Source: F. Durand


Light sources




https://en.wikipedia.org/wiki/Color_temperature

Same scene under different illuminations








https://en.wikipedia.org/wiki/Color_balance

White balancing




68

White balancing



69


Color constancy



Dale Purves, R. Beau Lotto, Surajit Nundy, "Why We See What We Do."

The dress





Two interpretations

Source: Wikipedia

Next classes: advanced topics