

Optimizing Scrip Systems

IAN A. KASH, ERIC J. FRIEDMAN, and JOSEPH Y. HALPERN

Cornell University

A model of providing service in a P2P network is analyzed. It is shown that by adding a scrip system, a mechanism that admits a reasonable Nash equilibrium that reduces free riding can be obtained. The effect of varying the total amount of money (scrip) in the system on efficiency (i.e., social welfare—the total utility of all the agents in the system) is analyzed, and it is shown that by maintaining the appropriate ratio between the total amount of money and the number of agents, efficiency is maximized. The work has implications for many online systems, not only P2P networks but also a wide variety of online forums for which scrip systems are popular, but formal analyses have been lacking. The tools we use in the analysis give us techniques for analyzing the effects of and dealing with hoarders, altruists, sybils, and collusion, as well as techniques for inferring the characteristics of agents in a system can be inferred from the equilibrium distribution of money.

Categories and Subject Descriptors: C.2.4 [Computer-Communication Networks]: Distributed Systems; I.2.11 [Artificial Intelligence]: Distributed Artificial Intelligence—*Multiagent systems*; J.4 [Social and Behavioral Sciences]: Economics; K.4.4 [Computers and Society]: Electronic Commerce

General Terms: Economics, Theory

Additional Key Words and Phrases: Game Theory, P2P Networks, Scrip Systems, Artificial Currency

1. INTRODUCTION

Historically, non-governmental organizations have issued their own currencies for a wide variety of purposes. These currencies, known as *scrip*, have been used in company towns where government issued currency was scarce [Timberlake 1987], in Washington DC to reduce the robbery rate of bus drivers [Washington Metropolitan Area Transit Commission 1970], and in Ithaca (New York) to promote fairer pay and improve the local economy [Ithaca Hours Inc. 2005]. Scrip systems are also becoming more prevalent in online systems.

To give some examples, market-based solutions using scrip systems have been suggested for applications such as system-resource allocation [Miller and Drexler 1988], managing replication and query optimization in a distributed database [Stonebraker et al. 1996], and allocating experimental time on a wireless sensor network test bed [Chun et al. 2005]; a number of sophisticated scrip systems have been proposed [Gupta et al. 2003; Ioannidis et al. 2002; Vishnumurthy et al. 2003] to allow agents to pool resources while avoiding what is known as *free riding*, where agents take advantage of the resources the system provides while providing none of their own (as Adar and Huberman [2000] have shown, this behavior certainly takes place in

Address?...

Permission to make digital/hard copy of all or part of this material without fee for personal or classroom use provided that the copies are not made or distributed for profit or commercial advantage, the ACM copyright/server notice, the title of the publication, and its date appear, and notice is given that copying is by permission of the ACM, Inc. To copy otherwise, to republish, to post on servers, or to redistribute to lists requires prior specific permission and/or a fee.

© 20YY ACM 0004-5411/YY/00-0001 \$5.00

systems such as Gnutella); and Yootles [Reeves et al.] uses a scrip system as a way of helping groups make decisions using economic mechanisms without involving real money.

Creating a scrip system creates a new market where goods and services can be exchanged that may have been impractical or undesirable to implement with standard currency. However, the potential benefits of a scrip system will not necessarily be realized simply by creating the framework to support one. The story of the Capitol Hill Baby Sitting Co-op [Sweeney and Sweeney 1977], popularized by Krugman [1999], provides a cautionary tale. The Capitol Hill Baby Sitting Co-op was a group of parents working on Capitol Hill who agreed to cooperate to provide babysitting services to each other. In order to enforce fairness, they issued a supply of scrip with each coupon worth a half hour of babysitting. At one point, the co-op had a recession. Many people wanted to save up coupons for when they wanted to spend an evening out. As a result, they went out less and looked for more opportunities to babysit. Since a couple could earn coupons only when another couple went out, no one could accumulate more, and the problem only got worse.

After a number of failed attempts to solve the problem, such as mandating a certain frequency of going out, the co-op started issuing more coupons. The results were striking. Since couples had a sufficient reserve of coupons, they were more comfortable spending them. This in turn made it much easier to earn coupons when a couple's supply got low. Unfortunately, the amount of scrip grew to the point that most of the couples felt "rich." They had enough scrip for the foreseeable future, so naturally they didn't want to devote their evening to babysitting. As a result, couples who wanted to go out were unable to find another couple willing to babysit.

This anecdote shows that the amount of scrip in circulation can have a significant impact on the effectiveness of a scrip system. If there is too little money in the system, few agents will be able to afford service. At the other extreme, if there is too much money in the system, people feel rich and stop looking for work. Both of these extremes lead to inefficient outcomes. This suggests that there is an optimal amount of money, and that nontrivial deviations from the optimum towards either extreme can lead to significant degradation in the performance of the system.

In this paper, we provide a formal model in which to analyze scrip systems. We describe a simple scrip system and show that, under reasonable assumptions, for each fixed amount of money there is a nontrivial equilibrium involving *threshold strategies*, where an agent accepts a request if he has less than $\$k$ for some threshold k .¹

An interesting aspect of our analysis is that, in equilibrium, the distribution of users with each amount of money is the distribution that minimizes relative entropy to an appropriate distribution (subject to the money supply constraint). This allows us to use techniques from statistical mechanics to explicitly compute the distribution of money and thus agents' best-reply functions.

An understanding of agents' best-reply functions allows us to compute the money supply that maximizes social welfare, given the number of agents. As we show,

¹Although we refer to our unit of scrip as the dollar, these are not real dollars nor do we view them as convertible to dollars.

adding more money decreases the equilibrium number of agents with no money, thus increasing social welfare. However, this only works up to a point. Once a critical amount of money is reached, the system experiences a monetary crash: there is so much money that, in equilibrium, everyone will feel rich and no agents are willing to work. We show that, to get optimal performance, we want the total amount of money in the system to be as close as possible to the critical amount, but not to go over it. If the amount of money in the system is over the critical amount, we get the worst possible equilibrium, where all agents have utility 0. This provides a significant tradeoff between efficiency and robustness.

Our equilibrium analysis assumes that all agents have somewhat similar motivation: in particular, they do not get pleasure simply from performing a service, and are interested in money only to the extent that they can use it to get services performed. But in real systems, not all agents have this motivation. Some of the more common “nonstandard” agents are *altruists* and *hoarders*. Altruists are willing to satisfy all requests, even if they go unpaid; hoarders value scrip for its own sake and are willing to accumulate amounts far beyond what is actually useful. Studies of the Gnutella peer-to-peer file-sharing network have shown that one percent of agents satisfy fifty percent of the requests [Adar and Huberman 2000; Hughes et al. 2005]. These agents are doing significantly more work for others than they will ever have done for them, so can be viewed as altruists. Anecdotal evidence also suggests that the introduction of any sort of currency seems to inspire hoarding behavior on the part of some agents, regardless of the benefit of possessing money. For example, SETI@home has found that contributors put in significant effort to make it to the top of their contributor rankings. This has included returning fake results of computations rather than performing them [Zhao et al. 2005].

Altruists and hoarders have opposite effects on a system: having altruists has the same effect as adding money; having hoarders is essentially equivalent to removing it. With enough altruists in the system, the system has a monetary crash, in the sense that standard agents stop being willing to provide service, just as when there is too much money in the system. We show that, until we get to the point where the system crashes, the utility of standard agents is improved by the presence of altruists. We show that the presence of altruists makes the critical point lower than it would without them. Thus, a system designer trying to optimize the performance of the system by making the money supply as close as possible to the critical point (but under it, since being over it would result in a “crash”) needs to be careful about estimating the number of altruists.

Similarly, we show that, as the fraction of hoarders increases, standard agents begin to suffer because there is effectively less money in circulation. On the other hand, hoarders can improve the stability of a system. Since hoarders are willing to accept an infinite amount of money, they can prevent the monetary crash that would otherwise occur as money was added. In any case, our results show how the presence of both altruists and hoarders can be mitigated by appropriately controlling the money supply.

Beyond nonstandard agents, we also consider two different manipulative behaviors in which standard agents may engage: creating multiple identities, known as *sybils* [Douceur 2002], and collusion. In scrip systems where each new user is given

an initial amount of scrip, there is an obvious benefit to creating sybils. Even if this incentive is removed, sybils are still useful: they can be used to increase the likelihood that an agent will be asked to provide service, which makes it easier for him to earn money. This increases the utility of the sybiling agent, at the expense of other agents. From the perspective of an agent considering creating sybils, the first few sybils can provide him with a significant benefit, but the benefits of additional sybils rapidly diminish. So if a designer can make sybiling moderately costly, the number of sybils actually created by rational agents will usually be relatively small.

If a small fraction of agents have sybils, the situation is more subtle. Agents with sybils still do better than those without, but the situation is not zero-sum. In particular, even agents without sybils might be better off, due to having more opportunities to earn money. Somewhat surprisingly, sybils can actually result in everyone being better off. However, exploiting this fact is generally not desirable. The same process that leads to an improvement in social welfare can also lead to a monetary crash, where all agents stop providing service. The system designer can achieve the same effects by increasing the average amount of money or biasing the volunteer selection process. In practice, it seems better to do this than to exploit the possibility of sybils.

In our setting, having sybils is helpful because it increases the likelihood that an agent will be asked to provide service. Our analysis of sybils applies no matter how this increase in likelihood occurs. In particular, it applies to advertising. Thus, our results suggest that there are tradeoffs involved in allowing advertising. For example, many systems allow agents to announce their connection speed and other similar factors. If this biases requests towards agents with high connection speeds, even when agents with lower connection speeds are perfectly capable of satisfying a particular request, then agents with low connection speeds will have an unnecessarily worsened experience in the system. This also means that such agents will have a strong incentive to lie about their connection speed.

While collusion is generally a bad thing, in the context of scrip systems with fixed prices, it is almost entirely positive. Without collusion, if a user runs out of money he is unable to request service until he is able to earn some. However, a colluding group can pool their money so that all members can make a request whenever the group as a whole has some money. This increases welfare for the agents who collude because agents who have no money receive no service. Collusion tends to benefit the non-colluding agents as well. Since colluding agents work less often, it is easier for everyone to earn money, which ends up making everyone better off. However, as with sybils, collusion does have the potential of crashing the system if the average amount of money is high.

While a designer should generally encourage collusion, we would expect that in most systems there will be relatively little collusion and what collusion exists will involve small numbers of agents. After all, scrip systems exist to try and resolve resource-allocation problems where agents are competing with each other. If they could collude to optimally allocate resources within the group, they would not need a scrip system in the first place. However, many of the benefits of collusion come from agents being allowed to effectively have a negative amount of money (by borrowing from their collusive partners). These benefits could also be realized if

agents are allowed to borrow money, so designing a loan mechanism could be an important improvement for a scrip system. Of course, implementing such a loan mechanism in a way that prevents abuse requires a careful design.

In order to effectively utilize our results, a system designer needs to be able to identify characteristics of agents (with what frequency do they make requests, how likely are they to be chosen to satisfy a request, and so on) and what strategies they are following. This is particularly useful because finding an amount of money close to the point of monetary crash, but not past it, relies on an understanding of the agents in the system. Of course, such information is also of great interest to social scientists and marketers. We show how relatively simple observations of the system can be used to infer this information.

The rest of the paper is organized as follows. In Section 2, we review related work. Then in Section 3, we present the formal model. We analyze the distribution of money in this model when agents are using threshold strategies in Section 4, and show that it is characterized by relative entropy. Using this analysis, we show in Section 5 that, under minimal assumptions, there is a nontrivial equilibrium where all agents use threshold strategy strategies. These results apply to a sufficiently large population of agents after a sufficiently long period of time, so in Section 6 we use simulations to demonstrate that these values are reasonable in practice. We begin applying our analysis in Section 7, where we show that the analysis leads to an understanding of how to choose the amount of money in the system (or, equivalently, the cost to fulfill a request) so as to maximize efficiency, and also shows how to handle new users. In Section 8, we discuss how the model can be used to understand the effects of altruists, hoarders, sybils, and collusion and provide guidance about how system designers can handle these user behaviors. All of this guidance relies on being able to understand what strategies agents are using and what their preferences are. In Section 9, we discuss how these can be inferred by examining the system. We conclude in Section 10.

2. RELATED WORK

Scrip systems have a long history in computer science, with two main thrusts: resource allocation and free-riding prevention. Early applications for resource allocation include agoric systems [Miller and Drexler 1988], which envisioned solving problems such as processor scheduling using markets, and Mariposa [Stonebraker et al. 1996], a market-driven query optimizer for distributed databases. More recently, scrip systems have been used to allocate the resources of research testbeds. Examples include Mirage [Chun et al. 2005] for wireless sensor networks, Bellagio [AuYoung et al. 2007] for PlanetLab, and Egg [Brunelle et al. 2006] for grid computing. Virtual markets have been used to coordinate the activity of nodes of a sensor network [Mainland et al. 2004]. Yootles [Reeves et al.] uses a scrip to help people make everyday decisions, such as where to have lunch, without involving real money.

Systems that use scrip to prevent free riding include KARMA [Vishnumurthy et al. 2003], which provides a general framework for P2P networks. Gupta et al. [2003] propose what they call a “debit-credit reputation computation” for P2P networks, which is essentially a scrip system. Fileteller [Ioannidis et al. 2002] uses

payments in a network file storage system. Dandelion [Sirivianos et al. 2007] uses scrip in a content distribution setting. Belenkiy et al. [Belenkiy et al. 2007] consider how a BitTorrent-like system can make use of e-cash. Antfarm [Peterson and Siler 2009] uses scrip to optimize content distribution across a number of BitTorrent-like swarms.

Despite this tremendous interest in scrip systems, there has been relatively little work studying how they behave. Chun et al. [2005] studied user behavior in a deployed scrip system and observed that users tried various (rational) manipulations of the auction mechanism used by the system. Their observations suggest that system designers will have to deal with game-theoretic concerns.

Hens et al. [2007] do a theoretical analysis of what can be viewed as a scrip system in a related model. There are a number of significant differences between the models. First, in the Hens et al. model, there is essentially only one type of agent, but an agent's utility for getting service (our γ_i) may change over time. Thus, at any time, there will be agents who differ in their utility. At each round, we assume that one agent is chosen (by nature) to make a request for service, while other agents decide whether or not to provide it. In the Hens et al. model, at each round, each agent decides whether to provide service, request service, or opt out, as a function of his utilities and the amount of money he has. They assume that there is no cost for providing service and everyone is able to provide service. Under this assumption, a system with optimal performance is one where half the agents request service and the other half are willing to provide it. Despite these differences, Hens et al. also show that agents will use a threshold strategy. However, although they have inefficient equilibria, because there is no cost for providing service, their model does not exhibit the monetary crashes that our model can exhibit.

Aperjis and Johari [2006] examine a model of a P2P filesharing system as an exchange economy. They associate a price (in bandwidth) with each file and find a market equilibrium in the resulting economy. However, they do not use an explicit currency.

The effects of altruists, sybils, and collusion on system behavior have all been studied in other contexts. Work on the evolution of cooperation stresses the importance of altruists willing to undertake costly punishment [Hauert et al. 2007]. Yokoo et al. [2004] studied the effects of sybils in auctions. Solution concepts such as *strong Nash equilibrium* [Aumann 1959] and *k-t robust equilibrium* [Abraham et al. 2006] explicitly address collusion in games; Hayrapetyan et al. [2006] study collusion in congestion games and find cases where, as with scrip systems, collusion is actually beneficial.

The ultimate goal of a scrip systems is to promote cooperation. While there is limited theoretical work on scrip systems, there is a large body of work on cooperation. Much of the work involves a large group of agents being randomly matched to pay a game such as prisoner's dilemma. Such models were studied in the economics literature [Kandori 1992; Ellison 1994] and first applied to online reputations in [Friedman and Resnick 2001]; Feldman et al. [2004] apply them to P2P systems.

These models fail to capture important asymmetries in the interactions of the agents. When a request is made, there are typically many people in the network who can potentially satisfy it (especially in a large P2P network), but not all can.

For example, some people may not have the time or resources to satisfy the request. The random-matching process ignores the fact that some people may not be able to satisfy the request. (Presumably, if the person matched with the requester could not satisfy the match, he would have to defect.) Moreover, it does not capture the fact that the decision as to whether to “volunteer” to satisfy the request should be made before the matching process, not after. That is, the matching process does not capture the fact that if someone is unwilling to satisfy the request, there will doubtless be others who can satisfy it. Finally, the actions and payoffs in prisoner’s dilemma game do not obviously correspond to actual choices that can be made. For example, it is not clear what defection on the part of the requester means. Our model addresses all these issues.

Scrip systems are not the only approach to preventing free riding. Two other approaches often used in P2P networks are barter and reputation systems. Perhaps the best-known example of a system that uses barter is BitTorrent [Cohen 2003], where clients downloading a file try to find other clients with parts they are missing so that they can trade, thus creating a roughly equal amount of work. Since the barter is restricted to users currently interested in a single file, this works well for popular files, but tends to have problems maintaining availability of less popular ones. An example of a barter-like system built on top of a more traditional file-sharing system is the credit system used by eMule. Each user tracks his history of interactions with other users and gives priority to those he has downloaded from in the past. However, in a large system, the probability that a pair of randomly-chosen users will have interacted before is quite small, so this interaction history will not be terribly helpful. Anagnostakis and Greenwald [2004] present a more sophisticated version of this approach, but it still seems to suffer from similar problems.

A number of attempts have been made at providing general reputation systems (e.g. [Guha et al. 2004; Gupta et al. 2003; Kamvar et al. 2003; Xiong and Liu 2002]). The basic idea is to aggregate each user’s experience into a global number for each individual that intuitively represents the system’s view of that individual’s reputation. However, these attempts tend to suffer from practical problems because they implicitly view users as either “good” or “bad”, assume that the “good” users will act according to the specified protocol, and that there are relatively few “bad” users. Unfortunately, if there are easy ways to game the system, once this information becomes widely available, rational users are likely to make use of it. We cannot count on only a few users being “bad” (in the sense of not following the prescribed protocol). For example, Kazaa uses a measure of the ratio of the number of uploads to the number of downloads to identify good and bad users. However, to avoid penalizing new users, they gave new users an average rating. Users discovered that they could use this relatively good rating to free ride for a while and, once it started to get bad, they could delete their stored information and effectively come back as a “new” user, thus circumventing the system (see [Anagnostakis and Greenwald 2004] for a discussion and [Friedman and Resnick 2001] for a formal analysis of this “whitewashing”). Thus, Kazaa’s reputation system is ineffective.

3. THE MODEL

Before specifying our model formally, we give an intuitive description of what our model captures. We model a scrip system where, as in the babysitting co-op, agents provide each other with service. There is a single service (babysitting) that agents occasionally want. In practice, at any given time, a number of agents will want service, but to simplify the formal description and analysis we model the scrip system as proceeding in a series of rounds where, in each round, a single agent wants service (the time between rounds will be adjusted to capture the growth in parallelism as the number of agents grows).² In each round, after an agent requests service, other agents have to decide whether they want to volunteer to provide service. However, not all agents may be able to satisfy the request (not everyone can babysit every night). While, in practice, the ability of agents to provide service at various times may be correlated for a number of reasons (some agents might have very young children that only certain agents are willing to babysit; being unavailable in one round might be correlated with being unavailable in the next round; and so on), for simplicity we model the ability to provide service using a single probability, and assume that the events of an agent being able to provide service in different rounds or two agents being able to provide service in the same or different rounds are independent. If there is at least one volunteer, someone is chosen from among the volunteers (uniformly at random) to satisfy the request. The requester then gains some utility (he was able to go out because he had a babysitter) and the volunteer loses some utility (people would rather do something other than babysit), and the requester pays the volunteer a fee that we fix at one dollar. As is standard in the literature, we assume that agents discount future payoffs. This captures the intuition that a util now is worth more than a util tomorrow, and allows us to compute the total utility derived by an agent in an infinite game. The amount of utility gained by having a service performed and the amount lost by performing it, as well as many other parameters may depend on the agent.

More formally, we assume that agents have a *type* t drawn from some finite set T of types. We can describe the entire population of agents using the pair (T, \vec{f}) , where \vec{f} is a vector of length $|T|$ and f_t is the fraction with type t . For most of the paper, we consider only what we call *standard agents*. These are agents who derive no pleasure from performing a service, and for whom money has no intrinsic value. We can characterize the type of a standard agent by a tuple $t = (\alpha_t, \beta_t, \gamma_t, \delta_t, \rho_t, \chi_t)$, where

- $\alpha_t > 0$ reflects the cost for an agent of type t to satisfy a request;
- $0 < \beta_t \leq 1$ is the probability that an agent of type t can satisfy a request;
- $\gamma_t > \alpha_t$ is the utility that an agent of type t gains for having a request satisfied;
- $0 < \delta_t < 1$ is the rate at which an agent of type t discounts utility;
- $\rho_t > 0$ represents the (relative) request rate (some people need babysitting more often than others). For example, if there are two types of agents with $\rho_{t_1} = 2$

²For large numbers of agents, our model converges to one in which agents make requests in real time, and the time between an agent's requests is exponentially distributed. In addition, the time between requests served by a single player is also exponentially distributed.

and $\rho_{t_2} = 1$ then agents of the first type will make requests twice as often as agents of the second type. Since these request rates are relative, we can multiply them all by a constant to normalize them. To simplify later notation, we assume the ρ_t are normalized so that $\sum_{t \in T} \rho_t f_t = 1$; and

- $\chi_t > 0$ represents the (relative) likelihood of an agent to be chosen when he volunteers (some babysitters may be more popular than others). In particular, this means the relative probability of two given agents being chosen is independent of which other agents volunteer.
- $\omega_t = \beta_t \chi_t / \rho_t$ is not part of the tuple, but is an important derived parameter that, as we will see in Section 4, helps determine how much money an agent will have.

Representing the population of agents in a system as (T, \vec{f}) captures the essential features of a scrip system we want to model: there are a large number of agents who may have different types. Note that some tuples (T, \vec{f}) may be incompatible with there being some number N of agents. For example, if there are two types, and \vec{f} says that half of the agents are of each type, then there cannot be 101 agents. Similar issues arise when we want to talk about the amount of money in example, by specifying how to assign to types to agents in a way that we could deal with this problem in a number of ways (for example, by having each agent determine his type at random according to the distribution \vec{f}). For convenience, we simply do not consider population sizes that are incompatible with \vec{f} . This is the approach used in the literature on *N-replica economies* [Mas-Colell et al. 1995].

Formally, we consider games specified by a tuple (T, \vec{f}, h, m, n) , where T and \vec{f} are as defined above, $h \in \mathbb{N}$ is the *base* number of agents of each type, $n \in \mathbb{N}$ is number of *replicas* of these agents and $m \in \mathbb{R}^+$ is the average amount of money. The total number of agents is thus hn . We ensure that the number of agents of type t is exactly f_t and that the average amount of money is exactly m by requiring that $f_t h \in \mathbb{N}$ and $m h \in \mathbb{N}$. Having created a base population satisfying these constraints, we can make an arbitrary number of copies of it. More precisely, we assume that agents $0 \dots f_{t_1} h - 1$ have type t_1 , agents $f_{t_1} h \dots (f_{t_1} + f_{t_2}) h - 1$ have type t_2 , and so on through agent $h - 1$. These base agents determine the types of all other agents. Each agent $j \in \{h, \dots, hn - 1\}$ has the same type as $j \bmod h$; that is, all the agents of the form $j + kh$ for $k = 1, \dots, n - 1$ are replicas of agent j . At the start of the game, we initially allocate each of the hmn dollars in the system to an agent chosen uniformly at random.

We now describe (T, \vec{f}, h, m, n) as an infinite extensive-form game. A non-root node in the game tree is associated with a round number (how many requests have been made so far), a phase number, either 1, 2, 3, or 4 (which describes how far along we are in determining the results of the current request), a vector \vec{x} where x_i is the current amount of money agent i has, and $\sum_i x_i = mhn$, and, for some nodes, some additional information whose role will be made clear below. We use $\tau(i)$ to denote the type of agent i .

- The game starts at a special root node, denoted Λ , where nature moves. Intuitively, at Λ , nature allocates money uniformly at random, so it transitions to a node of the form $(0, 1, \vec{x})$: round zero, phase one, and allocation of money \vec{x} , and

each possible transition is equally likely.

- At a node of the form $(r, 1, \vec{x})$, nature selects an agent to make a request in the current round. Agent i is chosen with probability $\rho_{\tau(i)}/hn$. If i is chosen, a transition is made to $(r, 2, \vec{x}, i)$.
- At a node of the form $(r, 2, \vec{x}, i)$, nature selects the set V of agents (not including i) able to satisfy the request. Each agent $j \neq i$ is included in V with probability $\beta_{\tau(j)}$. If V is chosen, a transition is made to $(r, 3, \vec{x}, i, V)$.
- At a node of the form $(r, 3, \vec{x}, i, V)$, each agent in V chooses whether to volunteer. If V' is the set of agents who choose to volunteer, a transition is made to $(r, 4, \vec{x}, i, V')$.
- At a node of the form $(r, 4, \vec{x}, i, V')$, if $V' \neq \emptyset$, nature chooses a single agent in V' to satisfy the request. Each agent j is chosen with probability $\chi_{\tau(j)}/\sum_{j' \in V'} \chi_{\tau(j')}$. If j is chosen, a transition is made to $(r+1, 1, \vec{x}')$, where

$$x'_j = \begin{cases} x_j - 1 & \text{if } i = j \text{ and } x_j > 0, \\ x_j + 1 & \text{if } j \text{ is chosen by nature and } x_i > 0, \\ x_j & \text{otherwise.} \end{cases}$$

If $V' = \emptyset$, nature has no choice; a transition is made to $(r+1, 1, \vec{x})$ with probability 1.

A strategy for agent j describes whether or not he will volunteer at every node of the form $(r, 3, \vec{x}, i, V)$ such that $j \in V$. (These are the only nodes where j can move.) A strategy profile \vec{S} consists of one strategy per agent. A strategy profile \vec{S} determines a probability distribution over paths $\Pr_{\vec{S}}$ in the game tree. Each path determines the value of the following random two variables:

- x_i^r , the amount of money agent i has during round r , defined as the value of x_i at the nodes with round number r and
- u_i^r , the utility of agent i for round r . If i is a standard agent, then

$$u_i^r = \begin{cases} \gamma_{\tau(i)} & \text{if a node } (r, 4, \vec{x}, i, V') \text{ is on the path with } V' \neq \emptyset \\ -\alpha_{\tau(i)} & \text{if } i \text{ is chosen by nature at node } (r, 4, \vec{x}, j, V') \\ 0 & \text{otherwise.} \end{cases}$$

$U_i(\vec{S})$, the total expected utility of agent i if strategy profile \vec{S} is played is the discounted sum of his per round utilities u_i^r , but the exact form of the discounting requires some explanation. As the number of agents increases, we would expect more requests to be made per unit time, and the expected number of requests an agent makes per unit time to be constant. Since only one agent makes a request per round, it seems that a reasonable way to model this is to take the time between rounds to be $1/n$, where n is the number of agents. The discount rate—which can be thought of as the present value of getting one util one round in the future—has to be modified as well. It turns out that the obvious choice of discount rate, $\delta_t^{1/n}$, is not appropriate. To understand why, consider an agent who has all of his requests satisfied. When there are just h agents, he is chosen to make a request each round with probability ρ_t/h . His total expected utility with a discount rate of δ is $\sum_{r=0}^{\infty} \delta^r \rho_t \gamma_t / h = (\rho_t \gamma_t / h) / (1 - \delta_t)$. With n replicas, scaling the discount rate as

$\delta_t^{1/n}$ gives $\sum_{r=0}^{\infty} \delta_t^{r/n} \rho_t \gamma_t / (hn) = (\rho_t \gamma_t / (hn)) / (1 - \delta_t^{1/n})$. Thus, using this scaling, the agent's utility for having all his requests satisfied decreases as n increases. This seems unnatural. If we instead use the discount rate $(1 - (1 - \delta_t)/n)$, his expected utility is $\sum_{r=0}^{\infty} (1 - (1 - \delta_t)/n)^r (\rho_t \gamma_t / (hn)) = (\rho_t \gamma_t / (hn)) / (1 - (1 - (1 - \delta_t)/n)) = (\rho_t \gamma_t / h) / (1 - \delta_t)$, which is independent of n , and seems much more reasonable.

Using the discount rate $(1 - (1 - \delta_t)/n)$ solves one problem, but leaves another. A larger δ_t is meant to reflect a more patient agent, who gives future utility a higher weight. However, as the preceding equation shows, increasing δ_t also increases total utility. To counteract this, we multiply the total discounted sum by $(1 - \delta_t)$. This is standard in economics, for example in the folk theorem for repeated games [Fudenberg and Tirole 1991]. With these considerations in mind, the total expected utility of agent i given the vector of strategies \vec{S} is

$$U_i(\vec{S}) = (1 - \delta_{\tau(i)}) \sum_{r=0}^{\infty} (1 - (1 - \delta_{\tau(i)})/n)^r E_{\vec{S}}[u_i^r], \quad (1)$$

In modeling the game this way, we have implicitly made a number of assumptions. For example, we have assumed that all of agent i 's requests that are satisfied give agent i the same utility, and that prices are fixed. We discuss the implications of these assumptions in Section 10.

Our solution concept is the standard notion of an approximate Nash equilibrium. As usual, given a strategy profile \vec{S} and agent i , we use (S'_i, \vec{S}_{-i}) to denote the strategy profile that is identical to \vec{S} except that agent i uses S'_i .

Definition 3.1. A strategy S'_i for agent i is an ϵ -best reply to a strategy profile \vec{S}_{-i} for the agents other than i in the game (T, \vec{f}, h, m, n) if, for all strategies S''_i ,

$$U_i(S''_i, \vec{S}_{-i}) \leq U_i(S'_i, \vec{S}_{-i}) + \epsilon.$$

Definition 3.2. A strategy profile \vec{S} for the game (T, \vec{f}, h, m, n) is an ϵ -Nash equilibrium if for all agents i , S_i is an ϵ -best reply to \vec{S}_{-i} . A Nash equilibrium is an epsilon-Nash equilibrium with $\epsilon = 0$.

As we show in Section 5, (T, \vec{f}, h, m, n) has equilibria where agents use a particularly simple type of strategy, called a *threshold strategy*. Intuitively, an agent with “too little” money will want to work, to minimize the likelihood of running out due to making a long sequence of requests before being able to earn more money. On the other hand, a rational agent with plenty of money will think it is better to delay working, thanks to discounting. These intuitions suggest that the agent should volunteer if and only if he has less than a certain amount of money. Let s_k be the strategy where an agent volunteers if and only if the requester has at least 1 dollar and the agent has less than k dollars. Note that s_0 is the strategy where the agent never volunteers. While everyone playing s_0 is a Nash equilibrium (nobody can do better by volunteering if no one else is willing to), it is an uninteresting one.

We frequently consider the situation where each agent of type t uses the same threshold s_{k_t} . In this case, a single vector \vec{k} suffices to indicate the threshold of each type, and we can associate with this vector the strategy $\vec{S}(\vec{k})$ where $\vec{S}(\vec{k})_i = s_{k_{\tau(i)}}$ (i.e., agent i of type $\tau(i)$ uses threshold $k_{\tau(i)}$).

For the rest of this paper, we focus on threshold strategies (and show why it is reasonable to do so). When we consider the threshold strategy $\vec{S}(\vec{k})$, for ease of exposition, we assume in our analysis that $mhn < \sum_t f_t k_t hn$. To understand why, note that mhn is the total amount of money in the system. If $mhn \geq \sum_t f_t k_t hn$, then if the agents use a threshold $\vec{S}(\vec{k})$, the system will quickly reach a state where each agent has k_t dollars, so no agent will volunteer. This is equivalent to all agents using a threshold of 0, and similarly uninteresting.

4. ANALYZING THE DISTRIBUTION OF WEALTH

Our main goal is to show that there exists an approximate equilibrium where all agents play threshold strategies. In this section, we examine a more basic question: if all agents play a threshold strategy, what happens? We show that there is some distribution over money (i.e., a distribution that describes what fraction of people have each amount of money) such that the system “converges” to this distribution in a sense to be made precise shortly. In addition to providing an understanding of system behavior that underpins our later results, this result also provides a strong guarantee about the stability of the economy.

Suppose that all agents of each type t use the same threshold k_t , so we can write the vector of thresholds as \vec{k} . For simplicity, assume that each agent has at most k_t dollars. We can make this assumption with essentially no loss of generality, since if someone has more than k_t dollars, he will just spend money until he has at most k_t dollars. After this point he will never acquire more than k_t . Thus, eventually the system will be in a state where, for all types t , no agent of type t has more than k_t dollars.

We are interested in the vectors \vec{x}^r that can be observed in round r (recall that x_i^r is the amount of money that agent i has at round r). By assumption, if agent i has type $\tau(i)$, then $x_i^r \in \{0, \dots, k_{\tau(i)}\}$. In addition, since the total amount of money is hmn ,

$$\vec{x}^r \in X_{T, \vec{f}, h, m, n, \vec{k}} = \{\vec{x} \in \mathbb{N}^{hn} \mid \forall i. x_i \leq k_{\tau(i)}, \sum_i x_i = hmn\}.$$

The evolution of \vec{x}^r can be described by a Markov chain $\mathcal{M}_{T, \vec{f}, h, m, n, \vec{k}}$ over the state space $X_{T, \vec{f}, h, m, n, \vec{k}}$. For brevity, we refer to the Markov chain and state space as \mathcal{M} and X , respectively, when the subscripts are clear from context. It is possible to move from state s to state s' in a single round if, by choosing a particular agent i to make a request and another agent j to satisfy it, i 's amount of money in s' is 1 more than in s ; j 's amount of money in s' is 1 less than in s , and all other agents have the same amount of money in s and s' . Therefore, the probability of a transition from a state \vec{x} to \vec{y} is 0 unless there exist two agents i and j such that $\vec{y}_{i'} = \vec{x}_{i'}$ for all $i' \notin \{i, j\}$, $\vec{y}_i = \vec{x}_i + 1$, and $\vec{y}_j = \vec{x}_j - 1$. In this case, the probability of transitioning from \vec{x} to \vec{y} is the probability of j being chosen to make a request and i being chosen to satisfy it. Let $\Delta_{\vec{f}, m, \vec{k}}$ denote the set of probability distributions d on $\cup_{t \in T} \{t\} \times \prod_t \{0, \dots, k_t\}$ such that for all types t , $\sum_{l=0}^{k_t} d(t, l) = f_t$. We can think of $d(t, l)$ as the fraction of agents of type t that have l dollars. We can associate each state \vec{x} with its corresponding distribution $d^{\vec{x}}$. This is a useful way of looking at the system, since we typically just care about the fraction of people with each

amount of money, not the amount that each particular agent has. We show that, if n is large, then there is a distribution $d^* \in \Delta_{\vec{f}, m, \vec{k}}$ such that, after a sufficient amount of time, the Markov chain \mathcal{M} is almost always in a state \vec{x} such that $d^{\vec{x}}$ is close to d^* . Thus, agents can base their decisions about what strategy to use on the assumption that they will be in a state where the distribution of money is essentially d^* .

We can in fact completely characterize the distribution d^* . Given two distributions $d, q \in \Delta_{\vec{f}, m, \vec{k}}$, let

$$H(d||q) = - \sum_{\{(t,j):q(t,j) \neq 0\}} d(t,j) \log d(t,j)/q(t,j)$$

denote the *relative entropy* of d relative to q ($H(d||q) = \infty$ if $d(t,j) = 0$ and $q(t,j) \neq 0$ or vice versa); this is also known as the *Kullback-Leibler divergence of q from d* [Cover and Thomas 1991]. If Δ is a closed convex set of distributions, then it is well known that, for each q , there is a unique distribution in Δ that minimizes the entropy relative to q . Since $\Delta_{\vec{f}, m, \vec{k}}$ is easily seen to be a closed convex set of distributions, in particular, this is the case for $\Delta_{\vec{f}, m, \vec{k}}$. We now show that there exists a q such that, for n sufficiently large, the Markov chain \mathcal{M} is almost always in a state \vec{x} such that $d^{\vec{x}}$ is close to the distribution $d_{q, \vec{f}, m}^* \in \Delta_{\vec{f}, m, \vec{k}}$ that minimizes entropy relative to q . (We omit some or all of the subscripts on d^* when they are not relevant.) The statement is correct under a number of senses of “close”. For definiteness, we consider the Euclidean distance. Given $\varepsilon > 0$ and q , let $X_{T, \vec{f}, h, m, n, \vec{k}, \varepsilon, q}$ (or $X_{\varepsilon, q}$, for brevity) denote the set of states $\vec{x} \in X_{T, \vec{f}, h, m, n, \vec{k}}$ such that $\sum_{(t,j)} |d^{\vec{x}}(t,j) - d_q^*|^2 < \varepsilon$.

Let $I_{q, n, \varepsilon}^r$ be the random variable that is 1 if $d^{\vec{x}^r} \in X_{\varepsilon, q}$, and 0 otherwise.

Theorem 4.1. *For all games $(T, \vec{f}, h, m, 1)$, all vectors \vec{k} of thresholds, and all $\varepsilon > 0$, there exist $q \in \Delta_{\vec{f}, m, \vec{k}}$ and n_ε such that, for all $n > n_\varepsilon$, there exists a round r^* such that, for all $r > r^*$, we have $\Pr(I_{q, n, \varepsilon}^r = 1) > 1 - \varepsilon$.*

The proof of Theorem 4.1 can be found in Appendix A. One interesting special case of the theorem is when there exist β, χ , and ρ such that for all types t , $\beta_t = \beta$, $\chi_t = \chi$, and $\rho_t = \rho$. In this case q is the distribution $q(t, j) = f_t / (k_t + 1)$ (i.e., q is uniform within each type t). We sketch the proof for this special case here.

PROOF. (Sketch) Using standard techniques, we can show that our Markov Chain has a *limit distribution* π such that for all \vec{y} , $\lim_{r \rightarrow \infty} \Pr(\vec{x}^r = \vec{y}) = \pi(\vec{y})$. Let $T_{\vec{x}\vec{y}}$ denote the probability of transitioning from state \vec{x} to state \vec{y} . It is easily verified by an explicit computation of the transition probabilities that (in this special case) $T_{\vec{x}\vec{y}} = T_{\vec{y}\vec{x}}$. It is well known that this symmetry implies that π is the uniform distribution [Resnick 1992]. Thus, after a sufficient amount of time, the distribution of \vec{x}^r will be arbitrarily close to uniform.

Since, for large r , $\Pr(\vec{x}^r = \vec{y})$ is approximately $1 / |X|$, the probability of \vec{x}^r being in a set of states is the size of the set divided by the total number of states. Using a straightforward combinatorial argument, it can be shown that the fraction of states not in $X_{\varepsilon, q}$ is bounded by $p(n)/e^{cn}$, where p is a polynomial. This fraction goes to 0 as n gets large. Thus, for sufficiently large n , $\Pr(I_{q, n, \varepsilon}^r = 1) > 1 - \varepsilon$. ■

The last portion of the proof sketch is actually a standard technique from statistical mechanics that involves showing that there is a *concentration phenomenon* around the maximum entropy distribution [Jaynes 1978]. In this special case, when π is the uniform distribution, the number of states corresponding to a particular distribution d is proportional to $e^{nH(d)}$ (where H here is the standard entropy function). In general, each state is not equally likely, which is why the general proof in Appendix A uses relative entropy.

Theorem 4.1 tells us that, after enough time, the distribution of money is almost always close to some d^* , where d^* can be characterized as a distribution that minimizes relative entropy subject to some constraints. Let $q(t, i) = (\omega_t)^i / (\sum_t \sum_{j=0}^{k_t} (\omega_t)^j)$. Then the value of d^* is given by the following lemma.

Lemma 4.1.

$$d^*(t, i) = \frac{f_t \lambda^i q(t, i)}{\sum_{j=0}^{k_t} \lambda^j q(t, j)}, \quad (2)$$

where λ is the unique value such that

$$\sum_t \sum_i i d^*(t, i) = m. \quad (3)$$

The proof of Lemma 4.1 is omitted because it can be easily checked using Lagrange multipliers in the manner of [Jaynes 1978] where the function to be minimized is the entropy of d^* relative to q and the constraints are there a f_t fraction of the agents are of type t and the average amount of money is m .

5. EXISTENCE OF EQUILIBRIA

We have seen that the system is well behaved if the agents all follow a threshold strategy; we now want to show that, if the discount factor δ is sufficiently large for all agents, there is a nontrivial approximate Nash equilibrium where they do so (that is, an approximate Nash equilibrium where all the agents use s_k for some $k > 0$.) To understand why we need δ to be sufficiently large, note that if δ is small, then agents have no incentive to work. Intuitively, if future utility is sufficiently discounted, then all that matters is the present, and there is no point in volunteering to work. Thus, for sufficiently small δ , s_0 is the only equilibrium. To show that there is a nontrivial equilibrium if the discount factor is sufficiently large, we first show that, if every other agent is playing a threshold strategy, then there is an approximate best reply that is also a threshold strategy. Furthermore, we show that the best-reply function is monotone; that is, if some agents change their strategy to one with a higher threshold, no other agent can do better by lowering his threshold. This makes our game one with what Milgrom and Roberts [1900] call *strategic complementarities*. Using results of Tarski [1955], Topkis [1979] showed that there are pure strategy equilibria in such games, since the process of starting with a strategy profile where everyone always volunteers (i.e., the threshold is ∞) and then iteratively computing the best-reply profile to it converges to a Nash equilibrium in pure strategies. This procedure also provides an efficient algorithm for explicitly computing equilibria.

To see that threshold strategies are approximately optimal, consider a single agent i of type t and fix the vector \vec{k} of thresholds used by the other agents. If

we assume that the number of agents is large, what an agent i does has essentially no effect on the behavior of the system (although it will, of course, affect that agent's payoffs). In particular, this means that the distribution q of Theorem 4.1 characterizes the distribution of money in the system. This distribution, together with the vector \vec{k} of thresholds, determines what fraction of agents volunteers at each step. This, in turn, means that from the perspective of agent i , the problem of finding an optimal response to the strategies of the other agents reduces to finding an optimal policy in a Markov decision process (MDP) $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$. The behavior of the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$ depends on two probabilities: p_u and p_d . Informally, p_u is the probability of i earning a dollar during each round if is willing to volunteer, and p_d is the probability that i will be chosen to make a request during each round. Note that p_u depends on m , \vec{k} , and t (although it turns out that p_d depends only on n , the number of agents in the system) and t ; if the dependence of p_u on m , \vec{k} , and/or t is important, we add the relevant parameters to the superscript, writing, for example, $p_u^{m, \vec{k}}$. We show that the optimal policy for i in $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$ is a threshold policy, and that this policy is an ε -optimal strategy for G . Importantly, the same policy is optimal independent of the value of n . This allows us to ignore the exact size of the system in our later analysis.

For many of our later results and discussion, it will be important to understand how p_u , p_d , and t affect the optimal policy for $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$, and thus the ε -optimal strategies in the game. We use this understanding in this section to show that there exist nontrivial equilibria in Lemma 5.3, to show that adding money increases social welfare in Section 7, to understand how agent behaviors affect social welfare in Section 8, and to identify agent types from their behavior in Section 9.

In the following lemma, whose proof (and the relevant formal definitions) are deferred to Appendix B, Equation (4), quantifies the effects of these parameters. When choosing whether he should volunteer with his current amount of money, an agent faces a choice of whether to pay a utility cost of α_t now in exchange for a discounted payoff of γ_t when he eventually spends the resulting dollar. His choice will depend on how much time he expects to pass before he spends that dollar, which in turn depends on his current amount of money k and the probabilities p_u and p_d . The following lemma quantifies this calculation.

Lemma 5.1. *Consider the games $G_n = (T, \vec{f}, h, m, n)$ (where T , \vec{f} , h , and m are fixed, but n may vary). There exists a k such that for all n , s_k is an optimal policy for $\mathcal{P}_{G_n, \vec{s}(\vec{k}), t}$. The threshold k is the maximum value of κ such that*

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u, p_d)}] \gamma_t, \quad (4)$$

where $J(\kappa, p_u, p_d)$ is a random variable whose value is the first round in which an agent starting with κ dollars, using strategy s_κ , and with probabilities p_u and p_d of earning a dollar and of being chosen given that he volunteers, respectively, runs out of money.

The following theorem shows that an optimal threshold policy for $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$ is an ε -optimal strategy for G . In particular, this means that Equation (4) allows us to understand how changing parameters affect an ε -optimal strategy for G , not just for $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$.

Theorem 5.1. *For all games $G = (T, \vec{f}, h, m, n)$, all vectors \vec{k} of thresholds, and all $\varepsilon > 0$, there exist n_ε^* and $\delta_{\varepsilon, n}^*$ such that for all $n > n_\varepsilon^*$, types $t \in T$, and $\delta_t > \delta_{\varepsilon, n}^*$, an optimal threshold policy for $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$ is an ε -best reply to the strategy profile $\vec{S}(\vec{k})_{-i}$ for every agent i of type t .*

We defer the proof of Theorem 5.1 to Appendix B. While, in this and later theorems, the acceptable values of $\delta_{\varepsilon, n}^*$ depend on n , they are independent if, as we suggest in Section 6, the Markov Chain from Section 4 is rapidly mixing.

Given a game $G = (T, \vec{f}, h, m, n)$ and a vector \vec{k} of thresholds, Lemma 5.1 gives an optimal threshold k'_t for each type t . Theorem 5.1 guarantees that $s_{k'_t}$ is an ε -best reply to $\vec{S}_{-i}(\vec{k})$, but does not rule out the possibility of other best replies. However, for ease of exposition, we will call k'_t the best reply to \vec{S}_{-i} and call $BR_G(\vec{k}) = \vec{k}'$ the best-reply function. The following lemma shows that this function is monotone: it is non-decreasing in \vec{k} and non-increasing in m . Along the way, we prove that several other quantities are monotone. First, we show that $\lambda_{m, \vec{k}}$, the value of λ from Lemma 4.1 given m and \vec{k} , is non-decreasing in m and non-increasing in \vec{k} . We use this to show that $p_u^{m, \vec{k}}$ is non-increasing in \vec{k} and non-decreasing in m . We defer the proof to Appendix B.

Lemma 5.2. *Consider the family of games $G_m = (T, \vec{f}, h, m, n)$ and the strategies $\vec{S}(\vec{k})$, for $mhn < \sum_t f_t k_t hn$. For this family of game, $\lambda_{m, \vec{k}}$ is non-decreasing in m and non-increasing in \vec{k} ; $p_u^{m, \vec{k}}$ is non-decreasing in m and non-increasing in \vec{k} ; and the function BR_G is non-decreasing in \vec{k} and non-increasing in m .*

Monotonicity is enough to guarantee the existence of an equilibrium. Unfortunately, we know that a trivial equilibrium always exists in threshold strategies: all agents choose a threshold of \$0, so no agent ever volunteers. To guarantee the existence of a nontrivial equilibrium, it is sufficient to show there is some vector \vec{k} of thresholds such that $BR_G(\vec{k}) > \vec{k}$. The following lemma, whose proof is again deferred to Appendix B, shows that we can always find such a point for sufficiently large δ_t .

Lemma 5.3. *For all games $G = (T, \vec{f}, h, m, n)$, there exists a $\delta^* < 1$ such that if $\delta_t > \delta^*$ for all t , there is a vector \vec{k} of thresholds such that $BR_G(\vec{k}) > \vec{k}$.*

We are now ready to prove our main theorem: there exists a non-trivial equilibrium where all agents play threshold strategies greater than zero.

Theorem 5.2. *For all games $G = (T, \vec{f}, h, m, 1)$ and all ε , there exist n_ε^* and $\delta_{\varepsilon, n}^*$ such that, if $n > n_\varepsilon^*$ and $\delta_t > \delta_{\varepsilon, n}^*$ for all t , then there exists a nontrivial vector \vec{k} of thresholds that is an ε -Nash equilibrium. Moreover, there exists a greatest such vector.*

PROOF. By Lemma 5.2, BR_G is a non-decreasing function on a complete lattice, so Tarski's fixed point theorem [Tarski 1955] guarantees the existence of a greatest and least fixed point; these fixed points are equilibria. The least fixed point is the trivial equilibrium. We can compute the greatest fixed point by starting with the strategy profile (∞, \dots, ∞) (where each agent uses the strategy S_∞ of always

volunteering) and considering ϵ -best-reply dynamics, that is, iteratively computing the ϵ -best-reply strategy profile. Monotonicity guarantees this process converges to the greatest fixed point, which is an equilibrium (and is bound to be an equilibrium in pure strategies, since the best reply is always a pure strategy). Since there is a finite amount of money, this process needs to be repeated only a finite number of times. By Lemma 5.3, there exists a \vec{k} such that $BR_G(\vec{k}) > \vec{k}$. Monotonicity then guarantees that $BR_G(BR_G(\vec{k})) \geq BR_G(\vec{k})$ and similarly for any number of applications of BR_G . If \vec{k}^* is the greatest fixed point of BR_G , then $\vec{k}^* > \vec{k}$. Thus, the greatest fixed point is a nontrivial equilibrium. ■

The proof of Theorem 5.2 also provides an algorithm for finding equilibria that seems efficient in practice: start with the strategy profile (∞, \dots, ∞) and iterate the best-reply dynamics until an equilibrium is reached.




Fig. 1. A hypothetical best-reply function with one type of agent.

There is a subtlety in our results. In general, there may be many equilibria. From the perspective of social welfare, some will be better than others. As we show in Section 7, strategies that use smaller (but nonzero) thresholds increase social welfare. Consider the best-reply function shown in Figure 1. In the game G in the example, there is only one type of agent, so $BR_G : \mathbb{N} \rightarrow \mathbb{N}$. In equilibrium, we must have $BR(k) = k$; that is, an equilibrium is characterized by a point on the line $y = x$. This example has three equilibria, where all agents play s_0 , s_5 , and s_{10} respectively. The strategy profile where all agents play s_5 is the equilibrium that maximizes social welfare, while s_{10} is the greatest equilibrium.

In the rest of this paper, we focus on the greatest equilibrium in all our applications (although a number of our results hold for all nontrivial equilibria). This equilibrium has several desirable properties. First, it is guaranteed to be stable; best-reply dynamics from nearby points converge to it. By way of contrast, best-reply dynamics moves the system away from the equilibrium S_5 in Figure 1. Unstable equilibria are difficult to find in practice, and seem unlikely to be maintained for any length of time. Second, the “greatest” equilibrium is the one found by the natural algorithm given in Theorem 5.2. The proof of the theorem shows that it

is also the outcome that will occur if agents adopt the reasonable initial strategy of starting with a large threshold and then using best-reply dynamics. Finally, by focusing on the worst nontrivial equilibrium, our results provide guarantees on social welfare, in the same way that results on *price of anarchy* [Roughgarden and Tardos 2002] provide guarantees (since price of anarchy considers the social welfare of the Nash equilibrium with the worst social welfare).

6. SIMULATIONS

Theorem 4.1 proves that, for a sufficiently large number n of agents, and after a sufficiently large number r of rounds, the distribution of wealth will almost always be close to the distribution that minimizes relative entropy. In this section, we simulate the game to gain an understanding of how large n and r need to be in practice. The simulations show that our theoretical results apply even to relatively small systems; we get tight convergence with a few thousand agents, and weaker convergence for smaller numbers, in very few rounds, indeed, a constant number per agent.




Fig. 2. Maximum distance from minimum relative entropy distribution over 10^6 timesteps.

The first simulation explores the tightness of convergence to the distribution that minimizes relative entropy for various values of n . We used a single type of agent, with $\beta = \rho = \chi = 1$, $m = 2$, and $k = 5$. For each value of n , the simulation was started with a distribution of money as close as possible to the distribution d^* that minimizes relative entropy to the distribution q defined in Theorem 4.1 that characterizes the distribution of money in equilibrium (when the threshold strategy 5 is used). We then computed the maximum Euclidean distance between d^* and the observed distribution over 10^6 rounds. As Figure 2 shows, the system does not move far from d^* once it is there. For example, if $n = 5000$, the system is never more than distance .001 from d^* . If $n = 25,000$, it is never more than .0002 from d^* .

Figure 2 does show a larger distance for $n = 1000$, although in absolute terms it is still small. The next simulation shows that, while the system may occasionally move away from d^* , it quickly converges back to it. We averaged 10 runs of the




Fig. 3. Distance from minimum relative entropy distribution with 1000 agents.

Markov chain, starting from an extreme distribution (every agent has either \$0 or \$5), and considered the average time needed to come within various distances of d^* . As Figure 3 shows, after 2 rounds per agent, on average, the Euclidean distance from the average distribution of money to d^* is .008; after 3 rounds per agent, the distance is down to .001.




Fig. 4. Average time to get within .001 of the minimum relative entropy distribution.

Finally, we considered more carefully how quickly the system converges to d^* for various values of n . There are approximately k^n possible states, so the convergence time could in principle be quite large. However, we suspect that the Markov chain that arises here is *rapidly mixing*, which means that it will converge significantly faster (see [Lovasz and Winkler 1995] for more details about rapid mixing). We believe that the actually time needed is $O(n)$. This behavior is illustrated in Figure 4, which shows that for our example chain (again averaged over 10 runs), after approximately $3n$ steps, the Euclidean distance between the actual distribution of money in the system and d^* is less than .001. This suggests that we should expect the system to converge in a constant number of rounds per agent.

7. SOCIAL WELFARE AND SCALABILITY

In this section, we consider a fundamental question faced by system designers: what is the optimal amount of money and how does it depend on the size of the system? We discuss how our theoretical results from Sections 4 and 5 show that in order to maximize social welfare, the optimal amount of money is some constant per agent. Thus, a system designer that wants to maximize social welfare should manage the average quantity of money appropriately. However, we also show that this must be done carefully. Specifically, we show that increasing the amount of money improves performance up to a certain point, after which the system experiences a monetary crash. Once the system crashes, the only equilibrium will be the trivial one where all agents play s_0 . Thus, optimizing the performance of the system involves discovering how much money the system can handle before it crashes.

In Section 3, we define the game using a tuple $G = (T, \vec{f}, h, m, n)$. Thus, our definition of a game uses the average amount of money m rather than the equally reasonable total amount of money hmn . The choice is motivated by our theoretical results. Theorem 4.1 shows that the long-term distribution of money d^* depends on the average amount of money, but is independent n , provided it is sufficiently large. Thus, since we normalize δ_i by the number of agents in computing utility, the optimal threshold policy for the MDP developed in Appendix B is also independent of n . Theorems 5.1 and 5.2 show that such policies constitute an ε -Nash equilibrium. Thus, modulo a technical issue regarding the rate of convergence of the Markov Chain towards its stationary distribution, to determine the optimal amount of money for a large system, it suffices to determine the optimal value of m , the average amount of money per agent.

We remark that, in practice, it may be easier for the designer to vary the price of fulfilling a request than to control the amount of money in the system. This produces the same effect. For example, changing the cost of fulfilling a request from \$1 to \$2 is equivalent to halving the amount of money that each agent has. Similarly, halving the the cost of fulfilling a request is equivalent to doubling the amount of money that everyone has. With a fixed amount hmn of money, there is an optimal product hnc of the number hn of agents and the cost c of fulfilling a request.

This also tells us how to deal with a dynamic pool of agents. Our system can handle newcomers relatively easily: simply allow them to join with no money. This gives existing agents no incentive to leave and rejoin as newcomers. (By way of contrast, in systems where each new agent starts off with a small amount of money, such an incentive clearly exists.) We then change the price of fulfilling a request so that the optimal ratio is maintained. This method has the nice feature that it can be implemented in a distributed fashion; if all nodes in the system have a good estimate of n , then they can all adjust prices automatically. (Alternatively, the number of agents in the system can be posted in a public place.) Approaches that rely on adjusting the amount of money may require expensive system-wide computations (see [Vishnumurthy et al. 2003] for an example), and must be carefully tuned to avoid creating incentives for agents to manipulate the system by which this is done.

Note that, in principle, the realization that the cost of fulfilling a request can change can affect an agent's strategy. For example, if an agent expects the cost

to increase, then he may want to defer volunteering to fulfill a request. However, if the number of agents in the system is always increasing, then the cost always decreases, so there is never any advantage in waiting. There may be an advantage in delaying a request, but it is far more costly, in terms of waiting costs than in providing service, since we assume the need for a service is often subject to real waiting costs. In particular, many service requests, such as those for information or computation, cannot be delayed without losing most of their value.

Issues of implementation aside, we have now reduced the problem of determining the optimal total amount of money for a large system to that of determining the optimal average amount of money, independent of the exact number of agents. Before we can determine the optimal value of m , we have to answer a more fundamental question: given an equilibrium that arose for some value of m , how good is it?

Consider a single round of the game with a population of a single type t and an equilibrium threshold k . If a request is satisfied, social welfare increases by $\gamma_t - \alpha_t$; the requester gains γ_t utility and the satisfier pays a cost of α_t . If no request is satisfied then no utility is gained. What is the probability that a request will be satisfied? This requires two events to occur. First, the agent chosen to make a request must have a dollar, which happens with probability approximately $1 - \zeta$, where $\zeta = d^*(t, 0)$ is the fraction of agents with no money. Second, there must be a volunteer able and willing to satisfy the request. Any agent who does not have his threshold amount of money is willing to volunteer. Thus, if $\theta = d^*(t, k_t)$ is the fraction of agents at their threshold, then the probability of having a volunteer is $1 - (1 - \beta_t)^{(1-\theta)n}$. We believe that in most large systems this probability is quite close to 1; otherwise, either β_t must be unrealistically small or θ must be very close to 1. For example, even if $\beta = .01$ (i.e., an agent can satisfy 1% of requests), 1000 agents will be able to satisfy 99.99% of requests. If θ is close to 1, then agents will have an easier time earning money than spending money (the probability of spending a dollar is at most $1/n$, while for large β the probability of earning a dollar if an agent volunteers is roughly $(1/n)(1/(1 - \theta))$). If an agent is playing s_4 and there are n rounds played a day, this means that for $\theta = .9$ he would be willing to pay α_t today to receive γ_t over 10 years from now. For most systems, it seems unreasonable to have δ_t or γ_t/α_t this large. Thus, for the purposes of our analysis, we approximate $1 - (1 - \beta_t)^{(1-\theta)n}$ by 1.

With this approximation, we can write the expected increase in social welfare each round as $(1 - \zeta)(\gamma_t - \alpha_t)$. Since $U_i(\vec{S})$ is normalized by the discount factor, the total expected social welfare summed over all rounds is also $(1 - \zeta)(\gamma_t - \alpha_t)$. If we have more than one type of agent, the situation is essentially the same. The equation for social welfare is more complicated because now the gain in welfare depends on the γ , α , and δ of the agents chosen in that round, but the overall analysis is the same, albeit with more cases. In the general case,

$$\zeta = \sum_t d^*(t, 0) \quad (5)$$

Thus our goal is clear: find the amount of money that, in equilibrium, minimizes ζ .

In general, as the following theorem shows, ζ decreases as m increases. More specifically, given our assumption that the system is starting at the greatest equi-

librium \vec{k} , increasing m and then following best response dynamics leads to the new greatest equilibrium \vec{k}' . As long as \vec{k}' is non-trivial, $\zeta_{m',\vec{k}'} \leq \zeta_{m,\vec{k}}$.

Theorems 4.1, 5.1, and 5.2 place requirements on the values of n and δ_t . Intuitively, the theorems require that the δ_t s is sufficiently large to ensure that agents are patient enough that their decisions are dominated by long-run behavior rather than short-term utility, and that n is sufficiently large to ensure that small changes in the distribution of money do not move it far from d^* . In the theorems in this section, assume that these conditions are satisfied. To simplify the statements of the theorems, we use “the standard conditions hold” to mean that the game $G = (T, \vec{f}, h, m, n)$ under consideration is such that $n > n^*$ and $\delta_t > \delta^*$ for the n^* and δ^* needed for the results of Theorems 4.1, 5.1, and 5.2 to apply.

Theorem 7.1. *Let $G = (T, \vec{f}, h, m, n)$ be such that the standard conditions hold, and let \vec{k} be the greatest equilibrium for G . Then if $m' > m$, the best-reply dynamics in $G' = (T, \vec{f}, h, m', n)$ starting at \vec{k} converge to some $\vec{k}' \leq \vec{k}$ that is the greatest equilibrium of G' . If \vec{k}' is a nontrivial equilibrium, then $\zeta_{m',\vec{k}'} \leq \zeta_{m,\vec{k}}$.*

PROOF. Since \vec{k} is an equilibrium, $BR_G(\vec{k}) = \vec{k}$. By Lemma 5.2, BR_G is non-increasing in m . Thus, $\vec{k} = BR_G(\vec{k}) \geq BR_{G'}(\vec{k})$. Applying best-reply dynamics to $BR_{G'}$ starting at \vec{k} as in Theorem 5.2 gives us an equilibrium \vec{k}' such that $\vec{k}' \leq \vec{k}$. By Lemma 5.2, $BR_G(\vec{k}'')$ is non-decreasing in \vec{k}'' , so this is the greatest equilibrium. Suppose \vec{k}' is nontrivial. By Equations (2) and (5),

$$\zeta_{m,\vec{k}} = \sum_t d^*(t, 0) = \sum_t \frac{f_t q(t, 0)}{\sum_{j=0}^{k_t} \lambda_{m,\vec{k}}^j q(t, j)}.$$

Again by Lemma 5.2, $\lambda_{m,\vec{k}}$ is non-decreasing in m and non-increasing in \vec{k} . Thus, $\zeta_{m',\vec{k}'} \leq \zeta_{m,\vec{k}}$. ■

Theorem 7.1 tells us that, as long as the system does not crash, more money is better. The following corollary tells us that such a crash is an essential feature; a sufficient increase in the amount of money leads to a monetary crash. Moreover, once the system has crashed, adding more money does not cause the system to become “uncrashed.”

Corollary 7.1. *Consider the family of games $G_m = (T, \vec{f}, h, m, n)$ such that the standard conditions hold. There exists a critical average amount m^* of money such that if $m < m^*$, then G_m has a nontrivial equilibrium, while if $m > m^*$, then G_m has no nontrivial equilibrium. (A nontrivial equilibrium may or may not exist if $m = m^*$.)*

PROOF. To see that there is some m for which G_m has no nontrivial equilibrium, fix m . If there is no nontrivial equilibrium in G_m , we are done. Otherwise, suppose that the greatest equilibrium in G_m is \vec{k} . Choose $m' > \sum_t f_t k_t$, and let \vec{k}' be the greatest equilibrium in $G_{m'}$. By Theorem 7.1, $\vec{k}' \leq \vec{k}$. But if \vec{k}' is a nontrivial equilibrium then, in equilibrium, each agent of type t has at most $k'_t \leq k_t$ dollars. But then $m' > \sum_t f_t k_t \geq \sum f_t k'_t$, a contradiction.

Let m^* be the infimum over all m for which no nontrivial equilibrium exists in the game G_m . Clearly, by choice of m^* , if $m < m^*$, there is a nontrivial equilibrium

in G_m . Now suppose that $m > m^*$. By the construction of m^* , there exists m' with $m > m' \geq m^*$ such that no nontrivial equilibrium exists in $G_{m'}$. Let the greatest equilibria with m' and m be \vec{k}' and \vec{k} , respectively. By Theorem 7.1, $\vec{k} \leq \vec{k}'$. Thus \vec{k} is also trivial. ■

Figure 5 shows an example of the monetary crash in the game

$$(\{(.05, 1, 1, .95, 1), (.15, 1, 1, .95, 1)\}, (.3, .7), 10, m, 100).$$




Fig. 5. Social welfare for various average amounts of money, demonstrating a monetary crash.

In light of Corollary 7.1, the system designer should try to find m^* , the point where social welfare is maximized. We discuss how she might go about finding m^* in practice in Section 9. The system designer may wish to choose the m somewhat less than m^* . Since there will be a crash for any $m > m^*$, small changes in the characteristics of the population or mistakes by the designer in modeling them could lead to a crash if she chooses m too close to m^* .

The phenomenon of a monetary crash is intimately tied to our assumption of fixed prices. We saw such a crash in practice in the babysitting co-op example. If the price is allowed to float freely, we expect that, as the money supply increases, there will be inflation; the price will increase so as to avoid a crash. However, floating prices can create other monetary problems, such as speculation, booms, and busts. Floating prices also impose transaction costs on agents. In systems where prices would normally be relatively stable, these transaction costs may well outweigh the benefits of floating prices, so a system designer may opt for fixed prices, despite the risk of a crash.

8. DEALING WITH NONSTANDARD AGENTS

The model in Section 3 defines the utility of standard agents, who value service and dislike using their resources to provide it to others. This seems like a natural description of the way most people use distributed systems. However, in a real system, not every user will behave the way the designer intends. A practical system needs to be robust to nonstandard behaviors. In this section, we show

how our model can be used to understand the effects of four interesting types of nonstandard behavior. First, an agent might provide service even when he will receive nothing in return, behaving as an altruist. Second, rather than viewing money as a means to satisfy future requests, an agent might place an inherent value on it and start hoarding it. Third, an agent might create additional identities, known as *sybils*, to try and manipulate the system. Finally, agents might collude with each other.

The results of this section give a system designer insight into how to design a scrip system that takes into account (and is robust to) a number of frequently-observed behaviors.

8.1 Altruists

P2P filesharing systems often have large numbers of free riders; they work because a small number of altruistic users provide most of the files. For example, Adar and Huberman [2000] found that, in the Gnutella network, nearly 50 percent of responses are from the top 1 percent of sharing hosts. A wide variety of systems have been proposed to discourage free riding (see Section 2). However, in our model, unless this system mostly eliminates the altruistic users, adding such a system will have no effect on rational users.

To make this precise, take an altruist to be someone who always volunteers to fulfill requests, regardless of whether the other agent can pay. Agent i might rationally behave altruistically if, rather than suffering a loss of utility when satisfying a request, i derives positive utility from satisfying it. Such a utility function is a reasonable representation of the pleasure that some people get from the sense that they provide the music that everyone is playing. For such altruistic agents, the strategy of always volunteering is dominant. While having a nonstandard utility function might be one reason that a rational agent might use this strategy, there are certainly others. For example a naive user of filesharing software with a good connection might well follow this strategy. All that matters for the following discussion is that there are some agents that use this strategy, for whatever reason. For simplicity, we assume that all such agents have the same type t_a .

Suppose that a system has a altruists. Intuitively, if a is moderately large, they will manage to satisfy most of the requests in the system even if other agents do no work. Thus, there is little incentive for any other agent to volunteer, because he is already getting full advantage of participating in the system. Based on this intuition, it is a relatively straightforward calculation to determine a value of a that depends only on the types, but not the number n , of agents in the system, such that the dominant strategy for all standard agents i is to never volunteer to satisfy any requests.

Proposition 8.1. *For all games $(T, \vec{f}, h, m, 1)$ with $f_{t_a} > 0$, there exists a value a such that, if $n > a/(f_{t_a}h)$, then never volunteering is a dominant strategy for all standard agents.*

PROOF. Consider the strategy for a standard agent i in the presence of a altruists. Even with no money, agent i will get a request satisfied with probability $1 - (1 - \beta_{t_a})^a$ just through the actions of the altruists. Consider a round when agent i is chosen to make a request. If he has no money (because he never volunteered), his expected

utility is $\gamma_{\tau(i)}(1 - (1 - \beta_{t_a})^a)$. His maximum possible utility for the round is $\gamma_{\tau(i)}$. Thus, a strategy where he volunteers can increase his utility for a round by at most $\gamma_{\tau(i)}(1 - \beta_{t_a})^a$. Thus, even if the agent gets every request satisfied, his expected utility can increase by at most

$$\begin{aligned} & (1 - \delta_{\tau(i)}) \sum_{r=1}^{\infty} (\rho_{\tau(i)}/hn) \gamma_{\tau(i)} (1 - \beta_{t_a})^a (1 - (1 - \delta_{\tau(i)})/n)^r \\ &= (1 - \delta_{\tau(i)}) (\rho_{\tau(i)}/h) \gamma_{\tau(i)} (1 - \beta_{t_a})^a (1 - \delta_{\tau(i)}) \\ &= (\rho_{\tau(i)}/h) \gamma_{\tau(i)} (1 - \beta_{t_a})^a. \end{aligned}$$

Clearly this expression goes to 0 as a goes to infinity. If we take a large enough that the expression is less than α_t for all types t , then the value of having every future request satisfied is less than the cost of volunteering now, so no agent will ever volunteer. ■

Consider the following reasonable values for our parameters: $\beta_t = .01$ (so that each player can satisfy 1% of the requests), $\gamma_t = 1$, $\alpha_t = .1$ (a low but non-negligible cost), $\delta_t = .9999/\text{day}$ (which corresponds to a yearly discount factor of approximately 0.95), and an average of 1 request per day per player. Then as long as $a > 1145$ to ensure that not volunteering is a dominant strategy. While this is a large number, it is small relative to the size of a large P2P network.

Proposition 8.1 shows that with enough altruists, the system eventually experiences a monetary crash, since all agents will use a threshold of zero. However, interesting behavior can still arise with smaller numbers of altruists. Consider the situation where an a fraction of requests are immediately satisfied at no cost without the requester needing to ask for volunteers. Intuitively, these are the requests satisfied by the altruists, although the following result also applies to any setting where agents occasionally have a (free) outside option. The following theorem shows that social welfare is increasing in a .

Let $G = (\{t\}, 1, h, m, n)$ be a game with a single type for which the standard conditions hold. Consider the family G_a of games (parameterized by a) that result from G if a fraction a of requests can be satisfied at no cost. That is, the game G_a is the same as G , except that if an agent i makes a request, with probability a , it is satisfied at no cost, and with probability $1 - a$, an agent is chosen among the volunteers to satisfy the request, just as in G , and the i is charged 1 dollar to have the request satisfied.

Theorem 8.1. *For the interval of values of a where there is no monetary crash in G_a , social welfare increases as a increases (assuming that the greatest equilibrium is played by all agents in G_a).*

PROOF. An agent's utility in a round where he makes a request and it is satisfied at no cost is γ_t . Since such rounds occur with probability a , by assumption, our normalization guarantees that the sum of an agent's expected utility in rounds where a request is satisfied at no cost is $a\gamma_t$. The same analysis as in Section 7 shows that the some of an agent's expected utility in the remaining rounds is $(1 - a)(1 - \zeta(a))(\gamma_t - \alpha_t)$, where, as before, $\zeta(a) = d^*(t, 0, a)$, the equilibrium value of $d^*(t, 0)$ in the game G_a . Thus, expected utility as a function of a is

$$a\gamma_t + (1 - a)(1 - \zeta(a))(\gamma_t - \alpha_t). \quad (6)$$

To see that this expression increases as a increases, we would like to take the derivative relative to a and show it is positive. Unfortunately, $\zeta(a)$ may not even be continuous. Because strategies are integers, there will be regions where $\zeta(a)$ is constant, and then a jump when a critical value of a is reached that causes the equilibrium to change. At a point a in a region where $\zeta(a)$ is constant, $\zeta'(a) = 0$, so the derivative of Equation (6) is $\gamma_t - (1 - \zeta(a))(\gamma_t - \alpha_t) > 0$. Hence, social welfare is increasing at such points.

Now consider a point a where $\zeta(a)$ is discontinuous. Such a discontinuity occurs when the greatest equilibrium, the greatest value \vec{k} for which $BR_{G_a}(\vec{k}) = \vec{k}$, changes. We show that, for a fixed \vec{k} , $BR_{G_a}(\vec{k})$ is non-increasing in a . Since increasing a can only cause the $BR_{G_a}(\vec{k})$ to decrease, the discontinuity must be caused by a change from an equilibrium \vec{k} to a new equilibrium $\vec{k}' < \vec{k}$. Fix a vector \vec{k} of thresholds, and let $p_u^{\vec{k},m,a}$ be the probability that i will be earn a dollar in a given round if he is willing to volunteer, given that a fraction a of requests is satisfied at no cost (so that $p_u^{\vec{k},m,0}$ is what we earlier called $p_u^{\vec{k},m}$); we similarly define $p_d^{\vec{k},m,a}$. It is easy to see that $p_u^{\vec{k},m,a} = (1 - a)p_u^{\vec{k},m,0}$ and $p_d^{\vec{k},m,a} = (1 - a)p_d^{\vec{k},m,0}$. The random variable $J(\kappa, p_u, p_d)$ in Equation (4) describes the first time at which an agent starting with κ dollars and using the threshold κ while earning a dollar with probability p_u and spending a dollar with probability p_d reaches zero dollars. As a increases, $p_u^{\vec{k},m,a}$ and $p_d^{\vec{k},m,a}$ both decrease, but the ratio $p_u^{\vec{k},m,a}/p_d^{\vec{k},m,a}$ remains constant. Intuitively, this means that the agent “slows down” his random walk on amounts of money by a factor of $1/(1 - a)$. Thus, the value of the expectation in Equation (4), and hence the right-hand side of Equation (4), is decreasing as a function of a . By Lemma 5.1, $(BR_{G_a}(\vec{k}))_t$ is the maximum value of κ such that Equation (4) is satisfied. Decreasing the right-hand side can only decrease the maximum value of κ , so $BR_{G_a}(\vec{k})$ is non-increasing as a function of a .

By Lemma 5.2, $\lambda_{m,\vec{k}}$ is non-increasing in \vec{k} (unless the system crashes). Since, as we have just shown, if there is a discontinuity at $\zeta(a)$ when a increases, the greatest equilibrium changes at a from \vec{k} to $\vec{k}' < \vec{k}$, we must have $\lambda_{m,\vec{k}'} \geq \lambda_{m,\vec{k}}$. In Equation (2) for $i = 0$, the value of the numerator is independent of λ , but the denominator with $\lambda_{m,\vec{k}'}$ is greater than or equal to the denominator with $\lambda_{m,\vec{k}}$. Thus $d^*(t, 0, a) = \zeta(a)$ is non-increasing at a . By Equation (6), this means that expected utility is increasing at a . Thus, in either case, social welfare is increasing in a . ■

Theorem 8.1 and Proposition 8.1 combine to tell us that a little altruism is good for the system, but too much causes a crash. Figure 6 demonstrates this phenomenon. As we saw in Section 7, such crashes are caused when m , the average amount of money, is too large. By decreasing m appropriately, even relatively large values of a can be exploited, as Figure 7 shows. The “social welfare without adjustment” plot is the same data from Figure 6, with the corresponding plot of the amount of money horizontal since m was held fixed. By decreasing the average amount of money appropriately as the number of altruists increases, a system designer can increase social welfare while avoiding a crash.





Fig. 6. Altruists can cause a crash.


 Fig. 7. m can be adjusted as a increases.

8.2 Hoarders

Whenever a system allows agents to accumulate something, be it work done, as in SETI@home, friends on online social networking sites, or “gold” in an online game, a certain group of users seems to make it their goal to accumulate as much of it as possible. In pursuit of this, they will engage in behavior that seems irrational. For simplicity here, we model hoarders as playing the strategy s_∞ . This means that they will volunteer under all circumstances. Our analysis would not change significantly if we also required that they never made a request for work. Our first result shows that, for a fixed money supply, having more hoarders makes standard agents worse off.

Consider a game $G = (T, \vec{f}, h, m, n)$ such that the standard conditions hold. Consider the family G_{f_h} of games (parameterized by f_h) that result from G if a fraction f_h of agents are hoarders. That is, $G_{f_h} = (T \times \{0, 1\}, \vec{f}', h', m, n)$ where at agent of type $(t, 0)$ is a standard agent of type t , but an agent of type $(t, 1)$ is

a hoarder and always uses the strategy s_∞ (his probabilities are still determined by β_t , ρ_t , and χ_t). Define \vec{f}' by taking $f'_{(t,0)} = (1 - f_h)f_t$ and $f'_{(t,1)} = f_h f_t$ for all types t . Let h' be the smallest multiple of h such that $f_{(t,i)}h'$ is an integer for all t and i . (We need to adjust h because otherwise the number of agents in the base game may not be well defined.) Finally, to account for the changed h , let $\delta_{(t,i)} = 1 - (1 - \delta_t)h/h'$.

Theorem 8.2. *In the family G_{f_h} of games, social welfare is non-increasing in f_h (if the greatest equilibrium is played by all agents in G_{f_h}).*

PROOF. Let $\vec{k}(f_h)$ denote the greatest equilibrium in G_{f_h} . An increase in f_h is equivalent to taking some number of standard agents and increasing their strategy to s_∞ . It follows from Lemma 5.2 that $BR_{G_{f_h}}$ is non-decreasing in f_h , and so $\vec{k}(f_h)$ is non-decreasing in f_h . Again by Lemma 5.2, $\lambda_{m,\vec{k}(f_h)}$ is non-increasing in f_h . Let $\zeta f_h = 1/(1 - f_h) \sum_t d^*((t,0), 0, f_h)$ be the fraction of non-hoarders with zero dollars, where $d^*((t,0), 0, f_h)$ is the value of $d^*((t,0), 0)$ at the greatest equilibrium of G_{f_h} . By Equation (2), $\zeta(f_h)$ is non-decreasing in f_h . Thus, social welfare is non-increasing in f_h . ■

Hoarders do have a beneficial aspect. As we have observed, a monetary crash occurs when a dollar becomes valueless, because there are no agents willing to take it. However, with hoarders in the system, there is always someone who will volunteer, so there cannot be a crash. Thus, for any m , the greatest equilibrium will be nontrivial and, by Theorem 7.1, social welfare keeps increasing as m increases. So, in contrast to altruism, where the appropriate response was to decrease m , the appropriate response to hoarders is to increase m . In fact, our results indicate that the optimal response to hoarders is to make m infinite. This is due to our unrealistic assumption that hoarders would use the strategy s_∞ regardless of the value of m . There is likely an upper limit on the value of m in practice, since it is unlikely that hoarders would be willing to hoard scrip if it is so easily available.

8.3 Sybils

Unless identities in a system are tied to a real world identity (for example by a credit card), it is effectively impossible to prevent a single agent from having multiple identities [Douceur 2002]. Nevertheless, there are a number of techniques that can make it relatively costly for an agent to do so. For example, Credence uses cryptographic puzzles to impose a cost each time a new identity wishes to join the system [Walsh and Sier 2006]. Given that a designer can impose moderate costs to sybiling, how much more need she worry about the problem? In this section, we show that the gains from creating sybils when others do not diminish rapidly, so modest costs may well be sufficient to deter sybiling by typical users. However, sybiling is a self-reinforcing phenomenon. As the number of agents with sybils gets larger, the cost to being a non-sybiling agent increases, so the incentive to create sybils becomes stronger. Therefore, measures to discourage or prevent sybiling should be taken early before this reinforcing trend can start. Finally, we examine the behavior of systems where only a small fraction of agents have sybils. We show that under these circumstances a wide variety of outcomes are possible (even when all agents are of a single type), ranging from a crash (where no service is provided)




Fig. 8. The effect of p_u on utility

to an increase in social welfare. This analysis provides insight into the tradeoffs between efficiency and stability that occur when controlling the money supply of the system's economy.

When an agent of type t creates sybils, the only parameter of his type that may change as a result is χ_t , if we redefine the likelihood of an agent being chosen to be the likelihood of the agent or any of his sybils being chosen. For simplicity, we assume that each sybil is as likely to be chosen as the original agent, so creating s sybils increases χ_t by $s\chi_t$. (Sybils may have other impacts on the system, such as increased search costs, but we expect these to be minor.)

Increasing χ_t benefits an agent by increasing his value of ω_t and thus p_u , his probability of earning a dollar (see Equation (9) in Appendix B). When $p_u < p_d$, the agent has more opportunities to spend money than to earn money, so he will regularly have requests go unsatisfied due to a lack of money. In this case, the fraction of requests he has satisfied is roughly p_u/p_d , so increasing p_u by creating sybils results in a roughly linear increase in utility. As Theorem 8.3 shows, when p_u is close to p_d , the increase in satisfied requests is no longer linear, so the benefit of increasing p_u begins to diminish. Finally, when $p_u > p_d$, most of the agent's requests are being satisfied, so the benefit from increasing p_u is very small. Figure 8 illustrates an agent's utility as p_u varies for $p_d = .0001$.³ We formalize the relationship between p_u , p_d , and the agent's utility in the following theorem, whose proof is deferred to Appendix C.

Theorem 8.3. *Fix a game G and vector of thresholds \vec{k} . Let $R_{\vec{k},t}^t = p_u^{\vec{k},t}/p_d^t$. In the limit as the number of rounds goes to infinity, the fraction of the agent's requests that have an agent willing and able to satisfy them that get satisfied is $(R_{\vec{k},t}^t - R_{\vec{k},t}^{k_t+1})/(1 - R_{\vec{k},t}^{k_t+1})$ if $R_{\vec{k},t}^t \neq 1$ and $k_t/(k_t + 1)$ if $R_{\vec{k},t}^t = 1$.*

³Except where otherwise noted, the remaining figures in this section assume that $m = 4$, $n = 10000$ and that there is a single type of rational agent with $\alpha = .08$, $\beta = .01$, $\gamma = 1$, $\delta = .97$, $\rho = 1$, and $\chi = 1$. These values are chosen solely for illustration, and are representative of a broad range of parameter values. The figures are based on calculations of the equilibrium behavior.




Fig. 9. The effect of sybils on utility

Theorem 8.3 gives insight into the equilibrium behavior with sybils. Clearly, if sybils have no cost, then creating as many as possible is a dominant strategy. However, in practice, we expect there is some modest overhead involved in creating and maintaining a sybil, and that a designer can take steps to increase this cost without unduly burdening agents. With such a cost, adding a sybil might be valuable if p_u is much less than p_d , and a net loss otherwise. This makes sybils a self-reinforcing phenomenon. When a large number of agents create sybils, agents with no sybils have their p_u significantly decreased. This makes them much worse off and makes sybils much more attractive to them. Figure 9 shows an example of this effect. This self-reinforcing quality means that it is important to take steps to discourage the use of sybils before they become a problem. Luckily, Theorem 8.3 also suggests that a modest cost to create sybils will often be enough to prevent agents from creating them because with a well chosen value of m , few agents should have low values of p_u .

We have interpreted Figures 8 and 9 as being about changes in χ due to sybils, but the results hold regardless of what caused differences in χ . For example, agents may choose a volunteer based on characteristics such as connection speed or latency. If these characteristics are difficult to verify and do impact decisions, our results show that agents have a strong incentive to lie about them. This also suggests that the decision about what sort of information the system should enable agents to share involves tradeoffs. If advertising legitimately allows agents to find better service or more services they may be interested in, then advertising can increase social welfare. But if these characteristics impact decisions but have little impact on the actual service, then allowing agents to advertise them can lead to a situation like that in Figure 9, where some agents have a significantly worse experience.

We have seen that when a large fraction of agents have sybils, those agents without sybils tend to be starved of opportunities to work (i.e. they have a low value of p_u). However, as Figure 9 shows, when a small fraction of agents have sybils this effect (and its corresponding cost) is small. Surprisingly, if there are few agents with sybils, an increase in the number of sybils these agents have can actually




Fig. 10. Sybils can improve utility




Fig. 11. Sybils can cause a crash

result in a *decrease* of their effect on the other agents. Because agents with sybils are more likely to be chosen to satisfy any particular request, they are able to use lower thresholds and reach those thresholds faster than they would without sybils, so fewer are competing to satisfy any given request. Furthermore, since agents with sybils can almost always pay to make a request, they can provide more opportunities for other agents to satisfy requests and earn money. Social welfare is essentially proportional to the number of satisfied requests (and is exactly proportional to it if everyone shares the same values of α and γ), so a small number of agents with a large number of sybils can improve social welfare, as Figure 10 shows. Note that, although social welfare increases, some agents may be worse off. For example, for the choice of parameters in this example, social welfare increases when twenty percent of agents create at least two sybils, but agents without sybils are worse off unless the twenty percent of agents with sybils create at least eight sybils. As the number of agents with sybils increases, they start competing with each other

for opportunities to earn money and so adopt higher thresholds, and this benefit disappears. This is what causes the discontinuity in Figure 9 when approximately a third of the agents have sybils.

This observation about the discontinuity also suggests another way to mitigate the negative effects of sybils: increase the amount of money in the system. This effect can be seen in Figure 11, where for $m = 2$ social welfare is very low with sybils but by $m = 4$ it is higher than it would be without sybils.

Unfortunately, increasing the average amount of money has its own problems. Recall from Section 7 that, if the average amount of money per agent is too high, the system will crash. It turns out that just a small number of agents creating sybils can have the same effect, as Figure 11 shows. With no sybils, the point at which social welfare stops increasing and the system crashes is between $m = 10.25$ and $m = 10.5$. If one-fifth of the agents each create a single sybil, the system crashes if $m = 9.5$, a point where, without sybils, the social welfare was near optimal. Thus, if the system designer tries to induce optimal behavior without taking sybils into account, the system will crash. Moreover, because of the possibility of a crash, raising m to tolerate more sybils is effective only if m was already set conservatively.

This discussion shows that the presence of sybils can have a significant impact on the tradeoff between efficiency and stability. Setting the money supply high can increase social welfare, but at the price of making the system less stable. Moreover, as the following theorem shows, whatever efficiencies can be achieved with sybils can be achieved without them, at least if there is only one type of agent. In the theorem, we consider a system where all agents have the same type t . Suppose that some subset of the agents have created sybils, and all the agents in the subset have created the same number of sybils. We can model this by simply taking the agents in the subsets to have a new type s , which is identical to t except that the value of χ increases. Thus, we state our results in terms of systems with two types of agents, t and s .

Theorem 8.4. *Suppose that t and s are two types that agree except for the value of χ , and that $\chi_t < \chi_s$. If $\vec{k} = (k_t, k_s)$ is an ε -Nash equilibrium for $G = (\{t, s\}, \vec{f}, h, m, n)$ with social welfare w , then there exist h' , m' , and n' such that $\vec{k}' = (k_s)$ is an ε -Nash equilibrium for $G'_{h', m', n'} = (\{t\}, \{1\}, h', m', n')$ with social welfare greater than w .*

We defer proof of Theorem 8.4 to Appendix C.

The analogous result for systems with more than one type of agent is not true. Figure 9 shows a game with a single type of agent, some of whom have created two sybils. However, we can reinterpret it as a game with two types of agents, one of whom has a larger value of χ . With this reinterpretation, Figure 9 shows that social welfare is higher when all the agents are of the type t_h with the higher value of χ than when only 40% are. Moreover, if only 40% of the agents have type t_h , social welfare would increase if the remaining agents created two sybils each (resulting in all agents having the higher value of χ). Note that this situation, where there are two types of agents, of which one has a higher value of χ , is exactly the situation considered by Theorem 8.4. Thus, the theorem shows that for any equilibrium with two such types of agents, there is a better equilibrium where one of those types creates sybils so as to effectively create only one type of agent.

While situations like this show that it is theoretically possible for sybils to increase

social welfare beyond what is possible to achieve by simply adjusting the average amount of money, this outcome seems unlikely in practice. It relies on agents creating just the right number of sybils. For situations where such a precise use of sybils would lead to a significant increase in social welfare, a designer could instead improve social welfare by biasing the algorithm agents use for selecting which volunteer will satisfy the request.

Thus far, we have assumed that when agents create sybils the amount of money in the system does not change. However, the presence of sybils increases the number of apparent agents in the system. Since social welfare depends on the average amount of money per agent, if the system designer mistakes these sybils for an influx of new users and increases the money supply accordingly, she will actually end up increasing the average amount of money in the system, and may cause a crash. This emphasizes the need for continual monitoring of the system rather than just using simple heuristics to set the average amount of money, an issue we discuss more in Section 9.

8.4 Collusion

Agents that collude gain two major benefits. The primary benefit is that they can share money, which makes them less likely to run out of money (and hence unable to make a request), and allows them to pursue a joint strategy for determining when to work. A secondary benefit, but important in particular for larger collusive groups, is that they can satisfy each other's requests. The effects of collusion on the rest of the system depend crucially on whether agents are able to volunteer to satisfy requests when they personally cannot satisfy the request but one of their colluding partners can. In a system where a request is for computation, it seems relatively straightforward for an agent to pass the computation to a partner to perform and then pass the answer back to the requester. On the other hand, if a request is a piece of a file it seems less plausible that an agent would accept a download from an unexpected source, and it seems wasteful to have the chosen volunteer download it for the sole purpose of immediately uploading it. If it is possible for colluders to pass off requests in this fashion, they are able to effectively act as sybils for each other, with all the consequences discussed in Section 8.3. However, if agents can volunteer only for requests they can personally satisfy, the effects of collusion are almost entirely positive.

Since we have already discussed the consequences of sybils, we will assume that agents are able to volunteer only to satisfy requests that they personally can satisfy. Furthermore, we make the simplifying assumption that agents that collude are of the same type, because if agents of different types collude their strategic decisions become more complicated. For example, once the colluding group has accumulated a certain amount of money, it may wish to have only members with small values of α volunteer to satisfy requests; or when it is low on money, it may wish to deny use of money to members with low values of γ . This results in strategies that involve sets of thresholds rather than a single threshold. While there seems to be nothing fundamentally different about the situation, it makes calculations significantly more difficult.

With these assumptions, we now examine how colluding agents will behave. Because colluding agents share money and types, it is irrelevant which members actu-

ally perform work and have money. All that matters is the total amount of money the group has. This means that when the group needs money, everyone in the group volunteers for a job; otherwise, no one does. Thus, the group essentially acts like a single agent, using a threshold that is somewhat less than the sum of the thresholds that the individual agents would have used, because it is less likely that c agents will make ck requests in rapid succession than a single agent making k . Furthermore, some requests will not require scrip at all because they can potentially be satisfied by other members of the colluding group. When deciding whether the group should satisfy a member's request or ask for an outside volunteer to fulfill it, the group must decide whether it should pay a cost of α to avoid spending a dollar. Since not spending a dollar is effectively the same as earning a dollar, the decision is already optimized by the threshold strategy; the group should always attempt to satisfy a request internally unless it is in a temporary situation where the group is above its threshold.




Fig. 12. The effect of collusion on utility

Figure 12 shows an example of the effects of collusion on agents' utilities as the size of collusive groups increases. As this figure suggests, the effects typically go through three phases. Initially, the fraction of requests colluders satisfy for each other is small. This means that each collusive group must work for others to pay for almost every request its members make. However, since they share money, the colluders do not have to work as often as individuals would. Thus, other agents have more opportunity to work, and every agent's p_u increases, making all agents better off.

As the number of colluders increases, the fraction of requests they satisfy internally grows significant. We can think of p_d as decreasing in this case, and view these requests as being satisfied "outside" the scrip system because no scrip changes hands. This is good for colluders, but is bad for other agents whose p_u is lower, since fewer requests are being made. Even in this range, non-colluding agents still tend to be better off than if there were no colluders, because the overall competition for opportunities to work is still lower. Finally, once the collusive group is large

enough, it will have a low p_d relative to p_u . This means the collusive group can use a very low threshold, which again begins improving utility for all agents. The analogous situation with sybils is transitory, and disappears when more agents create sybils. However, with collusion, this low threshold is an inherent consequence of colluders satisfying each other's requests, and so persists and even increases as the amount of collusion in the system increases. Since collusion is difficult to maintain (the problem of incentivizing agents to contribute is the whole point of using scrip), we would expect the size of collusive groups seen in practice to be relatively small. Therefore, we expect that for most systems collusion will make no agent worse off, and some better off. Note that, as with sybils, the decreased in competition that results from collusion can also lead to a crash. However, if the system designer is monitoring the system, and encouraging and expecting collusion, she can reduce m appropriately and prevent a crash.

These results also suggest that creating the ability to take out loans (with an appropriate interest rate) is likely to be beneficial. Loans gain the benefits of reduced competition without the accompanying cost of fewer requests being made in the system. However, implementing a loan mechanism requires addressing a number of other incentive problems. For example, *whitewashing*, where agents take on a new identity (in this case to escape debts) needs to be prevented [Friedman and Resnick 2001].

9. IDENTIFYING USER STRATEGIES

In Section 4, we used relative entropy to derive an explicit formula for the distribution of money d^* given a game (T, \vec{f}, h, n, m) and vector of strategies \vec{k} . In this section, we want to go in the opposite direction: given the distribution of money, we want to infer the strategies \vec{k} , the set of types present T , and the fraction of each type \vec{f} . For those interested in studying the agents of a scrip system, knowing the fraction of agents using each strategy can provide a window into the preferences of those agents. For system designers, this knowledge is useful because, as we show in Section 7, how much money the system can handle without crashing depends on the fraction of agents of each type.

In equilibrium, the distribution of money has the form described in Lemma 4.1. Note that, in general, we do not expect to see exactly this distribution at any given time, but it follows from Theorem 4.1 that, after sufficient time, the distribution is unlikely to be very far from it. Does this distribution help us identify the strategies and types of agents?

As a first step to answering this question, given a distribution of money d (where $d(i)$ is the fraction of agents with i dollars) such that $d(i)$ is rational for all i (this constraint is necessary of $d(i)$ is to represent the fraction of agents with i dollars in a real system), suppose that the maximum amount of money to which d gives positive probability is K_d . A vector \vec{f} of length $K_d + 1$ whose components are all rational numbers, where f_i is intuitively the fraction of agents playing the threshold strategy s_i , is an *explanation* of d if there exists a λ such that

$$d(j) = \sum_i d_\lambda(i, j),$$

where

$$d_\lambda(i, j) = f_i \lambda^j / \left(\sum_{l=0}^i \lambda^l \right) \quad (7)$$

if $j \leq i$ and 0 otherwise. Note that Equation (7) is very similar to Equation (2) from Lemma 4.1. In the following lemma, we show why we call \vec{f} an explanation: given a distribution d and an explanation \vec{f} we can find a game G where \vec{f} is the fraction of agents of each type and d is the equilibrium distribution of money (by which we mean that the value of d^* in Lemma 4.1 is such that $d(i) = \sum_t d^*(t, i)$).

Lemma 9.1. *If \vec{f} is an explanation for d , then there exists a game $G = (T, \vec{f}, h, m, n)$ and vector \vec{k} of thresholds such that \vec{k} is an ε -Nash equilibrium for G and the equilibrium distribution of money is d .*

PROOF. Let $T = \{0, \dots, K_d\}$, h be the minimum integer such that $hd(i)$ is an integer for all i , $m = \sum_i id(i)$, and \vec{k} be such that $k_i = i$. For each type i , choose β_i , χ_i , and ρ_i arbitrarily, subject to the constraint that $\beta\chi/\rho = 1$ (so that, by definition, $\omega_i = 1$ for all types i). Finally, choose an arbitrary n .

By Lemma 5.1, for any n , an optimal threshold policy in the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), i}$ for an agent of type i is s_κ , where κ is the maximum value such that

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u, p_d)}] \gamma_t. \quad (4)$$

Fix δ_i and γ_i , and let $g(\kappa)$ be the sequence of values of the right hand side of Equation (4) for natural numbers κ . Recall that the random variable $J(\kappa, p_u, p_d)$ represents the round at which an agent starting with k dollars runs out of money. Since $J(0, p_u, p_d) = 0$ for all histories, $g(0) = \gamma_t$. The time at which an agent runs out of money is increasing in his initial amount of money. Thus, $J(\kappa, p_u, p_d)$ is a strictly decreasing function of κ , so $g(\kappa)$ is also strictly decreasing. Choose α_i such that $g(i+1) < \alpha_i < g(i)$.

Thus, we have established parameters $(\alpha_i, \beta_i, \gamma_i, \delta_i, \chi_i, \rho_i)$ for each type i so that s_i an optimal policy for agents of type i in the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), i}$. By Theorem 5.1, taking n and the δ_i sufficiently large makes \vec{k} a ε -Nash equilibrium for G . By Lemma 4.1, the equilibrium distribution of money is d . ■

In general, there is not a unique explanation of a distribution d . Say that a distribution of money d is *fully-supported* if there do not exist i and j such that $i < j$, $d(j) > 0$, and $d(i) = 0$. For any game G , if all agents play threshold strategies then the resulting distribution will be fully-supported because it has the form given in Lemma 4.1. As the following lemma shows, a fully-supported distribution can be explained in an infinite number of different ways.

Lemma 9.2. *If d is a fully-supported distribution of money with finite support, there there exist an infinite number of explanations of d .*

We defer the proof of Lemma 9.2 to Appendix D.

Lemma 9.2 shows that d has an infinite number of explanations. Lemma 9.1 shows that we can find an (approximate) equilibrium corresponding to each of them. The explanations \vec{f} we construct in the proof of Lemma 9.2 seem unnatural;

typically $f_i > 0$ for all i . We are interested in a more parsimonious explanation, one that has a small support (i.e., the number of thresholds i for which $f_i > 0$ is small), for reasons the following lemma makes clear.

Lemma 9.3. *Let \vec{f} be an explanation for d . If s is the size of the support of \vec{f} , then any other explanation will have a support of size at least $K_d - s$.*

PROOF. Suppose that \vec{f} is an explanation for d . By Lemma 9.1, there is a game $G = (T, \vec{f}, h, m, n)$ and vector \vec{k} of thresholds such that \vec{k} is an ϵ -Nash equilibrium for G and the equilibrium distribution of money is d . Moreover, the proof of Lemma 9.1 shows that we can take $T = \{0, \dots, K_d\}$, $k_i = i$, and $\omega_i = 1$ for each type $i \in T$. By Equation (2) in Lemma 4.1, $d^*(t, i) = f_i \lambda^i q(t, i) / \sum_{j=0}^{k_t} \lambda^j q(t, j)$, where λ is the (unique) value that satisfies Equation (3). We first show that if $f_{i-1} = 0$, then $d(i)/d(i-1) = \lambda$. Since, for all i , $\omega_i = 1$, it is immediate from Equation (8) in the appendix that $q(i, j) = q(i, j')$ for all j and j' . Thus, the q terms cancel, so $d^*(i, j) = f_i \lambda^j / \sum_{l=0}^{k_i} \lambda^l$. Let $b_i = f_i / \sum_{l=0}^{k_i} \lambda^l$; then $d^*(i, j) = \lambda^j b_i$. Only agents with a threshold of at least j can have j dollars, so

$$d(j) = \sum_i d^*(i, j) = \sum_{\{t_l: l \geq j\}} d^*(l, j) = \sum_{\{t_l: l \geq j\}} b_l \lambda^j = B_j \lambda^j,$$

where $B_j = \sum_{\{t_l: l \geq j\}} b_l$. If $f_{i-1} = 0$, then $B_i = B_{i-1}$, so $d(i)/d(i-1) = \lambda$.

Since s strategies get positive probability according to \vec{f} , at least $k - s$ of the ratios $d(i)/d(i-1)$ with $1 \leq i \leq K_d$ must have value λ . Any other explanation \vec{f}' will have different coefficients f_i in Equation (7), so the value λ' satisfying it will also differ (since the requirement that $d(K_d) = d_{\lambda}(K_d, K_d)$ uniquely defines a value of λ). This means that the $K_d - s$ ratios with value λ must correspond to strategies i such that $f_i > 0$. Thus, the support of any other explanation must be at least $K_d - s$. ■

If $s \ll K_d$, Lemma 9.3 gives us a strong reason for preferring the minimal explanation (i.e., the one with the smallest support); any other explanation will involve significantly more types of agents being present. For $s = 3$ and $K_d = 50$, the smallest explanation has a support of three thresholds, and thus requires three types; the next smallest explanation requires at least 47 types. Thus, if the number of types of agents is relatively small, the minimal explanation will be the correct one.

The proof of Lemma 9.3 also gives us an algorithm for finding this minimal explanation. Since $d(i) = B_i \lambda^i$, taking logs of both sides, $\log d(i) = \log B_i + i \log \lambda$. Because B_i is constant in ranges of i where $f_{t_i} = 0$, a plot of $\log d(i)$ will be a line with slope λ in these ranges. Thus, the minimal explanation can be found by finding the minimum number of lines of constant slope that fit the data. For a simple example of how such a distribution might look, Figure 13 shows an equilibrium distribution of money for the game

$$(\{(.05, 1, 1, .95, 1), (.15, 1, 1, .95, 1)\}, (.3, .7), 10, 4, 100)$$

so the only difference between the types is that it costs the second type three times as much to satisfy a request) and the equilibrium strategy profile (20, 13). Figure 14 has the same distribution plotted on a log scale. Note the two lines with the same slope (λ) and the break at 13.




Fig. 13. Distribution of money with two types of agents.




Fig. 14. Log of the distribution of money with two types of agents.

This procedure allows us to infer a distribution of money the minimal explanation of the number of types of agents: the fraction of the population composed of each type, and the strategy each type is playing. (Note that we cannot distinguish multiple types with a shared ω_t playing the same strategy.) We would like to use this information to learn about the preferences of agents: their values of α_t , γ_t , and δ_t . Lemma 5.1 shows how we can do this. Once we find an explanation, the value of λ determines p_u^t and p_d^t for each type t . Then Equation 4 puts constraints on the values of α_t , β_t , and γ_t . Over time, if T , the set of types, remains constant, but \vec{f} , n , and m all vary as agents join and leave the system, a later observation with a slightly would give another equilibrium with new constraints on the types of the agents. A number of such observations potentially reveal enough information to allow strong inferences about agent types.

Thus far we have implicitly assumed that there are only a small number of types of agents in a system. Given that a type is defined by six real numbers, it is perhaps

more reasonable to assume that each agent has a different type, but there is a small number of “clusters” of agents with similar types. For example, we might believe that generally agents either place a high value or a low value on receiving service. While the exact value may vary, the types of two low-value agents or two high-value agents will be quite similar. We have also assumed in our analysis that all agents play their optimal threshold strategy. However, computing this optimum may be too difficult for many agents. Even ignoring computational issues, agents may have insufficient information about their exact type or the exact types of other agents to compute the optimal threshold strategy. Both the assumption that there are a few clusters of agents with similar, but not identical, types and the assumption that agents do not necessarily play their optimal threshold strategy, but do play a strategy close to optimal, lead to a similar picture of a system, which is one that we expect to see in practice: we will get clusters of agents playing similar strategies (that is, strategies with thresholds clustered around one value), rather than all agents in a cluster playing exactly the same strategy. This change has relatively little impact on our results. Rather than seeing straight lines representing populations with a sharp gap between them, as in Figure 14, we expect slightly curved lines representing a cluster of similar populations, with somewhat smoother transitions.

10. DISCUSSION

We have given a formal analysis of a scrip system and have shown that approximate equilibria exist in threshold strategies and that the distribution of money in these equilibria is given by relative entropy. As part of our equilibrium argument, we have shown that the best-reply function is monotone. This proves the existence of equilibria in pure strategies and permits efficient algorithms to compute these equilibria. We have also examined some of the practical implications of these theoretical results. For those interested in studying the agents of scrip systems, our characterization of equilibrium distribution of money forms the basis for techniques relevant to inferring characteristics of the agents of a scrip system from the distribution of money. For a system designer, our results on optimizing the money supply provide a simple maxim: keep adding money until the system is about to experience a monetary crash.

We have also seen that our model can be used to understand the effects of non-standard agent behavior on a scrip system. It provides insight into the effects of altruists and hoarders on a scrip system and guidance to system designers for dealing with them (less and more money respectively). Sybils are generally bad, but can typically be discouraged by imposing a moderate cost and possibly biasing the process for selecting a volunteer. On the other hand, collusion tends to be a net benefit and should be encouraged. Indeed, the entire purpose of the system is to allow users to collude and provide each other with service despite incentives to free ride.

Our model makes a number of assumptions that are worthy of further discussion. Some of the simplifying assumptions can be relaxed without significant changes to our results (albeit at the cost of greater strategic and analytic complexity). At a high level, our results show the system converges to a steady state when agents

follow simple threshold strategies and that there is in fact an equilibrium in these strategies. If, for example, rather than all requests having the same value to agent (γ_t), the value of a request is stochastic, agents might wish to have thresholds for each type of request. This would allow an agent to forgo a low-valued request if he is low on money. This makes the space of agent strategies larger and significantly complicates the proofs in the appendix, but this high-level characterization still holds.

The most significant assumption we make is that prices are fixed. However, our results provide insight even if we relax this assumption. With variable prices, the behavior of the system depends on the value of β , the probability that an agent can satisfy a request. For large β , where there are a large number of agents who can satisfy each request, we expect the resulting competition to effectively produce a fixed price, so our analysis applies directly. For small β , where there are few volunteers for each request, variable prices can have a significant impact.

However, allowing prices to be set endogenously, by bidding, has a number of negative consequences. For one thing, it removes the ability of the system designer to optimize the system using monetary policy. In addition, for small β , it is possible for colluding agents to form a cartel to fix prices on a resource they control. It also greatly increases the strategic complexity of using the system: rather than choosing a single threshold, agents need an entire pricing scheme. Finally, the search costs and costs of executing a transaction are likely to be higher with variable prices. Thus, we believe that adopting a fixed price or a small set of fixed prices is often a reasonable design decision.

We believe there is often a happy medium between a single, permanent fixed price and prices that change freely from round to round; indeed, our advice to system designers points naturally toward it. In particular, our advice about how to optimize the amount of money relies on experimentation and observation to determine what agents are doing and what their utilities are. This information then tells the designer how much money she should provide. Since adjusting the amount of money is equivalent to adjusting prices, the designer could incorporate this process into a price setting rule. Depending on the nature of the system, this could either be done manually over time (if the information is difficult to gather and analyze) or automatically (if the information gathering and analysis can itself be automated). From this perspective, a monetary crash, though real, is not something to be feared. Instead, it is just a strong signal that the current price, while probably not too far off from a very good price, requires adjustment. Naturally, this relies on a process that proceeds slow enough that agents myopically ignore the effects of future price changes in determining their current action.

Our model provides only the beginning of a full understanding of scrip systems. Many interesting open questions remain for future work. To name a few:

- Our model makes a number of strong predictions about the agent strategies, distribution of money, and effects of variations in the money supply. It also provides techniques to help analyze characteristics of agents of a scrip system. It would be interesting to test these predictions on a real functioning scrip system to either validate the model or gain insight from where its predictions are incorrect.
- In many systems there are overlapping communities of various sizes that are sig-

nificantly more likely to be able to satisfy each other’s requests. For example, in a P2P filesharing system, people are more likely to be able to satisfy the requests of others who share the same interests. It would be interesting to investigate the effect of such communities on the equilibrium of our system.

—It seems unlikely that altruism and hoarding are the only two types of “irrational” behavior we will find in real systems. Are there other major types that our model can provide insight into? Furthermore, it seems natural that the behavior of a very small group of agents should not be able to change the overall behavior of the system. Can we prove results about equilibria and utility when a small group follows an arbitrary strategy? This is particularly relevant when modeling attackers. See [Abraham et al. 2006] for general results in this setting.

11. ACKNOWLEDGMENTS

We would like to thank Randy Farmer, Peter Harremoës, Shane Henderson, Jon Kleinberg, David Parkes, Dan Reeves, Emin Gün Sirer, Michael Wellman, and anonymous referees for helpful suggestions, discussions, and criticisms. Eric Friedman, Ian Kash, and Joseph Halpern are supported in part by NSF under grant ITR-0325453. Joseph Halpern is also supported in part by NSF under grants CTC-0208535, IIS-0812045, and IIS-0534064; by ONR under grant N00014-01-10-511; by the DoD Multidisciplinary University Research Initiative (MURI) program administered by the ONR under grants N00014-01-1-0795 and N00014-04-1-0725; and by AFOSR under grants F49620-02-1-0101, FA9550-08-1-0438, FA9550-05-1-0055, and FA9550-09-1-0266. Eric Friedman is also supported in part by NSF under grant CDI-0835706.

A. PROOF OF THEOREM 4.1

Given a Markov chain \mathcal{M} over a state space X and state $s \in \mathcal{S}$, let $I_{\vec{x}, \vec{y}}^r$ be the random variable that is 1 if \mathcal{M} is in state \vec{y} at time r and the chain started in state \vec{x} and 0 otherwise. Then $\lim_{r \rightarrow \infty} \Pr(I_{\vec{x}, \vec{y}}^r = 1)$ is the limit probability of being in state \vec{y} given that the Markov chain starts in state \vec{x} . In general, this limit does not exist. However, there are well-known conditions under which the limit exists, and is independent of the initial state \vec{x} . A Markov chain is said to be *irreducible* if every state is reachable from every other state; it is *aperiodic* if, for every state \vec{x} , there exist two cycles from \vec{x} to itself such that the gcd of their lengths is 1.

Theorem A.1. [Resnick 1992] *If \mathcal{M} is a finite, irreducible, and aperiodic Markov chain over state space X , then there exists a d such that, for all \vec{x} and $\vec{y} \in X$, $\lim_{r \rightarrow \infty} \Pr(I_{\vec{x}, \vec{y}}^r = 1) = d$.*

Thus, if we can show that \mathcal{M} is finite, irreducible, and aperiodic, then the limit distribution exists and is independent of the start state \vec{x} . This is shown in the following lemma.

Lemma A.1. *If there are at least three agents, then \mathcal{M} is finite, irreducible, and aperiodic and therefore has a limit distribution π .*

PROOF. \mathcal{M} is clearly finite since X is finite. We prove that it is irreducible by showing that state \vec{y} is reachable from state \vec{x} by induction on the distance

$w = \sum_{i=1}^n |x_i - y_i|$ (i.e., the sum of the absolute differences in the amount of money each person has in states \vec{x} and \vec{y}). If $z = 0$, then $\vec{x} = \vec{y}$ so we are done. Suppose that $w > 0$ and all pairs of states that are less than w apart are reachable from each other. Consider a pair of states \vec{x} and \vec{y} such that the distance from \vec{x} to \vec{y} is w . Since $w > 0$ and the total amount of money is the same in all states, there must exist i_1 and i_2 such that $x_{i_1} > y_{i_1}$ and $x_{i_2} < y_{i_2}$. Thus, in state \vec{y} , i_1 is willing to work (since he has strictly less than the threshold amount of money) and i_2 has money to pay him (since i_2 has a strictly positive amount of money). The state \vec{z} that results from i_1 doing work for i_2 in state \vec{y} is of distance $w - 2$ from \vec{x} . By the induction hypothesis, \vec{z} is reachable from \vec{x} . Since \vec{y} is clearly reachable from \vec{z} , \vec{y} is reachable from \vec{x} .

Finally, we must show that \mathcal{M} is aperiodic. Suppose \vec{x} is a state such that there exist three agents i_1 , i_2 , and i_3 where i_1 has more than 0 dollars and i_2 and i_3 have less than their threshold amount of money. There must be such a state by our assumption that m is “interesting.” Clearly there is a cycle of length 2 from \vec{x} to itself: i_2 does work for i_1 and then i_2 does work for i_1 . There is also a cycle of length 3: i_2 does work for i_1 , i_3 does work for i_2 , then i_1 does work for i_3 . ■

We next give an explicit formula for the limit distribution. Recall that in the special case discussed in the main text, β_t , χ_t , and ρ_t were the same for all types, so the transition probabilities were symmetric and the limit distribution was uniform. While with more general values they are no longer symmetric, they still have significant structure that allows us to give a concise description of the limit distribution.

Lemma A.2. *For all states \vec{x} of \mathcal{M} , let $w_{\vec{x}} = \prod_i (\beta_{\tau(i)} \chi_{\tau(i)} / \rho_{\tau(i)})^{x_i}$, and let $Z = \sum_{\vec{y}} w_{\vec{y}}$. Then the limit distribution of \mathcal{M} is $\pi(\vec{x}) = w_{\vec{x}} / Z$.*

PROOF. Define π by taking $\pi(\vec{x}) = w_{\vec{x}} / Z$, where $w_{\vec{x}}$ and Z are as in the statement of the lemma. If $T_{\vec{x}\vec{y}}$ is the probability of transitioning from state \vec{x} to state \vec{y} , it is well known that it suffices to show that π satisfies the *detailed balance condition* [Resnick 1992], i.e., $\pi(\vec{x}) T_{\vec{x}\vec{y}} = \pi(\vec{y}) T_{\vec{y}\vec{x}}$ for all states \vec{x} and \vec{y} and π is a probability measure. The fact that π is a probability measure is immediate from its definition. To check the first condition, let \vec{x} and \vec{y} be adjacent states such that \vec{y} is reached from \vec{x} by i spending a dollar and j earning a dollar. This means that for the transition from \vec{x} to \vec{y} to happen, i must be chosen to spend a dollar and j must be able to work and chosen to earn the dollar. Similarly for the reverse transition to happen, j must be chosen to spend a dollar and i must be able to work and chosen to earn the dollar. All other agents have the same amount of money in each state, and so will make the same decision in each state. Thus the probabilities associated with each transition differ only in the relative likelihoods of i and j being chosen at each point. These may differ for three reasons: one might be more likely to be able to satisfy requests (β), to want to make requests (ρ), or to be chosen to satisfy requests (χ). Thus, for some p , which captures the effect of other agents volunteering on the likelihood of i and j being chosen, we can write the transition probabilities as $T_{\vec{x}\vec{y}} = \rho_{\tau(i)} \beta_{\tau(j)} \chi_{\tau(j)} p$ and $T_{\vec{y}\vec{x}} = \rho_{\tau(j)} \beta_{\tau(i)} \chi_{\tau(i)} p$. From

the definition of π , we have that

$$\frac{\pi(\vec{x})}{\pi(\vec{y})} = \frac{\beta_{\tau(i)}\chi_{\tau(i)}\rho_{\tau(j)}}{\rho_{\tau(i)}\beta_{\tau(j)}\chi_{\tau(j)}} = \frac{T_{\vec{y}\vec{x}}}{T_{\vec{x}\vec{y}}}.$$

Thus, $\pi(\vec{x})T_{\vec{x}\vec{y}} = \pi(\vec{y})T_{\vec{y}\vec{x}}$, as desired. ■

Note that for the special case considered in the main text, Lemma A.2 shows that the limit distribution is the uniform distribution.

The limit distribution tells us the long run probability of being in a given state. Theorem 4.1 does not mention states directly, but rather the distributions of money associated with a state. In order to prove the theorem, we need to know the probability of being in some state associated with a given distribution. This is established in the following lemma.

Lemma A.3. *Let π be the limit distribution from Lemma A.2, and let $V(d) = H(d) - H(\vec{f}) - \log Z + \sum_t \sum_{i=0}^{k_t} id(t, i) \log \omega_t$ (where H is the standard entropy function; that is, $H(d) = \sum_{t,i} d(t, i) \log d(t, i)$). For all $d \in \Delta_{\vec{f}, m, \vec{k}}$, either $\pi(\{\vec{x} \mid d^{\vec{x}} = d\}) = 0$ or $F(hn)e^{hnV(d)} \leq \pi(\{\vec{x} \mid d^{\vec{x}} = d\}) \leq G(hn)e^{hnV(d)}$, where F and G are polynomials.*

PROOF. Before computing the probability of being in such a state, we first compute the number of states. It is possible that there is no state \vec{x} such that $d = d^{\vec{x}}$ (e.g., if hn is odd and d has half the agents with 0 dollars). If there is such a state \vec{x} , each such state has $hnd(t, i)$ agents of type t with i dollars. Thus, the number of states \vec{x} with $d = d^{\vec{x}}$ is the number of ways to divide the agents into groups of these sizes. Since there are hnf_t agents of type t , the number of such states is

$$\prod_t \binom{hnf_t}{hnd(t, 0), \dots, hnd(t, k_t)}.$$

To complete the proof, we use the fact (shown in the proof of Lemma 3.11 of [Grove et al. 1994]) that

$$\frac{1}{F(hn)} e^{hnf_t H(d_t)} \leq \binom{hnf_t}{hnd(t, 0), \dots, hnd(t, k_t)} \leq G(hn) e^{hnf_t H(d_t)},$$

where F and G are polynomial in hn , and d_t is the distribution restricted to a single type t (i.e., $d_t(i) = d(t, i) / \sum_i d(t, i)$). The (*generalized*) *grouping property* [Cover and Thomas 1991] of entropy allows us to express $H(d)$ in terms of the entropy of the distributions for each fixed t , or the $H(d_t)$. Because $f_t = \sum_i d(t, i)$, this has the particularly simple form $H(d) = H(\vec{f}) + \sum_t f_t H(d_t)$. Thus, up to a polynomial factor, the number of such states is

$$\prod_t e^{hnf_t H(d_t)} e^{hn(\sum_t f_t H(d_t))} = e^{hn(H(d) - H(\vec{f}))}.$$

By Lemma A.2, each of these states has the same probability $\pi(\vec{x})$. Thus, dropping the superscript \vec{x} on $d^{\vec{x}}$ for brevity, the probability of being in such a

state is:

$$\begin{aligned}
e^{hn(H(d)-H(\vec{f}))}\pi(\vec{x}) &= e^{hn(H(d)-H(\vec{f}))} \prod_i (\beta_i \chi_t / \rho_i)^{x_i} / Z \\
&= e^{hn(H(d)-H(\vec{f}))} Z^{-hn} \prod_i (\omega_{t_i})^{x_i} \\
&= e^{hn(H(d)-H(\vec{f}))} Z^{-hn} \prod_t \prod_{i=0}^{k_t} (\omega_t)^{hni d(t,i)} \\
&= e^{hn(H(d)-H(\vec{f}))} Z^{-hn} \prod_t \prod_{i=0}^{k_t} e^{hni d(t,i) \log \omega_t} \\
&= e^{hn(H(d)-H(\vec{f}))- \log Z + \sum_t \sum_{i=0}^{k_t} id(t,i) \log \omega_t} \\
&= e^{hnV(d)}
\end{aligned}$$

■

Theorem 4.1 says that there exists a $q \in \Delta_{\vec{f},m,\vec{k}}$ (i.e., a probability distribution on agent types t and amounts of money i) with certain properties. We now define the appropriate q . Let

$$q(t, i) = (\omega_t)^i / \left(\sum_t \sum_{j=0}^{k_t} (\omega_t)^j \right). \quad (8)$$

It is not immediately clear why this is the right choice of q . As the following lemma shows, this definition allows us to characterize the distribution that maximizes the probability of being in a state corresponding to that distribution (as given by Lemma A.3) in terms of relative entropy.

Lemma A.4. *The unique maximum of $V(d) = H(d) - H(\vec{f}) - \log Z + \sum_t \sum_{i=0}^{k_t} id_i^t \log \omega_t$ on $\Delta_{\vec{f},m,\vec{k}}$ occurs at d_q^* .*

PROOF. For brevity, we drop the superscript \vec{x} on d and let $Y = \sum_t \sum_i (\omega_t)^i$.

$$\begin{aligned}
 \operatorname{argmax}_d V(d) &= \operatorname{argmax}_d (H(d) - H(\vec{f}) - \log Z + \sum_t \sum_{i=0}^{k_t} i d(t, i) \log \omega_t) \\
 &= \operatorname{argmax}_d (H(d) + \sum_t \sum_{i=0}^{k_t} i d(t, i) \log \omega_t) \\
 &= \operatorname{argmax}_d \sum_t \sum_{i=0}^{k_t} [-d(t, i) \log d(t, i) + i d(t, i) \log \omega_t] \\
 &= \operatorname{argmax}_d \sum_t \sum_{i=0}^{k_t} [-d(t, i) \log d(t, i) + d(t, i) \log(q(t, i)Y)] \\
 &= \operatorname{argmax}_d \sum_t \sum_{i=0}^{k_t} [-d(t, i) \log d(t, i) + d(t, i) \log q(t, i) + d(t, i) \log Y] \\
 &= \operatorname{argmax}_d \log Y + \sum_t \sum_{i=0}^{k_t} [-d(t, i) \log d(t, i) + d(t, i) \log q(t, i)] \\
 &= \operatorname{argmin}_d \sum_t \sum_{i=0}^{k_t} [d(t, i) \log d(t, i) - d(t, i) \log q(t, i)] \\
 &= \operatorname{argmin}_d \sum_t \sum_{i=0}^{k_t} d(t, i) \log \frac{d(t, i)}{q(t, i)} \\
 &= \operatorname{argmin}_d H(d||q).
 \end{aligned}$$

By definition, d_q^* minimizes $H(d||q)$. It is unique because H (and thus V) is a strictly concave function on a closed convex set. ■

Lemma A.4 tells us that the most likely distributions of money to be observed are those with low relative entropy to q . Among all distributions in $\Delta_{\vec{f}, m, k}$, relative entropy is minimized by d_q^* . However, given n , it is quite possible that d_q^* is not $d^{\vec{x}}$ for any \vec{x} . For example, if $d_q^*(t, i) = 1/3$ for some t and i , but $f_t h n = 16$, then $d^{\vec{x}}(t, i) = d_q^*(t, i)$ only if exactly $16/3$ agents of type t to have i dollars, which cannot be the case. However, as the following lemma shows, for sufficiently large n , we can always find a $d^{\vec{x}}$ that is arbitrarily close to d_q^* . For convenience, we use the 1-norm as our notion of distance.

Lemma A.5. *For all ϵ , there exists n_ϵ such that, if $n > n_\epsilon$, then for some state \vec{x} , $\|d^{\vec{x}} - d_q^*\| < \epsilon$.*

PROOF. Given n , we construct $d \in \Delta_{\vec{f}, m, k}$ that is of the form $d^{\vec{x}}$ and is close to d_q^* in a number of steps. As a first step, for all t and i , let $d_1(t, i)$ be the result of rounding $d_q^*(t, i)$ to the nearest $1/hn$ (where ties are broken arbitrarily). The function d_1 may not be in $\Delta_{\vec{f}, m, k}$; we make minor adjustments to it to get a function in $\Delta_{\vec{f}, m, k}$. First, note that we may not have as is $d_1(t, i)$ for all t and i , we $\sum_i d_1(t, i) \neq f_t$. Since f_t is a multiple of $1/hn$, can get a function d_2 that

satisfies these constraints by modifying each term $d_1(t, i)$ by either adding $1/hn$ to it, subtracting $1/hn$ from it, or leaving it alone. Such a function d_2 may still violate the final constraint that $\sum_{t,i} id_2(t, i) = m$. We construct a function d_3 that satisfies this constraint (while continuing to satisfy the constraint that $\sum_i d_3(t, i) = f_t$) as follows. Note that if we increase $d_2(t, i)$ by $1/hn$ and decrease $d_2(t, j)$ by $1/hn$, then we keep the keep $\sum_i d_2(t, i) = f_t$, and change $\sum_i id_2(t, i)$ by $(i - j)/hn$. Since each term $d_2(t, i)$ is a multiple of $1/hn$ and m is a multiple of $1/h$, we can perform these adjustments until all the constraints are satisfied.

The rounding to create d_1 changed each $d_1(t, i)$ by at most $1/hn$, so $\|d_q^* - d_1\|_1 \leq (\sum_t k_t + 1)/hn$. Since, each term $d_1(t, i)$ was changed by at most $1/hn$ to obtain $d_2(t, i)$, we have $\|d_1 - d_2\|_1 \leq (\sum_t k_t + 1)/hn$. Let $c = \max_t(\max(k_t - m, m))$. Each movement of up to $1/hn$ in the creation of d_1 and d_2 altered m by at most c/hn . Thus at most $2c$ movements are needed in the creation of d_3 for each pair (t, i) . Therefore, $\|d_2 - d_3\|_1 \leq (\sum_t k_t + 1)2c/hn$. By the triangle inequality, $\|d_q^* - d_3\|_1 \leq (\sum_t k_t + 1)(2c + 2)/hn$, which is $O(1/n)$. Hence, for n_ϵ sufficiently large, the resulting d_3 will always be within distance ϵ of d_q^* .

Finally, we need to show that $d_3 = d^{\vec{x}}$ for some \vec{x} . Each $d_3(t, i)$ is a multiple of $1/hn$. There are hn agents in total, so we can find such an \vec{x} by taking any allocation of money such that $d_3(t, i)hn$ agents of type t have i dollars. ■

We are now ready to prove Theorem 4.1. We repeat the statement here for the reader's convenience.

THEOREM 4.1. *For all games $(T, \vec{f}, h, m, 1)$, all vectors \vec{k} of thresholds, and all $\epsilon > 0$, there exist $q \in \Delta_{\vec{f}, m, \vec{k}}$ and n_ϵ such that, for all $n > n_\epsilon$, there exists a round r^* such that, for all $r > r^*$, we have $\Pr(I_{q, n, \epsilon}^r = 1) > 1 - \epsilon$.*

PROOF. From Lemma A.2, we know that, after a sufficient amount of time, the probability of being in state \vec{x} will be close to $\pi_{\vec{x}} = w_{\vec{x}}/Z$. Since \mathcal{M} converges to a limit distribution, it is sufficient to show that the theorem holds in the limit as $r \rightarrow \infty$. If the theorem holds in the limit for some $\epsilon' < \epsilon$, then we can take r large enough that the L1 distance between the distribution of the chain at time r and the limit distribution (i.e. treating the distributions as vectors and computing the sum of the absolute values of their differences) is at most $\epsilon - \epsilon'$.

The remainder of the proof is essentially that of Theorem 3.13 in [Grove et al. 1994] (applied in a very different setting). Let $V(d) = H(d) - H(\vec{f}) - \log Z + \sum_t \sum_{i=0}^{k_t} id(t, i) \log \omega_t$. We show there exists a value v_L such that, for all states \vec{x} such that $d^{\vec{x}}$ is not within ϵ of d_q^* , we have $V(d^{\vec{x}}) \leq v_L$, and a value $v_H > v_L$ such that $v_H = V(d^{\vec{y}})$ for some point \vec{y} such that $d^{\vec{y}}$ is within distance ϵ of d_q^* . Lemma A.3 then shows that it is exponentially more likely that $d^{\vec{x}^r} = d^{\vec{y}}$ than any distribution d such that $V(d) \leq v_L$. If $\vec{x}^r = \vec{y}$ then $I_{q, n, \epsilon}^r = 1$, and if $I_{q, n, \epsilon}^r = 0$ then $V(d^{\vec{x}^r}) \leq v_L$, so this suffices to establish the theorem.

By Lemma A.4, the unique maximum of V on $\Delta_{\vec{f}, m, \vec{k}}$ occurs at d_q^* . The set $\{d \in \Delta_{\vec{f}, m, \vec{k}} \mid \|d_q^* - d\|_2 \geq \epsilon\}$ is closed. V is a continuous function, so it takes some maximum v_L on this set. Pick some v_H such that $v_L < v_H < V(d_q^*)$. By the continuity of V , there exists an ϵ such that if $\|d_q^* - d\|_1 < \epsilon$ then $V(d) \geq v_H$. By Lemma A.5, for sufficiently large n , there is always some \vec{x} such that $\|d_q^* - d^{\vec{x}}\|_1 < \epsilon$.

Thus, for some $\vec{x} \in X_{\varepsilon, q}$, $V(d^{\vec{x}}) \geq v_H$.

$\Pr(I_{q, n, \varepsilon}^r = 1) \geq \Pr(\vec{x}^r \in \{\vec{y} \mid d^{\vec{y}} = d^{\vec{x}}\})$. By Lemma A.3, $\Pr(I_{q, n, \varepsilon}^r = 1)$ is at least $1/F(hn)e^{hnV(d^{\vec{x}})} \geq 1/F(hn)e^{hnv_H}$. Now consider a \vec{y} such that $I_{q, n, \varepsilon}(\vec{y}) = 0$. By Lemma A.3, the probability that $d^{\vec{x}^r} = d^{\vec{y}}$ is at most $G(hn)e^{hnV(d^{\vec{y}})} \leq G(hn)e^{hnv_L}$. There are at most $(hn + 1)^{\sum_t (k_t + 1)}$ such points, a number which is polynomial in hn . Thus, for $G'(hn) = G(hn)(hn + 1)^{\sum_t (k_t + 1)}$, the probability that $I_{q, n, \varepsilon}^r = 0$ is at most $G'(hn)e^{hnv_L}$. The ratio of these probabilities is at most

$$\frac{G'(hn)e^{hnv_L}}{\frac{1}{F(hn)}e^{hnv_H}} = \frac{G'(hn)F(hn)}{e^{hn(v_H - v_L)}}.$$

This is the ratio of a polynomial to an exponential, so the probability of seeing a distribution of distance greater than ε from d_q^* goes to zero as n goes to infinity. ■

B. PROOFS FROM SECTION 5

In this appendix, we provide the omitted proofs from Section 5.

The proof of Theorem 5.1 relies on modeling the game from the perspective of a single agent. Consider a vector \vec{k} of thresholds and the corresponding strategy profile $\vec{S}(\vec{k})$. Fix an agent i of type t . Assume that all the agents other than i continue play their part of $\vec{S}(\vec{k})$. What is i 's best response? Since the set of agents is large, i 's choice of strategy will have (essentially) no impact on the distribution of money. By Theorem 4.1, the distribution of money will almost always be close to a distribution d^* . Suppose, the distribution were exactly d^* . Since we know the exact distribution of money and the thresholds used by the other agents, we can calculate the number of each type of agent that wish to volunteer and thus the probabilities that our single agent will be able to earn or spend a dollar. Thus, by assuming the distribution of money is always exactly d^* , we can model the game from the perspective of agent i as a Markov Decision Process (MDP). We show in Lemma B.2 that this MDP has an optimal threshold policy. (Threshold policies are known as *monotone* policies in the more general setting where there are more than two actions.) We then prove that any optimal policy for the MDP is an ε -best reply to the strategies of the other agents in the actual game.

Taking notation from Puterman [1994], we formally define the MDP $\mathcal{P}_{G, \vec{S}(\vec{k}), t} = (S, A, p(\cdot \mid s, a), r(s, a))$ that describes the game where all the agents other than i are playing $\vec{S}(\vec{k})_{-i}$ and i has type t .

- $S = \{0, \dots, mhn\}$ is the set of possible states for the MDP (i.e., the possible amounts of money compatible with the distribution d^*).
- $A = \{0, 1\}$ is the set of possible actions for the agent, where 0 denotes not volunteering and 1 denotes volunteering iff another agent who has at least one dollar makes a request.
- p_u is the probability of earning a dollar, assuming the agent volunteers (given that all other agents have fixed their thresholds according to \vec{k} and the distribution of money is exactly d^*). Each agent of type t' who wishes to volunteer can do so with probability $\beta_{t'}$. Assuming exactly the expected number of agents are able to volunteer, $v_{t'} = \beta_{t'}(f_{t'} - d^*(t', k_{t'}))n$ agents of type t' volunteer. Note that

we are disregarding the effect of i in computing the $v_{t'}$, since this will have a negligible effect for large n . Using the $v_{t'}$ s, we can express p_u as the product of two probabilities: that some agent other than i who has a dollar is chosen to make a request and that i is the agent chosen to satisfy it. Thus,

$$p_u = \left(\sum_{t'} \rho_{t'} (f_{t'} - d^*(t', 0)) \right) \left(\frac{\chi_t \beta_t}{\sum_{t'} \chi_{t'} v_{t'}} \right). \quad (9)$$

— p_d is the probability of agent i having a request satisfied, given that agent i has a dollar. Given that all agents are playing a threshold strategy, if the total number n of agents is sufficiently large, then it is almost certainly the case that some agent will always be willing and able to volunteer. Thus, we can take p_d to be the probability that agent i will be chosen to make a request; that is,

$$p_d = \frac{\rho_t}{n} \quad (10)$$

— $r(s, a)$ is the (immediate) expected reward for performing action a in state s . Thus, $r(s, 0) = \gamma_t p_d$ if $s > 0$; $r(0, 0) = 0$; $r(s, 1) = \gamma_t p_d - \alpha_t p_u$ if $s > 0$; and $r(0, 1) = -\alpha_t p_u$.

— $p(s' | s, a)$ is the probability of being in state s' after performing action a in state s ; $p(s' | s, a)$ is determined by p_u and p_d ; specifically, $p(s + 1 | s, 1) = p_u$, $p(s - 1 | s, a) = p_d$ if $s > 0$, and the remainder of the probability is on $p(s | s, a)$ (i.e., $p(s | s, a) = 1 - (p(s + 1 | s, 1) + p(s - 1 | s, a))$).

— $u^*(s)$ is the expected utility of being in state s if agent i uses the optimal policy for the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$

— $u(s, a)$ is the expected utility for performing action a in state s , given that the optimal strategy is followed after this action;

$$u(s, a) = r(s, a) + \delta \sum_{s'=0}^{mhn} p(s' | s, a) u^*(s').$$

To prove Theorem 5.1, we need two preliminary lemmas about the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$.

Lemma B.1. *For the MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$, $u^*(s + 2) + u^*(s) \leq 2u^*(s + 1)$.*

PROOF. The MDP $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$ has an optimal stationary policy [Puterman 1994, Theorem 6.2.10] (a policy where the chosen action depends only on the current state). Let π be such a policy. Consider the policy π' starting in state $s + 1$ that “pretends” it actually started in state s and is following π . More precisely, if $s_0 = s + 1$ and $s_j > 0$ for $j = 0, \dots, k$, define $\pi'(s_0, s_1, \dots, s_k) = \pi(s_k - 1)$; otherwise, if $j \leq k$ is the least index such that $s_j = 0$, define $\pi'(s_0, \dots, s_k) = \pi(s_k)$. Given a history (s_0, \dots, s_k) , j is the random variable whose value is the minimum i such that $s_i = 0$ or ∞ if no such value exists. The definition of π' from π creates a bijection between histories that start in state $s + 1$ and histories that start in state s , such that if h' corresponds to h , the probability of history h' with policy π' is the same as the probability of h with policy π . Technically, making the mapping a bijection requires the introduction of a new state $0'$, which intuitively represents the state where the agent has zero dollars and missed an

opportunity to have a request satisfied last round because of it. More formally, we let $p(0' | 0, a) = p_d$ and $p(s | 0', a) = p(s | 0, a)$. With this change, the probabilities of corresponding histories are the same because the probability of transitioning from a state to the one “immediately below” it (where $s - 1$ is immediately below s , $0'$ is immediately below 0 , and $0'$ is immediately below itself) is always p_d , and the probability of transitioning from a state to the one “immediately above” it (where $s + 1$ is immediately above s , and 1 is immediately above $0'$) is always p_u .⁴

This argument shows that an agent starting with $s + 1$ dollars “pretending” to start with s will have the same expected reward each round as an agent who actually started with s dollars, except during the first round j in a history such that $s_j = 0$. Thus (treating j as a random variable), we have

$$u^*(s + 1) \geq u^*(s) + E[\delta^j \gamma_t].$$

Similarly, we can use π starting from state $s + 2$ to define a policy π'' starting from state $s + 1$, where i “pretends” he has one more dollar and is using π , up to the first round j' that he is chosen to make a request with π in a state where he has no money (in which case he can make the request with π started from $s + 2$, but cannot make it with π'' started from $s + 1$); from that point on, he uses π . For corresponding histories, the utilities of an agent starting with $s + 1$ dollars and following π'' and an agent starting with $s + 2$ dollars and following π will be the same, except during round j' the agent following π will have a request satisfied but the agent following π'' will not. Thus,

$$u^*(s + 1) \geq u^*(s + 2) - E[\delta^{j'} \gamma_t].$$

Since, if i uses π , he will run out of money sooner if he starts with s dollars than if he starts with $s + 2$ dollars,

$$E[\delta^j \gamma_t] > E[\delta^{j'} \gamma_t].$$

Thus, $u^*(s + 2) + u^*(s) \leq 2u^*(s + 1)$. ■

Lemma B.2. $\mathcal{P}_{G, \vec{s}(\vec{k}), t}$ has an optimal threshold policy.

PROOF. As shown by Puterman [1994, Lemma 4.7.1], it suffices to prove that $u(s, a)$ is subadditive. That is, we need to prove that, for all states s ,

$$u(s + 1, 1) + u(s, 0) \leq u(s + 1, 0) + u(s, 1). \quad (11)$$

We consider here only the case that $s > 0$ (the argument is essentially the same if $s = 0$). Because $s > 0$, $r(s + 1, a) = r(s, a)$, so (11) is equivalent to

$$\begin{aligned} & p_u u^*(s + 2) + p_d u^*(s) + (1 - p_u - p_d) u^*(s + 1) + p_d u^*(s - 1) + (1 - p_d) u^*(s) \\ & \leq p_d u^*(s) + (1 - p_d) u^*(s + 1) + p_u u^*(s + 1) + p_d u^*(s - 1) + (1 - p_u - p_d) u^*(s). \end{aligned}$$

This simplifies to

$$u^*(s + 2) + u^*(s) \leq 2u^*(s + 1),$$

⁴Note that this means that $0'$ is immediately below 0 but 1 is immediately above $0'$. This is intended, because $0'$ intuitively represents the state where the agent has 0 dollars and had a request go unsatisfied due to a lack of money in the previous round, so if he then earns a dollar he will have 1 dollar regardless of whether or not his request of two rounds previous was satisfied.

which follows from Lemma B.1. ■

We can now prove Lemma 5.1 and Theorem 5.1.

LEMMA 5.1. *Consider the games $G_n = (T, \vec{f}, h, m, n)$ (where T , \vec{f} , h , and m are fixed, but n may vary). There exists a k such that for all n , s_k is an optimal policy for $\mathcal{P}_{G_n, \vec{S}(\vec{k}), t}$. The threshold k is the maximum value of κ such that*

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u, p_d)}] \gamma_t, \quad (4)$$

where $J(\kappa, p_u, p_d)$ is a random variable whose value is the first round in which an agent starting with κ dollars, using strategy s_κ , and with probabilities p_u and p_d of earning a dollar and of being chosen given that he volunteers, respectively, runs out of money.

PROOF. Fix n . Suppose that an agent is choosing between a threshold of κ and a threshold of $\kappa + 1$. These policies only differ when the agent has κ dollars: he will volunteer with the latter but not with the former. If he volunteers when he has κ dollars and is chosen, he will pay a cost of α_t and he will have $\kappa + 1$ dollars. As in the proof of Lemma B.1, we can define a bijection on histories such that, in corresponding histories of equal probability, an agent who started with κ dollars and is using s_κ will always have one less dollar than an agent who started with $\kappa + 1$ dollars and is using $s_{\kappa+1}$, until the first round r in which the agent using $s_{\kappa+1}$ has zero dollars. This means that in round $r - 1$ the agent using $s_{\kappa+1}$ had a request satisfied but the agent using s_κ was unable to because he had no money. Thus, if the agent volunteers when he has κ dollars and pays a cost of α_t in the current round, the expected value of being able to spend that dollar in the future is $E[(1 - (1 - \delta_t)/n)^{J(\kappa+1, p_u, p_d)}] \gamma_t$. Since this expectation is strictly increasing in κ (an agent with more money takes longer to spend it all), the maximum κ such that Equation (4) holds is an optimal threshold policy.

Taking the maximum value of κ that satisfies Equation (4) ensures that, for the n we fixed, we chose the maximum optimal threshold. We now need to show that this maximum optimal threshold is independent of n , which we do by showing that the expecting utility of every threshold policy s_k is independent of n . The expected utility of a policy depends on the initial amount of money, but since an agent's current amount of money is a random walk whose transition probabilities are determined by p_u and p_d , there is a well-defined limit probability

$$x_i^* = \lim_{r \rightarrow \infty} \Pr(\text{agent has } i \text{ dollars in round } r)$$

determined by the ratio p_u/p_d (this is because the limit distribution satisfies the detailed balance condition: $x_i^* p_u = x_{i+1}^* p_d$). This distribution has the property that if the agent starts with i dollars with probability x_i^* , then in every round the probability he has i dollars is x_i^* . Thus, in each round his expected utility is $\gamma p_d(1 - x_0^*) - \alpha p_u(1 - x_k^*)$. We can factor out n to write $p_u = p'_u/n$ and $p_d = p'_d/n$ where p'_u and p'_d are independent of n . Note that $p_u/p_d = p'_u/p'_d$, so the x_i^* 's are independent of n . Thus, we can rewrite the agent's expected utility for each round as c/n , where $c = \gamma p'_d(1 - x_0^*) - \alpha p'_u(1 - x_k^*)$ is independent of n . Therefore, the

expected utility of s_k is

$$\sum_{r=0}^{\infty} \left(1 - \frac{1 - \delta_t}{n}\right)^r \frac{c}{n} = \frac{c}{1 - \delta_t},$$

which is independent of n . ■

THEOREM 5.1. *For all games $G = (T, \vec{f}, h, m, n)$, all vectors \vec{k} of thresholds, and all $\varepsilon > 0$, there exist n_ε^* and $\delta_{\varepsilon, n}^*$ such that for all $n > n_\varepsilon^*$, types $t \in T$, and $\delta_t > \delta_{\varepsilon, n}^*$, an optimal threshold policy for $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$ is an ε -best reply to the strategy profile $\vec{S}(\vec{k})_{-i}$ for every agent i of type t .*

PROOF. By Lemma B.2, $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$ has an optimal threshold policy. However, this might not be a best reply for agent i in the actual game if the other agents are playing $\vec{S}(\vec{k})$. $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$ assumes that the probabilities of earning or spending a dollar in a given round are always exactly p_u and p_d respectively. Theorem 4.1 guarantees only that, in the game, the corresponding probabilities are close to p_u and p_d with high probability after some amount of time that can depend on n . A strategy S for player i in G defines a policy π_S for $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$ in the obvious way; similarly, a policy for the MDP determines a strategy for player i in the game. The expected utility of π_S is close to $U_i(S, \vec{S}(\vec{k})_{-i})$, but is, in general, not equal to it, because, as we noted, p_u and p_d may differ from the corresponding probabilities in the game. They differ for three reasons: (1) they are close, but not identical; (2) they are only close with high probability, and (3) they are only close after some amount of time. As we now show, given ε , the difference in the expected utility due to each reason can be bounded by $\varepsilon/6$, so the expected utility of any strategy is within $\varepsilon/2$ of the value the corresponding policy in $\mathcal{P}_{G, \vec{S}(\vec{k}), t}$. Thus, an optimal strategy for the MDP is an ε -best reply.

As we have seen, the probabilities p_u and p_d are determined by the number of agents of each type that volunteer (i.e., the expressions $v_{t'}$ for each type t'). The distance between $d^{\vec{x}^r}$ and d^* bounds how much the actual number of agents of type t' that wish to volunteer in round r can differ from $v_{t'}/\beta_{t'}$. Even if exactly $v_{t'}/\beta_{t'}$ agents wish to volunteer for each type t , there might not be exactly $v_{t'}$ agents who actually volunteer because of the stochastic decision by nature about who can volunteer and because i cannot satisfy his own requests. However, for sufficiently large n , the effect on p_u and p_d from these two factors is arbitrarily close to zero. Applying Theorem 4.1, there exist n_1 and r_1 such that if there are at least n_1 agents, for all round $r > r_1$, $d^{\vec{x}^r}$ and d^* are sufficiently close that the difference between the utility of policy $\pi_{S'}$ in the MDP and $U_i((S', \vec{S}_{-i}))$ in rounds $r > r_1$ where d^* is sufficiently close is at most $\varepsilon/6$.

Note that the maximum possible difference in utility between a round of the MDP and a round of the game is $\gamma + \alpha$ (if agent i spends a dollar rather than earning one). Applying Theorem 4.1 again, for $e = \varepsilon/6(\gamma + \alpha)$, there exist n_2 and r_2 such that the probability of the distribution not being within e of d^* is less than e . Thus, the difference between the expected utility of policy $\pi_{S'}$ in the MDP and $U_i((S', \vec{S}_{-i}))$ in rounds $r > r_2$ where d^* is not sufficiently close is at most $e(\gamma + \alpha) = \varepsilon/6$.

Let $n_\varepsilon^* = \max(n_1, n_2)$ and $r^* = \max(r_1, r_2)$. The values of n_ε^* and r^* do not depend on δ , so we can take $\delta_{\varepsilon, n}^*$ to be sufficiently close to 1 that the total utility

from the first r^* rounds is at most $\varepsilon/6$, completing the proof of the theorem. ■

Recall that BR_G maps a vector \vec{k} describing the threshold strategy for each type to a vector \vec{k}' of best replies.

LEMMA 5.2. *Consider the family of games $G_m = (T, \vec{f}, h, m, n)$ and the strategies $\vec{S}(\vec{k})$, for $mhn < \sum_t f_t k_t hn$. For this family of game, $\lambda_{m, \vec{k}}$ is non-decreasing in m and non-increasing in \vec{k} ; $p_u^{m, \vec{k}}$ is non-decreasing in m and non-increasing in \vec{k} ; and the function BR_G is non-decreasing in \vec{k} and non-increasing in m .*

PROOF. We first show that that $\lambda_{m, \vec{k}}$ is monotone in m and \vec{k} . We then show that $p_u^{m, \vec{k}}$ is a monotone function of $\lambda_{m, \vec{k}}$ and that BR_G is a monotone function of $p_u^{m, \vec{k}}$, completing the proof.

We now show that $\lambda_{m, \vec{k}}$ is non-decreasing in m . Fix a vector of thresholds \vec{k} and let

$$g_{\vec{k}}(\lambda) = \sum_{t, i} i \frac{f_t \lambda^i q_{\vec{k}}(t, i)}{\sum_{j=0}^{k_t} \lambda^j q_{\vec{k}}(t, j)}, \quad (12)$$

where $q_{\vec{k}}$ is the value of q from Equation (8) (we add the subscript \vec{k} to stress the dependence on \vec{k}). The definition of $\lambda_{m, \vec{k}}$ in Equations (2) and (3) in Lemma 4.1 ensures that, for all m , $m = g_{\vec{k}}(\lambda_{m, \vec{k}})$. A relatively straightforward computation shows that $g'_{\vec{k}}(\lambda) > 0$ for all λ . Thus, if $m' > m$, $g_{\vec{k}}(\lambda) = m$, and $g_{\vec{k}}(\lambda') = m'$, we must have $\lambda' > \lambda$. It follows that $\lambda_{m, \vec{k}}$ is increasing in m . (Note that $\lambda_{m, \vec{k}}$ is undefined for $m \geq \sum_t f_t k_t$, which is why monotonicity holds only for values of m such that $mhn < \sum_t f_t k_t$.)

We next show that $\lambda_{m, \vec{k}}$ is non-increasing in \vec{k} . Since we have a finite set of types, it suffices to consider the case where a single type t^* increases its threshold by 1. Let \vec{k} denote the initial vector of thresholds, and let \vec{k}' denote the vector of thresholds after agents of type t^* increase their threshold by 1; that is, $k_t = k'_t$ for $t \neq t^*$, and $k'_{t^*} = k_{t^*} + 1$.

The first step in showing that $\lambda_{m, \vec{k}}$ is non-increasing in \vec{k} is to show that $g_{\vec{k}'}(\lambda_{m, \vec{k}}) > g_{\vec{k}}(\lambda_{m, \vec{k}}) = m$. We do this by breaking the sum in the definition of g in Equation (12) into two pieces; those terms were $t \neq t^*$, and those where $t = t^*$.

It follows immediately from Equation (8) that there exists a constant c such that, for all i and $t \neq t^*$, we have $q_{\vec{k}'}(t, i) = cq_{\vec{k}}(t, i)$. It follows from Equation (2) that for all i and $t \neq t^*$, since $k_t = k'_t$, we have

$$i \frac{f_t \lambda_{m, \vec{k}}^i q_{\vec{k}'}(t, i)}{\sum_{j=0}^{k'_t} \lambda_{m, \vec{k}}^j q_{\vec{k}'}(t, j)} = i \frac{f_t \lambda_{m, \vec{k}}^i c q_{\vec{k}}(t, i)}{\sum_{j=0}^{k'_t} \lambda_{m, \vec{k}}^j c q_{\vec{k}}(t, j)} = i \frac{f_t \lambda_{m, \vec{k}}^i q_{\vec{k}}(t, i)}{\sum_{j=0}^{k_t} \lambda_{m, \vec{k}}^j q_{\vec{k}}(t, j)}; \quad (13)$$

that is, the corresponding terms in the sum for $g_{\vec{k}'}(\lambda_{m, \vec{k}})$ and $g_{\vec{k}}(\lambda_{m, \vec{k}})$ are the same if $t \neq t^*$.

Now consider the corresponding terms for type t^* . First observe that for all

$i < k'_t$,

$$\frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}'}(t^*, i)}{\sum_{j=0}^{k'_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}'}(t^*, j)} < \frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}}(t^*, i)}{\sum_{j=0}^{k_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}}(t^*, j)}; \quad (14)$$

the two terms have essentially the same numerator (the use of $q_{\vec{k}'}$ instead of $q_{\vec{k}}$ cancels out as in Equation (13)), but the first has a larger denominator because $k'_{t^*} = k_{t^*} + 1$, so there is one more term in the sum. Since $f_{t^*} = \sum_{i=0}^{k_{t^*}} d_{q_{\vec{k}}}^*(t^*, i) = \sum_{i=0}^{k'_{t^*}} d_{q_{\vec{k}'}}^*(t^*, i)$, by Equations (2) and (3),

$$\sum_{i=0}^{k_{t^*}} \frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}}(t^*, i)}{\sum_{j=0}^{k_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}}(t^*, j)} = \sum_{i=0}^{k'_{t^*}} \frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}'}(t^*, i)}{\sum_{j=0}^{k'_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}'}(t^*, j)}. \quad (15)$$

It follows that

$$\sum_{i=0}^{k'_{t^*}} i \frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}'}(t, i)}{\sum_{j=0}^{k'_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}'}(t, j)} > \sum_{i=0}^{k_{t^*}} i \frac{f_{t^*} \lambda_{m, \vec{k}}^i q_{\vec{k}}(t, i)}{\sum_{j=0}^{k_{t^*}} \lambda_{m, \vec{k}}^j q_{\vec{k}}(t, j)}. \quad (16)$$

To see this, note that the two expressions above have the form $\sum_{i=0}^{k'_{t^*}+1} i c_i$ and $\sum_{i=0}^{k_{t^*}} i d_i$, respectively. By Equation (15), $\sum_{i=0}^{k'_{t^*}+1} c_i = \sum_{i=0}^{k_{t^*}} d_i = f_{t^*}$; by Equation (14), $c_i < d_i$ for $i = 0, \dots, k_{t^*}$. Thus, in going from the right side to the left side, weight is being transferred from lower terms to $k_{t^*} + 1$.

Combining Equations (13) and (16) gives us $g_{\vec{k}'}(\lambda_{m, \vec{k}}) > g_{\vec{k}}(\lambda_{m, \vec{k}}) = m$, as desired. Since $g_{\vec{k}'}(\lambda_{m, \vec{k}'}) = m$, by definition, it follows that $g_{\vec{k}'}(\lambda_{m, \vec{k}}) > g_{\vec{k}'}(\lambda_{m, \vec{k}'})$. Since, as shown above, $g_{\vec{k}'}$ is an increasing function, it follows that $\lambda_{m, \vec{k}} > \lambda_{m, \vec{k}'}$. Thus, $\lambda_{m, \vec{k}}$ is decreasing in \vec{k} .

We now show that the monotonicity of $\lambda_{m, \vec{k}}$ implies the monotonicity of $p_u^{m, \vec{k}}$. To do this, we show that, for all types t , $p_u^{m, \vec{k}} = p_d \lambda_{m, \vec{k}} \omega_t$. Since ω_t and p_d are independent of m and \vec{k} , it then follows that the monotonicity of $\lambda_{m, \vec{k}}$ implies the monotonicity of $p_u^{m, \vec{k}}$. (Recall that $\omega_t = \beta_t \chi_t / \rho_t$ was defined in Section 3.)

Fix a type t' . Then, dropping superscripts and subscripts on p_u , d , and λ for brevity, we have the following sequence of equalities (where the explanation for

some of these lines is given following the equations):

$$p_u = \left(\sum_t \rho_T(f_t - d(t, 0)) \right) \left(\frac{\chi_{t'} \beta_{t'}}{n \sum_t \chi_t \beta_t (f_t - d(t, k_t))} \right) \quad (17)$$

$$= \left(\frac{\sum_t \sum_{i=1}^{k_t} \rho_t d(t, i)}{\sum_t \sum_{i=0}^{k_t-1} \chi_t \beta_t d(t, i)} \right) \left(\frac{\chi_{t'} \beta_{t'}}{n} \right) \quad (18)$$

$$= \left(\frac{\sum_t \sum_{i=0}^{k_t-1} \rho_t \lambda \omega_t d(t, i)}{\sum_t \sum_{i=0}^{k_t-1} \chi_t \beta_t d(t, i)} \right) \left(\frac{\chi_{t'} \beta_{t'}}{n} \right) \quad (19)$$

$$= \lambda \left(\frac{\sum_t \sum_{i=0}^{k_t-1} \chi_t \beta_t d(t, i)}{\sum_t \sum_{i=0}^{k_t-1} \chi_t \beta_t d(t, i)} \right) (\omega_{t'} p_d) \quad (20)$$

$$= \lambda \omega_{t'} p_d$$

Equation (17) is just the definition of p_u from Equation (9). Equation (18) follows from the observation that, by Equation (2), $f_t = \sum_i d(t, i)$. Equation (19) follows from the observation that, again by Equation (2), $d(t, i) = \omega_t \lambda d(t, i-1)$. Equation (20) follows from the definitions of ω_t and p_d (see Equation (10)). Thus, as required, $p_u^{m, \vec{k}} = p_d \lambda_{m, \vec{k}} \omega_t$.

Finally, we show that the monotonicity of $p_u^{m, \vec{k}}$ implies the monotonicity of BR_G . Let $\vec{k}'' = BR_G(\vec{k})$. By Lemma 5.1, k_t'' is the maximum value of κ such that

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u^{m, \vec{k}}, p_d)}] \gamma_t.$$

We (implicitly) defined the random variable $J(\kappa, p_u, p_d)$ as a function on histories. Instead, we can define $J(\kappa, p_u, p_d)$ as a function on random bitstrings (which intuitively determine a history). With this redefinition, it is clear that, if $p_u < p'_u$, for all bitstrings b , we have $J(\kappa, p_u, p_d)(b) < J(\kappa, p'_u, p_d)(b)$. It easily follows that

$$E[(1 - (1 - \delta_t)/n)^{J(\kappa, p'_u, p_d)}] < E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u, p_d)}]$$

for all κ . Thus, the monotonicity of BR_G follows from the monotonicity of $p_u^{m, \vec{k}}$. ■

LEMMA 5.3. *For all games $G = (T, \vec{f}, h, m, n)$, there exists a $\delta^* < 1$ such that if $\delta_t > \delta^*$ for all t , there is a vector \vec{k} of thresholds such that $BR_G(\vec{k}) > \vec{k}$.*

PROOF. Take \vec{k} to be such that $k_t = \lceil m \rceil + 1$ for each type t . Then by Theorem 5.1, there exists a \vec{k}' such that $BR_G(\vec{k}) = \vec{k}'$. By Lemma 5.1, k_t' is the maximum value of κ such that

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u^{\vec{k}}, p_d)}] \gamma_t. \quad (4)$$

As δ_t approaches 1, $E[(1 - (1 - \delta_t)/n)^{J(\kappa, p_u^{\vec{k}}, p_d)}]$ approaches 1, and so the right hand side of Equation (4) approaches γ_t . For any standard agent, $\alpha_t < \gamma_t$. Thus, there exists a δ_t such that

$$\alpha_t \leq E[(1 - (1 - \delta_t)/n)^{J(k_t, p_u^{\vec{k}}, p_d)}] \gamma_t.$$

For this choice of δ_t , we must have $k_t' \geq k_t + 1 > k_t$. Take $\delta^* = \max_t \delta_t$. ■

C. PROOFS FROM SECTION 8

THEOREM 8.3. *Fix a game G and vector of thresholds \vec{k} . Let $R_{\vec{k},t} = p_u^{\vec{k},t}/p_d^t$. In the limit as the number of rounds goes to infinity, the fraction of the agent's requests that have an agent willing and able to satisfy them that get satisfied is $(R_{\vec{k},t} - R_{\vec{k},t}^{k_t+1})/(1 - R_{\vec{k},t}^{k_t+1})$ if $R_{\vec{k},t} \neq 1$ and $k_t/(k_t + 1)$ if $R_{\vec{k},t} = 1$.*

PROOF. Consider the Markov chain \mathcal{M} that results from fixing the agent's policy to s_{k_t} in $\mathcal{P}_{G,\vec{s}(\vec{k}),t}$. \mathcal{M} satisfies the requirements given in Theorem A.1 to have a limit distribution. It can be easily verified that the distribution gives the agent probability $R^i(1 - R)/(1 - R^{k+1})$ of having i dollars if $R \neq 1$ and probability $1/(k + 1)$ if $R = 1$ satisfies the detailed balance condition and thus is the limit distribution. This gives the probabilities given in the theorem. ■

THEOREM 8.4. *Suppose that t and s are two types that agree except for the value of χ , and that $\chi_t < \chi_s$. If $\vec{k} = (k_t, k_s)$ is an ε -Nash equilibrium for $G = (\{t, s\}, \vec{f}, h, m, n)$ with social welfare w , then there exist m' , and n' such that $\vec{k}' = (k_s)$ is an ε -Nash equilibrium for $G'_{m',n'} = (\{t\}, \{1\}, h, m', n')$ with social welfare greater than w .*

PROOF. We prove the theorem by finding m' , and n' such that agents in $G'_{m'}$ that play some strategy k get essentially the same utility that an agent with sybils would by playing that strategy in G . Since k_s was the optimal strategy for agents with sybils in G , it must be optimal in $G'_{m',n'}$ as well. Since agents with sybils have utility at least as great as those without, social welfare will be at least as large in $G'_{m',n'}$ as in G .

Since an agent can earn a dollar only if he is able to satisfy the current request, $0 < p_u^{m,\vec{k},s} < \beta_s$. The constraint that $hm'n'$, the total amount of money, is a natural number means that m' must be a rational number. For the moment, we ignore that constraint and allow m' to take on any value in $[0, k'_t]$. From Equation (9), $p_u^{m',\vec{k}',t}$ is continuous in $d_{q_{\vec{k}'}}^*$, which, by Lemma 4.1, is continuous in $\lambda_{m',\vec{k}'}$ and thus m' . We use this continuity to show that we can find a value of m' such that $p_u^{m,\vec{k},s} = p_u^{m',\vec{k}',t}$. By Equation (3), if $m' = 0$ then $d_{q_{\vec{k}'},m}^*(t,0) = 1$, and if $m' = k'_t$ then $d_{q_{\vec{k}'},m}^*(t,k'_t) = 1$. Combining these with Equation (9) gives $p_u^{0,\vec{k}',t} = 0$ and $p_u^{m',\vec{k}',t} = \beta_t$. Thus, by the Intermediate Value Theorem, there exists an m' such that $p_u^{m,\vec{k},s} = p_u^{m',\vec{k}',t}$. For this choice of m' , observe that by Lemma 5.1, $\mathcal{P}_{G,\vec{s}(\vec{k}),s}$ and $\mathcal{P}_{G_{m'},\vec{s}(\vec{k}'),t}$ have the same optimal threshold policy.

If m' is rational, say $m' = a/b$, take $n' = bn$; then $hm'n'$ is an integer and, by the argument above, \vec{k}' is an equilibrium for $G'_{m',n'}$. Since $p_u^{m',\vec{k}'} = p_u^{\vec{k},s} > p_u^{\vec{k},t}$, we must have $\zeta_{G_{m',n'}} < \zeta_G$. Thus, social welfare has increased. If m' is not rational, we instead use a rational value m'' sufficiently close to m' that \vec{k}' is still an equilibrium for $G'_{m'',n'}$ and $\zeta_{G_{m'',n'}} < \zeta_G$. ■

D. PROOF OF LEMMA 9.2

LEMMA 9.2. *If d is a fully-supported distribution of money with finite support, there exist an infinite number of explanations of d .*

PROOF. Fix λ . The distribution d and λ determine an explanation \vec{f} as follows. By Equation (7), we need \vec{f} to satisfy $d(j) \sum_{i=0}^{K_d} d_\lambda(i, j)$.

Recall that K_d is the maximum value for which $d(K_d) > 0$. Start by considering f_{K_d} . By the definition of d_λ , $d_\lambda(i, j) = 0$ if $j > i$. Thus, the constraint becomes

$$d(K_d) = d_\lambda(K_d, K_d) = f_{K_d} \lambda^{K_d} / \left(\sum_{l=0}^{K_d} \lambda^l \right).$$

Take f_{K_d} to be the unique value that satisfies this equation. Once we have defined f_{K_d} , again apply the constraint and take f_{K_d-1} to be the unique value that satisfies

$$d(K_d-1) = d_\lambda(K_d, K_d-1) + d_\lambda(K_d-1, K_d-1) = f_{K_d} \lambda^{K_d-1} / \left(\sum_{l=0}^{K_d} \lambda^l \right) + f_{K_d-1} \lambda^{K_d-1} / \left(\sum_{l=0}^{K_d-1} \lambda^l \right).$$

Iterating this process uniquely defines \vec{f} as the unique value that satisfies

$$d(j) = \sum_{i=j}^{K_d} d_\lambda(i, j) = \sum_{i=j}^{K_d} f_i \lambda^j / \left(\sum_{l=0}^i \lambda^l \right),$$

or

$$f_i = \left(\sum_{l=0}^i \lambda^l \right) / \lambda^i \left(d(i) - \sum_{j=i+1}^{K_d} f_j \lambda^j / \left(\sum_{l=0}^j \lambda^l \right) \right).$$

However, \vec{f} may not be an explanation, since some f_j may be negative. This happens exactly when

$$d(i) < \sum_{j=i+1}^{K_d} f_j \lambda^j / \left(\sum_{l=0}^j \lambda^l \right). \quad (21)$$

As λ grows large, the right-hand side of (21) tends to 0. Since d is fully-supported, we must have $d(i) > 0$. Thus, we can ensure that (21) does not hold for any i by taking λ sufficiently large. Thus, for sufficiently large λ , \vec{f} provides an explanation for d . Continuing to increase λ will give an infinite number of different explanations. ■

REFERENCES

- ABRAHAM, I., DOLEV, D., GONEN, R., AND HALPERN, J. 2006. Distributed computing meets game theory: Robust mechanisms for rational secret sharing and multiparty computation. In *Proc. 25th ACM Symp. on Principles of Distributed Computing (PODC)*. 53–62.
- ADAR, E. AND HUBERMAN, B. A. 2000. Free riding on Gnutella. *First Monday* 5, 10.
- ANAGNOSTAKIS, K. G. AND GREENWALD, M. 2004. Exchange-based incentive mechanisms for peer-to-peer file sharing. In *International Conference on Distributed Computing Systems (ICDCS)*. 524–533.
- APERJIS, C. AND JOHARI, R. 2006. A peer-to-peer system as an exchange economy. In *GameNets '06: Proceeding from the 2006 Workshop on Game Theory for Communications and Networks*. 10.
- AUMANN, R. J. 1959. Acceptable points in general cooperative n -person games. In *Contributions to the Theory of Games IV, Annals of Mathematical Studies 40*, A. Tucker and R. Luce, Eds. Princeton University Press, Princeton, N. J., 287–324.

- AUYOUNG, A., CHUN, B., NG, C., PARKES, D., VAHDAT, A., AND SNOEREN, A. 2007. Practical market-based resource allocation. Tech. rep., UC San Diego. CS2007-0901.
- BELENKIY, M., CHASE, M., ERWAY, C. C., JANNOTTI, J., KÜPÇÜ, A., LYSYANSKAYA, A., AND RACHLIN, E. 2007. Making p2p accountable without losing privacy. In *Proceedings of the 2007 ACM Workshop on Privacy in the Electronic Society, WPES 2007, Alexandria, VA, USA, October 29, 2007*. 31–40.
- BRUNELLE, J., HURST, P., HUTH, J., KANG, L., NG, C., PARKES, D., SELTZER, M., SHANK, J., AND YOUSSEF, S. 2006. Egg: An extensible and economics-inspired open grid computing platform. In *Third Workshop on Grid Economics and Business Models (GECON)*. 140–150.
- CHUN, B., BUONADONNA, P., AUYOUNG, A., NG, C., PARKES, D., SCHNEIDMAN, J., SNOEREN, A., AND VAHDAT, A. 2005. Mirage: A microeconomic resource allocation system for sensor network testbeds. In *Second IEEE Workshop on Embedded Networked Sensors*. 19–28.
- COHEN, B. 2003. Incentives build robustness in BitTorrent. In *First Workshop on the Economics of Peer-to-Peer Systems (P2PECON)*.
- COVER, T. AND THOMAS, J. 1991. *Elements of Information Theory*. John Wiley & Sons, Inc., New York.
- DOUCEUR, J. R. 2002. The sybil attack. In *First International Workshop on Peer-to-Peer Systems (IPTPS)*. 251–260.
- ELLISON, G. 1994. Cooperation in the prisoner’s dilemma with anonymous random matching. *Review of Economic Studies* 61, 567–588.
- FELDMAN, M., LAI, K., STOICA, I., AND CHUANG, J. 2004. Robust incentive techniques for peer-to-peer networks. In *ACM Conference on Electronic Commerce (EC)*. 102–111.
- FRIEDMAN, E. J. AND RESNICK, P. 2001. The social cost of cheap pseudonyms. *Journal of Economics and Management Strategy* 10, 2, 173–199.
- FUDENBERG, D. AND TIROLE, J. 1991. *Game Theory*. MIT Press.
- GROVE, A. J., HALPERN, J. Y., AND KÖLLER, D. 1994. Random worlds and maximum entropy. *J. Artif. Intell. Res. (JAIR)* 2, 33–88.
- GUHA, R., KUMAR, R., RAGHAVAN, P., AND TOMKINS, A. 2004. Propagation of trust and distrust. In *Conference on the World Wide Web (WWW)*. 403–412.
- GUPTA, M., JUDGE, P., AND AMMAR, M. H. 2003. A reputation system for peer-to-peer networks. In *Network and Operating System Support for Digital Audio and Video (NOSSDAV)*. 144–152.
- HAUERT, C., TRAUlsen, A., BRANDT, H., NOWAK, M. A., AND SIGMUND, K. 2007. Via freedom to coercion: The emergence of costly punishment. *Science* 316, 1905–1907.
- HAYRAPETYAN, A., TARDOS, É., AND WEXLER, T. 2006. The effect of collusion in congestion games. In *Proceedings of the 38th Annual ACM Symposium on Theory of Computing, Seattle, WA, USA, May 21-23, 2006*. 89–98.
- HENS, T., SCHENK-HOPPE, K. R., AND VOGT, B. 2007. The great Capitol Hill baby sitting co-op: Anecdote or evidence for the optimum quantity of money? *J. of Money, Credit and Banking* 9, 6, 1305–1333.
- HUGHES, D., COULSON, G., AND WALKERDINE, J. 2005. Free riding on Gnutella revisited: The bell tolls? *IEEE Distributed Systems Online* 6, 6.
- IOANNIDIS, J., IOANNIDIS, S., KEROMYTIS, A. D., AND PREVELAKIS, V. 2002. Fileteller: Paying and getting paid for file storage. In *Financial Cryptography*. 282–299.
- ITHACA HOURS INC. 2005. Ithaca hours. <http://www.ithacahours.org/>.
- JAYNES, E. T. 1978. Where do we stand on maximum entropy? In *The Maximum Entropy Formalism*, R. D. Levine and M. Tribus, Eds. MIT Press, Cambridge, Mass., 15–118.
- KAMVAR, S. D., SCHLOSSER, M. T., AND GARCIA-MOLINA, H. 2003. The Eigentrust algorithm for reputation management in P2P networks. In *Conference on the World Wide Web (WWW)*. 640–651.
- KANDORI, M. 1992. Social norms and community enforcement. *Review of Economic Studies* 59, 63–80.
- KRUGMAN, P. 1999. *The Accidental Theorist*. Norton.

- LOVASZ, L. AND WINKLER, P. 1995. Mixing of random walks and other diffusions on a graph. In *Surveys in Combinatorics, 1993, Walker (Ed.), London Mathematical Society Lecture Note Series 187, Cambridge University Press*.
- MAINLAND, G., KANG, L., LAHAIE, S., PARKES, D. C., AND WELSH, M. 2004. Using virtual markets to program global behavior in sensor networks. In *Proceedings of the 11st ACM SIGOPS European Workshop, Leuven, Belgium, September 19-22, 2004*. 1.
- MAS-COLELL, A., WHINSTON, M. D., AND GREEN, J. R. 1995. *Microeconomic Theory*. Oxford University Press, Oxford, U.K.
- MILGROM, P. AND ROBERTS, J. 1900. Rationalizability, learning, and equilibrium in games with strategic complement- arities. *Econometrica* 58, 6, 1255–1277.
- MILLER, M. S. AND DREXLER, K. E. 1988. Markets and computation: Agoric open systems. In *The Ecology of Computation*, B. Huberman, Ed. Elsevier, 133–175.
- NG, C., BUONADONNA, P., CHUN, B., SNOEREN, A., AND VAHDAT, A. 2005. Addressing strategic behavior in a deployed microeconomic resource allocator. In *Third Workshop on Economics of Peer-to-Peer Systems (P2PECON)*. 99–104.
- PETERSON, R. S. AND SIRER, E. G. 2009. Antfarm: Efficient content distribution with managed swarms. In *Proceedings of Networked Systems Design and Implementation (NSI), Boston, Massachusetts, April 2009*.
- PUTERMAN, M. L. 1994. *Markov Decision Processes*. Wiley.
- REEVES, D. M., SOULE, B. M., AND KASTURI, T. Group decision making with Yootles. <http://ai.eecs.umich.edu/people/dreeves/yootles/yootles.pdf>.
- RESNICK, S. I. 1992. *Adventures in Stochastic Processes*. Birkhauser.
- ROUGHGARDEN, T. AND TARDOS, É. 2002. How bad is selfish routing? *J. ACM* 49, 2, 236–259.
- SIRIVIANOS, M., PARK, J. H., YANG, X., AND JARECKI, S. 2007. Dandelion: Cooperative content distribution with robust incentives. In *Proceedings of the 2007 USENIX Annual Technical Conference, June 17-22, 2007, Santa Clara, CA, USA*. 157–170.
- STONEBRAKER, M., AOKI, P. M., LITWIN, W., PFEFFER, A., SAH, A., SIDELL, J., STELIN, C., AND YU, A. 1996. Mariposa: a wide-area distributed database system. *The VLDB Journal* 5, 1, 48–63.
- SWEENEY, J. AND SWEENEY, R. J. 1977. Monetary theory and the great capitol hill babysitting co-op crisis: Comment. *J. of Money, Credit and Banking* 9, 1, 86–89.
- TARSKI, A. 1955. A lattice-theoretical fixpoint theorem and its applications. *Pacific Journal of Mathematics* 5, 285–309.
- TIMBERLAKE, R. H. 1987. Private production of scrip-money in the isolated community. *J. of Money, Credit and Banking* 19, 4, 437–447.
- TOPKIS, D. M. 1979. Equilibrium points in nonzero-sum n-person submodular games. *Siam Journal of Control and Optimization* 17, 773–787.
- VISHNUMURTHY, V., CHANDRAKUMAR, S., AND SIRER, E. G. 2003. KARMA: a secure economic framework for peer-to-peer resource sharing. In *First Workshop on Economics of Peer-to-Peer Systems (P2PECON)*.
- WALSH, K. AND SIRER, E. G. 2006. Experience with an object reputation system for peer-to-peer filesharing. In *Third Symp. on Network Systems Design & Implementation (NSDI)*. 1–14.
- WASHINGTON METROPOLITAN AREA TRANSIT COMMISSION. 1970. Scrip system of the D.C. transit system.
- XIONG, L. AND LIU, L. 2002. Building trust in decentralized peer-to-peer electronic communities. In *International Conference on Electronic Commerce Research (ICECR)*.
- YOKOO, M., SAKURAI, Y., AND MATSUBARA, S. 2004. The effect of false-name bids in combinatorial auctions: new fraud in internet auctions. *Games and Economic Behavior* 46, 1, 174–188.
- ZHAO, S., LO, V. M., AND GAUTHIERDICKEY, C. 2005. Result verification and trust-based scheduling in peer-to-peer grids. In *Fifth IEEE International Conference on Peer-to-Peer Computing (P2P 2005), 31 August - 2 September 2005, Konstanz, Germany*. 31–38.